



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:


- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>







Per. 19282 e $\frac{+7}{-11}$

THE
ROTHAMSTED MEMOIRS
ON
AGRICULTURAL CHEMISTRY
AND
PHYSIOLOGY.

BY
SIR JOHN BENNET LAWES, BART., D.C.L., LL.D., F.R.S., F.C.S., &c.,
OF ROTHAMSTED, HERTS.,
AND
SIR JOSEPH HENRY GILBERT, M.A., PH.D., LL.D., F.R.S., V.P.C.S., &c.

VOLUME IV.
CONTAINING REPORTS OF EXPERIMENTS,
ON THE FEEDING OF ANIMALS,
SEWAGE UTILISATION,
ENSILAGE,
&c., &c.

PUBLISHED 1864—1885 INCLUSIVE.

LONDON:
PRINTED BY WILLIAM CLOWES AND SONS, LIMITED,
STAMFORD STREET AND CHARING CROSS.
1893.



ROTHAMSTED MEMOIRS.

CONTENTS OF VOL. IV.

REPORTS OF EXPERIMENTS ON THE FEEDING OF ANIMALS, SEWAGE UTILISATION, ENSILAGE, &c., &c.

	Published
1. On the Chemistry of the Feeding of Animals for the production of Meat and Manure (read before the Royal Dublin Society, March 31, 1864)	1864
2. On the Sewage of Towns (Third Report, and Appendices 1, 2, & 3, of the Royal Commission. Presented to Parliament)	1865
3. Report (Presented to Parliament) of Experiments undertaken by Order of the Board of Trade to determine the relative Values of Unmalted and Malted Barley as Food for Stock	1866
4. On the Composition, Value, and Utilisation of Town Sewage (Jour. Chem. Soc., New Series, vol. iv., Entire Series, vol. xix., p. 80.)	1866
5. Food in its relations to the various exigencies of the Animal Body (Phil. Mag., July, 1866)	1866
6. On the Sources of the Fat of the Animal Body (Abstract—Rep. Brit. Ass. for 1866)	1866
7. On the Sources of the Fat of the Animal Body (Phil. Mag., December, 1866)	1866
8. Note—On Sewage Utilisation (Proceedings of the Institution of Civil Engineers, vol. xiv., part 3.)	1876
9. On some points in connection with Animal Nutrition (South Kensington Science Conferences, 1876)	1876
10. On the Formation of Fat in the Animal Body (Journal of Anatomy and Physiology, vol. xi., part 4.)	1877
11. Supplement to former Paper entitled :—"Experimental Inquiry into the Composition of some of the Animals fed and slaughtered as Human Food." <i>Composition of the Ash of the Entire Animals, and of certain Separated Parts. Abstract</i> (Proc. Roy. Soc. No. 226, 1883)	1883
12. Experiments on Ensilage conducted at Rothamsted, Season 1884-5	1885

THE CHEMISTRY OF THE FEEDING OF ANIMALS FOR THE PRODUCTION OF MEAT AND MANURE.

LECTURE DELIVERED BEFORE THE ROYAL DUBLIN SOCIETY, MARCH 31, 1864.

BY JOHN BENNET LAWES, F.R.S., F.C.S.

ing and feeding of stock must always be an important branch of the agricultural science of this Island. With a climate rarely so hot in summer, or so cold in winter, as to materially affect the vegetation, Ireland may not less truly be styled the Emerald Isle. A succession of seasons more than usually unfavourable for agriculture has greatly reduced the profits, and even the industry of your farmers. It is natural, therefore, that there should be, at the present time, more attention directed to the production of meat, and less of corn; more especially as with the dearth of grain that of meat has considerably increased and has probably not yet reached its maximum.

The application of science to agriculture has long been regarded with much favour by practical agriculturists; there are still very many who feel how important it would be to know more of the principles of the various operations than they do at present. The principles involved even in old-established modes of agriculture are frequently but little understood; and it is every year becoming less and less an intuitive business than it was formerly: new manures, improved descriptions of seeds, and new mechanical appliances are continually being introduced, requiring more knowledge and discrimination in their selection and use. The principal branch of agriculture upon which I propose to address you this evening is that which relates to the production of meat and manure.

When fattening animals are supplied with an amount of proper food they increase in weight, and the portion of the food being fixed or stored up in the animal is rejected by the animal in solid form, and serve as manure; the portion which is expended or lost in the processes of respiration and cutaneous exhalation. Experience has shown that some foods have higher feeding value than others, and it is generally supposed that the knowledge of the feeding properties of foods will also be of great value in the estimation of the value of the manure.

It is the province of agricultural chemistry to determine what proportion of the several constituents of the food consumed will be stored up in the form of meat, and how much will remain as manure, according to the description of animal, and the kind of food employed; and so to provide the means of estimating the value of the respective products of the feeding operation. To this end, it is necessary to determine, by means of careful analysis, the composition of the foods consumed, of animals in the store or lean and in the fat condition, and of the manurial matters voided. Such an undertaking is, however, by no means a light one, and it can only be carried out with any prospect of success by the conjoint aid of experiments on a large scale in the feeding-shed, and of investigations in the laboratory, involving a great amount of analytical labour, and requiring the observance of all the refinements of method which modern science permits.

I propose to bring before you a condensed summary of some of the results which have been obtained in experiments made at different times during the last twenty years, at my farm and laboratory, at Rothamsted, in Hertfordshire. There are, it is true, many points which are not as yet satisfactorily cleared up, and some of these are still under investigation. The figures given in the tables, in most cases, however, represent the results obtained in careful experiments with large numbers of animals of each of the descriptions indicated, and they may be taken as showing what should be the average result obtained in ordinary farm practice, when animals of fair quality are fed liberally for the butcher.

Composition of Oxen, Sheep, and Pigs, in the Store and Fat Condition.

For the purposes of my illustration, I shall assume that an ox or a sheep will increase in weight by about one-half, and that a pig will double its weight during the so-called fattening period. Accordingly, I shall direct your attention to the composition of each of these descriptions of animal when in the lean or store condition.

ity, or stinted in amount.
 iminary remarks, I will now direct
 the tables.

ition, per cent., of Oxen, Sheep, and
 store, and in the fat condition.

	Oxen.		Sheep.		Pigs.	
	Store	Fat.	Store	Fat.	Store	Fat.
e	18.0	16.0	18.0	12.5	14.0	10.5
stance	16.0	30.0	18.0	33.0	22.0	44.0
.....	5.2	4.0	3.5	3.0	2.8	1.8
ce	39.2	49.0	36.5	48.5	38.8	56.3
.....	60.8	51.0	63.5	51.5	61.2	43.7
.....	100.0	100.0	100.0	100.0	100.0	100.0

the composition, per cent., of oxen,
 th in the store and in the fat con-
 nents given being the nitrogenous
 the non-nitrogenous substance or
 incombustible matter, the sum of
 substance, and the water.
 nitrogenous substance, it is seen
 ription of animal there are seven
 of it in the fat than in the store
 on the other hand, there is in the
 oxen and the sheep nearly, and in
 , twice as much in 100 lbs. live
 as in the same weight of the store

is generally attributable to the character of t
 and is found to result when too much oily r
 given, or when pigs are fed freely with roots
 succulent food.

*Proportion of Parts, in Animals of different
 tions, and in different Conditions of Matu*

Passing from the question of the chemical
 sition of oxen, sheep, and pigs, it will be d
 before considering the relation of the incre
 manure produced to that of the food co
 briefly to point out some characteristic differ
 structure or relative proportion of certain
 internal organs, as in these will be found the k
 difference in the character and amount of foc
 the three descriptions of animal respectively
 Table II. illustrates this part of the subject.

TABLE II.—*Relation of Parts in Animals of
 Descriptions, and in different Conditions of Ma*

	Per Cent.				
	In different animals.			In Sheep	
	Oxen.	Sheep.	Pigs.	Store.	fat
Average of	16	249	59	5	1
Stomachs and contents	11.8	7.5	1.2	0.1	0.1

sheep contains only $7\frac{1}{2}$, and that of the ox. Of intestines and contents, on the other hand, the ox contains only $2\frac{3}{4}$, the sheep $3\frac{1}{2}$, and the pig 11 per cent. Again, of stomachs and intestines (with their respective contents) taken together, the ox contains about $14\frac{1}{2}$, sheep about 11, and the pig 11 per cent. Thus, of the receptacles and contents of the food, the oxen contain the largest, and pigs by far the smallest proportion. It would appear to indicate a great difference in requirement for bulk of food, such, indeed, as actually exists. Oxen require a larger proportion of body fibre in their food than sheep, and more than pigs. On the other hand, the pig contains much more starch, or allied matter, than that of sheep, and that of sheep is more than that of oxen, reckoned in relation to the weight of the animal; and it is known that starch is the primary change (into sugar) almost throughout the length of the intestinal canal. Accordingly, that the pig has a larger proportion of food than the sheep, and the sheep more than

the ox, is further elaborating, or what we may call labour organs of the body, and their contents, liver, lungs, blood, &c.—the proportion to be nearly the same in the three descriptions of animal.

The proportion of internal or loose fat is greater than in the oxen; but it should be observed that a large proportion of the sheep contribute to the average result given in the table were in an advanced state of fatness than the oxen. The relatively small proportion of internal fat is accounted for by the peculiarities of the constitution of its internal organs is very small, and its speciality is to lay on fat in a proportion outside the frame.

The proportion of the table shows the varying amounts of the different parts in one and the same animal, according to its degree of maturity. The animals selected for illustration of this are the sheep. Records not given in the table are those of the animals grew and fattened, the proportion, per head, of stomachs and contents is considerably; that the intestines and contents are a much less degree; that the internal

loose fat was more than trebled; and that the other internal parts, and their fluids, collectively, increased in nearly the same proportion as the stomachs and contents. The general result was, that the total offal parts increased in actual amount from the store to the very fat condition in the proportion of about 1 to $1\frac{1}{2}$; but the total carcass parts augmented from 1 to nearly $2\frac{1}{2}$ —much more, therefore, than the total offal parts.

Turning now to the figures in the table, it is seen that the per cent., or proportion in 100 parts, of all the internal organs and parts, excepting the loose fat, diminished very considerably as the animals matured and fattened. Whilst the total offal parts diminished from 45.2 in the store, to 40.6 in the fat, and to 35.5 per cent. in the very fat condition, the carcass parts increased from 53.4 in the store, to 58.7 in the fat, and to 64.1 per cent. in the very fat condition. That is to say, the so-called offal parts, which are chiefly composed of the organs of reception, elaboration, and transmission of the food constituents, increase in very much less proportion than those parts which it is the object of the feeder should be produced from the food consumed.

Relation of the Increase, Manure, and Loss by Respiration, to the Food consumed, by different Animals.

We now come to the question of the description and amount of food consumed by the different animals to produce a given amount of increase, and to the collateral questions of the relation of the constituents in the increase and in the manure to those in the food consumed.

Table III. shows the amounts of certain foods assumed to be required for the production of 100 lbs. of increase in live weight—of oxen, sheep, and pigs, respectively. The amounts will, of course, vary, according to the quality of the animal, the stage of its development, the external conditions to which it is subjected, the description and quality of the food, and so on; but the quantities assumed are approximately those which will be required, taking the average of large numbers of animals over the whole period of fattening, and supposing foods of the descriptions indicated, and of good quality, are employed, and that other conditions are moderately favourable.

TABLE III.

E III.—Food, Increase, Manure, &c., of Fattening Animals.

OXEN.

250 lbs. Oil-cake 600 lbs. Clover Chaff 3,500 lbs. Swedes and Supply—				100 Total Dry Substance of Food supply—			Amount of each con- stituent stored up, for 100 of it con- sumed.
In Food.	In 100 lbs. Increase.	In Manure.	To Respi- ration, &c.	In Increase.	In Manure.	To Respi- ration, &c.	
lbs.	lbs.	lbs.	lbs.				
218	9.0	} 323.0	636	} 0.8 5.2	} 29.1	} 57.3	} 4.1 7.2
808	58.0		—				
83	1.6	81.4	—	0.2	7.4	—	1.9
1109	68.6	404.4	636	6.2	36.5	57.3	

SHEEP.

250 lbs. Oil-cake 300 lbs. Clover Chaff 4,000 lbs. Swedes and Supply—				100 Total Dry Substance of Food Supply—			Amount of each con- stituent stored up, for 100 of it con- sumed.
In Food.	In 100 lbs. Increase.	In Manure.	To Respi- ration, &c.	In Increase.	In Manure.	To Respi- ration, &c.	
lbs.	lbs.	lbs.	lbs.				
177	7.5	} 229	548.5	} 0.8 7.0	} 25.1	} 60.1	} 4.2 9.4
671	63.0		—				
61	2.0	62	—	0.2	6.8	—	8.1
912	72.5	291	548.5	8.0	31.9	60.1	

PIGS.

500 lbs. Barley meal produce 100 lbs. Increase, and Supply—				100 Total Dry Substance of Food Supply—			Amount of each constituent stored up, for 100 of it consumed.
In Food.	In 100 lbs. Increase.	In Manure.	To Respi- ration, &c.	In Increase.	In Manure.	To Respi- ration, &c.	
lbs.	lbs.	lbs.	lbs.				
52	7.0	} 59.8	276.2	} 1.7 15.7	} 14.3	} 65.7	} 13.5 18.5
367	66.0		—				
11	0.8	10.2	—	0.2	2.4	—	7.3
420	73.8	70.0	276.2	17.6	16.7	65.7	

ferent foods recorded in
after a very careful con-
sumed numerous experiments on
and after the illustrations
different proportions of
descriptions of animal, it
are the variations in the
food recorded as required
lbs, to produce the same

amount of increase, oxen consume a much larger
proportion of hay, containing so much indigestible
matter, than sheep; whilst pigs are fattened on a diet
as concentrated and containing as little indigestible
substances as corn alone. The actual amounts of food
assumed to be required for the production of 100 lbs.
increase in live weight are—for oxen, 250 lbs. of oil-
cake, 600 lbs. of hay-chaff, and 3,500 lbs. of swedes;
for sheep, 250 lbs. of oil-cake, 300 lbs. of hay-chaff

and 4,000 lbs. of swedes; and for pigs, 500 lbs. of barley meal.

It will be remembered that when speaking of the composition of the animals themselves, their constituents were grouped under the heads of nitrogenous substance, non-nitrogenous substance, mineral matter, and total dry substance, and the same classification is, for convenience of comparison, adopted in reference to the composition of the food, increase, and manure, of the different animals as recorded in Table III.

As the food of the pig is the most simple, I will direct your attention to the figures relating to it in the first place. These will be found in the lowest division of the table.

The 500 lbs. of barley meal consumed in increasing the weight of the pig from 100 to 200 lbs. contains 420 lbs. of dry substance, and the 100 lbs. increase in live weight produced by it not quite 74 lbs.; about 70 lbs. remain in the manure, and 276 out of the 420 lbs. consumed are expended in respiration, and other exhalations from the body. Nearly two-thirds of the whole dry substance consumed have, therefore, been expended in keeping in working order the living meat and manure-making machine.

Looking to the column showing the composition of the 100 lbs. of increase, it is seen that it contains only 7 lbs. of nitrogenous substance, and 66 lbs., or more than 9 times as much non-nitrogenous substance or fat, whilst the mineral matter does not amount to 1 per cent. The general result is, then, that nearly two-thirds of the fattening increase in live weight were pure fat itself, and only about one-fourteenth of it nitrogenous substance or lean meat.

But to produce the 7 lbs. of nitrogenous substance in increase, 52 lbs. were consumed in food; by far the greater part of the remainder being found in the manure. To produce the 66 lbs. of fat, 357 lbs. of non-nitrogenous substance were consumed; but as it existed in the food almost entirely in the form of starch, and as it requires about $2\frac{1}{2}$ parts of starch to form 1 of fat, it may be said that at least 165 lbs. of the non-nitrogenous substance consumed contributed pretty directly to the formation of the 66 lbs. of fat. Lastly in reference to the increase: of the 11 lbs. of mineral matter consumed, only about $\frac{3}{4}$ lb. were stored up in the increase of the animal.

It is observed, then, that a comparatively small proportion of either the nitrogenous substance, or the mineral matter of the food, is retained in the increase; the manure, on the other hand, retains a very large proportion of the former, and nearly the whole of the latter.

Of 100 parts of gross dry substance consumed, 1.7 parts of nitrogenous substance, 15.7 of fat, and 0.2 of mineral matter—in all 17.6 parts—are stored up in the increase; 14.3 parts, consisting of highly nitro-

genous organic matter, and 2.4 parts of mineral matter, making a total of 16.7 parts, are in the manure; and 65.7 parts, consisting chiefly of carbon, hydrogen, and oxygen, are lost by respiration &c. Or, if we reckon the proportion of each constituent consumed, which is stored up in the increase, the last column of the table shows that of nitrogenous substance consumed, 13.5 parts; of non-nitrogenous substance consumed, 18.5 parts; of 100 mineral matter consumed, 7.3 parts retained in the increase.

It will not be necessary to follow so far the figures in the table relating to the sheep and oxen. It will suffice to direct attention to the differences of result obtained with the three different animals.

Whilst the pig required only 420 lbs., the sheep required 912 lbs., and the oxen 1,109 lbs. of dry substance in food to produce 100 lbs. increase in live weight. In other words, the sheep consumed more than twice as much, and the oxen more than two and a half times as much, to produce a given amount of increase in the pig. But the food of the pig was of a much simpler character than that of the other animals. The food consisted entirely of highly elaborated grain, resembling human food, the food of the sheep contained a large amount both of woody fibre and crude succulent roots; that of the ox contained the largest proportion of hay, with its high percentage of indigestible woody matter.

Turning to the columns giving the composition of 100 parts of the increase, they show that the food of the pig contained 73.8 parts of dry substance, that of the sheep contained rather less, and that of the oxen rather less still. The proportion of dry substance was greater in the increase of the pig than in that of the sheep, and greater in that of the sheep than in that of the oxen. The contrary was, however, the case with the proportion of nitrogenous substance, which was the greatest (9 per cent.) in that of the oxen, less (7.5 per cent.) in that of the sheep, and less still (7 per cent.) in that of the pig. As to the percentage of mineral matter in the increase of the ox and sheep, it was considerably higher than in that of the pig, even rather higher in the case of sheep than in that of the ox. Independently of any essential differences in the different animals, this result is due to the fact that sheep and oxen, especially the ox, develop bony structure during the fattening more than pigs. It is true that both sheep and oxen, compared with oxen, fattened at an early stage of their development; but not only are they more naturally disposed to fatten instead of frame very early in his career, if only supplied with proper food, but the practical

The last column of the table shows that nitrogenous substance of food consumed, and sheep stored up little more than 4, about 13.5 parts; that for 100 non-nitrogenous substance in food, the oxen yielded 7.2, 3.4, and the pigs 18.5 parts of fat in food that for 100 mineral matter consumed oxen stored up 1.9, the sheep 3.1, and the pigs 16.7 parts.

The very much larger proportion of the consumed food of the pig than of that of oxen could be stored up as increase is, however, what we should expect, when we consider that the food of the pig consists of matured grain, and the food of the ox consists of comparatively immatured vegetable matter containing a large proportion of indigestible matter, and also a larger amount of nitrogenous matter in proportion to its digestible non-nitrogenous constituents.

Thus the pig, with his much higher character of food, should increase so much more than the sheep. The amount consumed, and the sheep more so, the ox returned as manure 36.5 per cent of dry substance he consumed, the sheep 16.7 per cent. The proportion of the consumed matter that was lost by respiration, on the other hand, rather the lowest, namely, 57.3 per cent.; whilst with the ox it was 60.1, and with the pig it was 65.7 per

The first column of this table shows that with the pig increases from 6 to $6\frac{1}{2}$ per cent. of its weight in a week, the sheep increases only $1\frac{1}{2}$, and the ox more than 1 per cent. No wonder, then (taking into account the difference in the character of the food) that the oxen and sheep, requiring so much more time to add a given proportion to the weight of their bodies, should consume so much more food and much more as manure, and expend so much more on respiration, for a given amount of increase in weight as we have seen they do.

The other columns of the table show, however, that neither the amount of dry substance of food consumed nor the amount lost by respiration, by a given weight of animal within a given time, is in excess with the pig in anything like the proportion that its increase exceeds that of the other animals. In other words, the much higher character of the food of the pig is itself in the much greater rapidity, and in the greater proportion, of its conversion into the most valuable product of the feeding operation.

Lastly, in regard to the results in this table, it is remarkable that, whilst, for a given weight of animal within a given time, the amounts of increase in weight and of dry substance consumed in food, and lost by respiration, are so very different for the different animals, the amounts of dry substance voided in excreta are almost identical. I shall show further that the limit of consumption is much regulated by the amount of excreta.

at during the fattening process the proportion of a given weight of the body, of water, mineral and nitrogenous compounds decreases, whilst the fat very considerably increases.

at the carcass parts or saleable meat increase more rapidly than the internal parts or offal.

at the amount of dry substance of food required to produce a given weight of increase is less with the ox than with the sheep, and less with the sheep than with the pig.

at the dry substance of the food of the ox is a larger proportion of indigestible matter than of sheep, and that of sheep more than that

of oxen require from 5 to 6, and sheep from 3 to 4, as much time to add a given proportion of weight of their bodies as pigs.

at the greater portion of the nitrogenous and mineral matters of the food is recovered in the manure and that the greater part of the non-nitrogenous substance is lost by respiration, and other excretion—a much smaller proportion being retained in the crease, or voided in the manure.

at for a given amount of increase produced, the ox gives out considerably more substance as manure, and more in respiration, &c., than sheep, and the pig much more than pigs.

at for a given weight of dry substance consumed the oxen void more as manure than sheep, and the pig much more than pigs; but oxen respire rather less than sheep, and sheep rather less than pigs.

at in proportion to a given weight of animal, in a given time, oxen both consume and respire a greater substance of food than sheep, and sheep very much more than pigs; but they void almost identical quantities of dry substance as manure.

Comparative Feeding Value of different Foods, according to their Composition.

far I have endeavoured to indicate the characteristic points of distinction between the food of the sheep, and the pig, and to show in what its constituents are differently disposed of by different animals; and for the purposes of my illustration I have supposed the animals to be fed on such food as is recognised as appropriate to them, and in proportion and amount as experience justifies. I propose to say a few words on the relative properties of different foods, according to their composition.

coming out of view, just now, the incombustible mineral constituents, it will be convenient, as we are to consider the other constituents of food to be grouped under the heads of nitrogenous and non-nitrogenous substances.

among the nitrogenous substances, the most im-

portant of those which enter into our stock foods are albumen, casein, legumin, and gluten; and chemists and physiologists are accustomed to speak of these—the nitrogenous compounds—as the flesh-forming substances.

The non-nitrogenous constituents of our stock foods are starch, sugar, gum, pectin, oil, and cellulose, or woody fibre in different conditions of digestibility or induration. The non-nitrogenous compounds are spoken of as the respiratory or heat-producing, and fat-forming substances.

Now, writers on agricultural chemistry and physiology have generally assumed that it is chiefly the proportion of the nitrogenous or so-called flesh-forming substances contained in them, which determines the comparative value, for feeding purposes, of different foods.

The coloured diagram before you will enable you to judge whether or not this supposition is justified by the practical experience of feeding. This diagram has been constructed by the animals themselves. They know nothing about nitrogenous or non-nitrogenous constituents, digestible or indigestible cellulose, and so on; but they are gifted with an unerring instinct which enables them not only to distinguish between substances which are and are not food, but also to select from a variety of food stuffs those which are most suitable for the requirements of the system, and so to indicate to us the proper amounts and proportions of the different constituents.

In the experiments to which the diagram refers, as well as in many others, the plan has been to select foods containing very different proportions of nitrogenous and non-nitrogenous compounds; in fact, some containing two or three times as much nitrogen as others. We have then given to one set of animals a small fixed amount daily, of food containing a low percentage of nitrogen, and allowed them to take as much as they chose of another food, different in composition in this respect. To another set we have given a limited amount of food, rich in nitrogenous compounds, and allowed the animals to take, *ad libitum*, of a different description of food, and so on. In this way they have been enabled to fix for themselves the limit of their consumption of nitrogenous and non-nitrogenous constituents respectively, according to their wants.

The diagram shows the results of such experiments with pigs; and the foods employed were Indian corn meal, barley meal, bean meal, lentil meal, bran, and dried cod-fish, used alone, or in combination, as the case might be. Black being taken to represent nitrogenous substance, red non-nitrogenous substance, and green total dry organic matter (nitrogenous and non-nitrogenous together), the diagram is constructed as follows:—The smallest quantity of nitrogenous

non-nitrogenous, or total organic matter consumed in any one experiment is reckoned as 100; and the several lines above the base line, which is marked 100, indicate larger amounts, corresponding to the figures given at the side of the diagram.

The upper portion shows the relative amounts of each constituent consumed in each experiment per 100 lbs. live weight per week; that is to say, by a given weight of animal within a given time. A glance shows you that the height to which the colours representing the non-nitrogenous, or the total organic substance reach is very much more uniform than that indicating the consumption of nitrogenous substance. In fact, it is perfectly clear that the animals were guided in the amount of food which they consumed by the amount of non-nitrogenous, and not by that of the nitrogenous, constituents which it supplied.

But, according to current theories, the amount of nitrogenous substance ought at least to determine the amount of increase produced. The lower portion of the diagram shows what the animals have to say on this point. The arrangement is the same as before; but the results show not how much of each class of constituents was consumed by a given weight of animal within a given time, but how much was consumed to produce a given weight (100 lbs.) of increase.

Here again we see that the amount of either non-nitrogenous or total organic substance consumed varied comparatively little, whilst that of the nitrogenous substance consumed for the production of a given amount of increase varied from 100 to over 300 parts.

It is obvious, therefore, that both the amount of food consumed by a given weight of animal within a given time, and that required to produce a given weight of increase, were determined by the amount of available non-nitrogenous substance which the food supplied. The quantities required would, doubtless, have varied within even narrower limits, had all the foods contained equal proportions of indigestible woody matter.

It may be observed that it is doubtful whether pigs are able to digest cellulose, or woody fibre, at all; but here is no doubt, as the investigations of ourselves and others on the point sufficiently prove, that oxen and sheep are able to digest a considerable portion of such matter, when it is not in too indurated a condition.

It will, of course, be understood, that a certain amount and proportion of nitrogenous substance is essential in the food of animals; and if I were asked to state, in general terms, what was the approximate proportion of the nitrogenous to the digestible non-nitrogenous substances, below which they should not exist in the food of our stock, I should say

(though with reservations) about such as we find them in the cereal grains; and since few of stock foods are below, and many above, this in the proportion of nitrogenous substance, it results that we are more likely to give an excess than a deficiency of such constituents, so far as the requirements of the animal are concerned. The value of the nature depends, however, very much on the amount of the nitrogen which the food contains; but to this point I shall recur after directing attention to a few more points in connection with the comparative value of different foods as such.

Some years ago we published the results of our experiments on the equivalency of starch and sugar as food, pigs being the subject of the trial. Several having each a fixed and limited quantity of lentils and bran allowed, one was permitted to take as much starch, another as much sugar, and another as much of the mixture of the two as they chose; whilst in another experiment the animals were allowed to select at discretion from lentils, bran, sugar, or starch each placed separately within their reach. The result was, that sugar and starch were found to have weight for weight, practically the same value as constituents of food.

These results would, *a priori*, lead to an answer to the negative to the much agitated question, whether there is any advantage in malting barley for feeding purposes. The chief effect of the malting process is to convert starch into sugar—not, it is true, sugar of exactly the same description as that used in our experiments; but there is good reason for supposing that malt sugar would have a lower value than cane sugar as a food constituent; and direct experiments, made many years ago at Rothamsted, have shown that a given amount of malt, mixed with other food, gave less rather than more increase than the amount of barley from which it was produced. It is obvious, too, that as the conversion of barley into malt is a manufacturing process, attended with considerable cost, as well as actual loss of substance, the remission of the duty on malt employed for feeding purposes would not be likely to be of benefit to the farmer, unless either a given amount of malt sugar proved to be of considerably higher feeding value than the starch from which it was produced, or the other constituents were rendered more digestible and assimilable by the process.

This leads me, before leaving the subject of food, to make a few remarks on some other manufactured foods for stock. Many complaints are made, and justly made, of the adulteration of oil-cakes; and it is sometimes asserted that cheaper and better food than the average of cakes now in use could be manufactured with advantage both to the maker and the feeder. Linseed and other cakes are themselves

in one sense, manufactured foods. But the object of the manufacturer is not the production of cake, but of oil. If the farmer did not use the cake at all, it would still be made, and the oil would be sold for a higher price. As it is, the manufacturer makes the cake as a bye-product, and the price he gets for it enables him to sell his oil so much the cheaper.

But if manufactories were set up for the special purpose of preparing foods for stock, the whole cost of the undertakings must be charged upon the food. Lentils, beans, peas, Indian meal, barley meal, linseed, and other good staple foods must be used; and although it might be possible so to combine foods together that a given weight of the mixture would possess a somewhat higher feeding value than the component parts used singly, there is every reason to suppose that the increased cost would more than counterbalance any slight benefit that could be derived in that way. Nor do I anticipate that the progress of science will aid us much in this direction. Condimental foods have been tried, and found wanting; and I have little doubt that a similar result will attend the manufacture and use of simpler food mixtures. Our hopes as feeders must be in increased and cheap supplies of ordinary cattle foods of good quality, rather than in submitting those we have to costly processes of manufacture.

The results arrived at in regard to this portion of the subject may be briefly summed up as follows:—

1. The comparative feeding value of our current stock foods depends more upon the proportion of the digestible non-nitrogenous substances they contain than upon their richness in nitrogenous compounds; but the richer the food in nitrogen, the more valuable will be the manure.

2. Of the non-nitrogenous constituents of food, starch and cane sugar have, weight for weight, nearly equal feeding values; malt sugar has probably rather a lower value than either cane sugar or starch; digestible cellulose, in moderate proportion, has, for ruminant animals, probably nearly the same value as starch; and fat or oil have probably about $2\frac{1}{2}$ times the value of starch for the purposes of respiration, or the storing up of fat in the body.

3. Some advantage results in a feeding point of view from the judicious mixture of a variety of ordinary stock foods; but the benefit to be derived in this way is not such as to compensate for the extra cost of a special manufacturing process to attain it.

Connection between the Value of the Manure and the Composition of the Food consumed.

The next and last branch of the subject relates to the comparative value of the different constituents in the liquid and solid excretions of the animals, and to

the connection between the value of the manure and the composition of the food from which it is derived.

I have already pointed out that the great value of the carbon, hydrogen, and oxygen of either passes into the increase or off in respiration that comparatively little of any of them is in manure. By far the larger portion of the carbon and nearly the whole of the mineral matter are, however, so recovered.

To show the economic connection between feeding of stock for the production of manure, and the growth of corn, I propose to give a few results obtained in experiments on the wheat by different manures. In the experiment question, wheat has been grown for 20 seasons on the same land.

In Table V. are given the average annual yield of corn and straw, and the estimated yield per acre, over the last 12 years, respectively, with mineral manure alone, with ammoniacal manure (ammonia salts), and farm-yard manure.

TABLE V.—Average annual Produce of Wheat sown on the same land, estimated yield of Carbon, per acre, over 12 years.

Manures, per acre, per annum.	Average annual produce		
	Dressed Corn.	Total Corn.	Straw
	Bushels.	lbs.	lbs.
Unmanured ...	16½	964	166
Mineral manure alone ...	18½	1167	189
Mineral manure and 400 lbs. ammonia salts }	36½	2275	421
14 tons farm-yard manure	35½	2232	388

Where the farm-yard manure was employed, as well as more of every other crop was annually applied in manure than removed. In the other cases no carbon was supplied in the manure; and yet it will be seen that where the mineral manure and ammonia were employed (the latter containing a large amount of nitrogen), the yield of carbon was greater than where a large amount of that substance was supplied by means of farm-yard manure. This carbon has been derived from the atmosphere.

In experiments in this field last year, from 1½ tons of carbon per acre were removed in the crop, any being supplied in manure; but in the other large quantities of nitrogen were supplied.

The quantity of carbonic acid required for the growth of the crop is about as much as is given off into the atmosphere in a year by the individuals of a mixed population of both sexes, and it will be seen that it is under the action of ammoniacal or nitrogenous manure that the amount of carbon has been fixed in the vegetable carbonic acid of the atmosphere.

Table III. showed how small was nitrogen consumed by an animal stored up in its increase and at. If there were none of the lost in the various exhalations whole of that not stored up in found in the manure. But the selves and others show that a nitrogen is so lost. Our own nine the limit of this loss, and der which it is greater or less, ar back as 1847, and have been from that time to the present ; few years we have collected a mental data on the subject; but analytical work is not yet con- that I am in a position to give ent of the results obtained. It ted as beyond doubt, that by far the nitrogen consumed in food limals in their liquid and solid e higher the proportion of nitro- cher will be the excrements in itment of manures.

published a table showing the the manure obtained from the on of different articles of food a practice. The valuation was ledge of the average composition iptions of food, and upon infor- the course of the experiments the probable average amount of od valuable for manure which e solid and liquid excrements of

ts of these valuations in very ay be said that the estimated from 1 ton of oil-cake was con- that from the same quantity of , beans, or peas; from two to as that from 1 ton of oats, wheat, or hay; from seven to ten times e same weight of oat, wheat, or about twenty times as much as

efore, that in the selection of stock, it is very important to ring as well as their feeding value. this point will suffice. A ton of rtainly not yield nitrogen in the nals consuming it equal to more as, $\frac{1}{2}$ cwt., or 28 lbs., of ammonia ; ke will yield 1 cwt., or four times ore, we take the ammonia in the , the amount of it obtained from : ton of locust beans will be

worth only 16s. 4d. ; whilst that from the ton o cake will be £3 5s. 4d.

There is, in fact, far greater difference in t nuring than in the feeding value of most ordinary stock foods in the market.

In illustrating the comparative value of the obtained from different foods, by reference me the amounts of nitrogen or ammonia-yielding ; which they supply, it will not be understood in any way ignore or under-rate the value mineral constituents associated with the nitro- matter in the excrements. But, inasmuch amount of mineral constituents voided is ge in excess of that required for the due effect as n of the nitrogen with which they are accompa results that the amount of the nitrogen or am- yielding matter is practically the best index value of the manure.

Appropriateness of Animal Food in the Diet of

It will be obvious that the importance of th ject which I have brought before you this e rests upon the assumption that animal food important element in the diet of man. The indeed, some who maintain that a purely ve diet would be more suitable and natural th mixed vegetable and animal one so generall ferred. If their view were adopted, we n longer trouble ourselves about the connection b the food, the increase, and the manure of fa oxen, sheep, and pigs. There are, however, circumstances, economical and physiological, p to the appropriateness of admitting a certain g tion of animal food into the diet of man. To two of these I will briefly refer.

Walking is for man undoubtedly a very means of progression. Still, it is often very tageous to ride, and so to employ the legs of rruped instead of our own. In eating meat v be said to employ the stomachs of other anil do that which we could not so well do with o. As a few ounces of gold are separated fro tons of rock by the combined aid of mechanil chemical processes, so the animals feeding crude, and often to us indigestible, vegetable eliminate from it, and store up in their bodies, its constituents in a form at once much more trated than that in which they consumed the much more easily appropriated by the hun nomy. A given amount of nitrogenous con in the form of meat is undoubtedly more ea gested and assimilated by man than if t amount were supplied in the form of beans. again, the animals convert starch, sugar, d probably some of them cellulose, which we c digest at all), into fat, which has twice and

respiratory and fat-storing capacity of the substances from which they produce it. It is, doubtless, true that man can produce fat, and keep up his respiratory function, from starch and sugar; but it can hardly be doubted that there is some economy to his system in having a portion of fat supplied to him ready made.

Apart from the strong testimony of common experience on the subject, there is evidence in the comparative structure of man that he is adapted for a concentrated form of food. One illustration, in passing, may be adduced on this point. Table VI. shows the proportion of the stomach, by weight, in a given live weight of oxen, sheep, pigs, and man.

TABLE VI.—*Proportion of stomach in different animals.*

Stomach in 100 lbs. live weight:—

Oxen ...	51 ounces.	Sheep ...	89 ounces.
Pigs ...	14 "	Man ...	6 "

Relative weight does not, of course, necessarily represent with numerical exactness relative capacity or size. But there is little doubt that there is a gradation in the capacity of the stomach relatively to a given weight of the body in the animals enumerated in the order, and to a great extent in the degree indicated by the figures given in the table. Admitting this to be the case, we have seen that the sheep, with its less proportion of stomach than the ox, takes a somewhat more concentrated food; and that the pig, with its much less proportion of stomach than the sheep, requires a much more concentrated food than the latter. May we not conclude that man in his turn, with his less proportion of stomach than the pig, will also appropriately take a more concentrated food than his useful friend?

The food of man is, indeed, very closely allied, in a chemical point of view, to that of the pig. The staple of the food of both the fattening pig, and man, is cereal grain. The pig, it is true, consumes the husk as well as the farinal portion, whilst man does not; but we know that this proportion of indigestible woody matter is very nearly the limit of that which is appropriate for the fattening pig; and that on the addition of a small quantity of bran the proportion of increase diminishes, and that of the dry substance of the food voided as excrement increases. The only other essential difference is, that the pig takes, as a rule, the whole of his nitrogenous compounds in the form of vegetable products, and a much larger proportion of starch, and other non-nitrogenous compounds, more bulky in relation to their respiratory and fat-forming capacity than fat itself. Not, indeed, that the pig is at all unapt or unwilling to adopt even still more closely the diet of man; for he will take *animal flesh* and fat when he can get them, and, what is more, he likes them better cooked than raw.

Were it not, then, that man separates from the flour, and that he gets lower eliminate in an easily digestible form a p nitrogenous aliment, from foods which h himself readily digest, and that he gets to provide him with a portion of his resp fat-storing food in the concentrated i itself, we could hardly account for the tion to a given weight of the body of the the receptacle and first laboratory of the i case than in that of the pig. We kno that in the cases where man is reduced for nearly the whole of the non-nitrogen tuents of his food upon starch, in the fo toes or rice, that there is a disposition to ment of the abdominal organs, and to a in physical and mental energy.

To conclude on this point, there can t whatever that the food of the labouring i proved when he can add to his bread a po bacon, or butter, or fat in some other it is better still if he can substitute or su little butcher's meat. Indeed, that whic experience recognises as high quality i within certain limits, high proportion o vegetable food, and with it high proporti starch and other non-nitrogenous compou

But not only do the animals which we our own food convert vegetable produce either could not digest at all, or could d easily than they, into concentrated and gestible and assimilable material for our t doing this they supply carbonic acid to i phere, and return the most important ma stituents of their food in their excren providing, to both the soil and the atmos crude vegetable products, that which is ne the luxuriant growth of cereal grain, and c table produce suited for the direct use i man.

Were it not for such compensations, by t of man and other animals upon the sur earth (if it could take place at all), by th quantities of carbonic acid evolved into the from the combustion of coal and from oth and by the gradual destruction of forests, w chief natural agents for restoring the b purity of the atmosphere would becom But the grasses, which supply so large a of the food of beasts, and the cereals an plants of the same great family, which sup man in almost every climate, serve to carbon given into the atmosphere in th carbonic acid. It may seem at first i that the humble grasses, and the corr ing only a few feet from the surface

should be able to take up more carbonic acid, and evolve more oxygen, over an acre of land than an acre covered with forest trees. Still, there can be little doubt that more carbon is fixed in an acre of luxuriant wheat than over the same area of woodland; and there can be as little that an acre of sugar cane would fix more than an equal area of the most luxuriant tropical forest.

Conclusion.

With a few general remarks of a practical nature, I will conclude my discourse. The great change which has taken place in the practice of feeding stock in modern times has consisted in bringing the animals much earlier to maturity, by means of careful breeding, and more liberal feeding. Scales and weights were seldom used in agricultural experiments until comparatively recently; but there are some few records of the results of feeding as practised at the latter end of the last century, which will serve as in instituting a comparison between the results then obtained and those which are possible, or even common, at the present day.

In 1794 the Duke of Bedford made some experiments to determine the comparative feeding qualities of South Down, Leicester, Worcester, and Wiltshire sheep. Twenty of each were selected and weighed on November 19, 1794. To each lot were allotted 16 acres of pasture, and in the winter some turnips were thrown upon the pasture, and a small quantity of hay was also provided. On February 16, 1796, after a period of 65 weeks of feeding, the experiment was concluded, and the sheep sent to market.

Over the whole period the sheep gave an average increase of between 40 and 50 lbs. per head; and as their original weight was nearly 100 lbs. per head, they increased nearly 50 per cent. from the store or lean to the fat condition, which is the same proportion as that assumed in the illustrations to which Table III. refers.

Some years ago, I tried a set of experiments upon the comparative fattening qualities of South Downs, Hampshire Downs, Cotswolds, Leicesters, and cross-bred wethers, and cross-bred ewes, each lot consisting of between 40 and 50 sheep. They were put up in November, when their weights averaged very nearly the same as those of the Duke of Bedford's sheep; and when fat, they had increased in about the same degree, namely, to an average of about 150 lbs. each. The Duke of Bedford's sheep were about 65 weeks in adding 50 lbs. to their weight, and mine in some cases 20, and in others a little more, or about one-third the

time. It is somewhat singular that in May—the at which my sheep were consumed as mutton the Duke of Bedford's were weighed for the first time the commencement of the experiment, and were to have increased only about 6 lbs. per head.

The difference of result in these two cases almost entirely due to the difference in the feeding. Formerly, sheep received perhaps turnips on their pasture, and but little dry food that not of high feeding quality; and the consequence was, that during the colder months of the year either lost weight or increased but little. Now have a liberal allowance of good food, and are consequently protected from the inclemency of the weather. In my own experiments, just referred to, the sheep were allowed from $\frac{3}{4}$ lb. to 1 lb. of oil-cake head per day, according to their weight, about the same amount of clover chaff, and as many as they chose to eat; and they gave an average increase of nearly 2 per cent. upon their weight.

There is no doubt that in rapidly fattening stock at an early age, quality of meat is to some extent sacrificed to quantity. But it is only by means of the system of liberal feeding and early maturation that meat can be brought within the reach of the masses of the population. The farmer, too, must adopt that system which will pay him the best; and the difference between the price which the consumer will give for a pound of four-year-old and one-year-old mutton will, only under very exceptional circumstances of locality, remunerate him for the extra cost of production.

In conclusion, I have only now to thank you for the very kind attention with which you have followed me through what I fear may be thought by you somewhat tedious detail. The subject of the chemistry of feeding is, however, essentially a complicated one; and I think you will have learnt from my lecture, if you did not know it before, that there remains much to be determined by careful investigation respecting it. But if I have in any degree succeeded in indicating the proper points of view in which this, at once practical and scientific, subject should be studied, and in impressing upon you some prominent and important facts regarding it, as to lead to improvement in practice by the knowledge of principle, or to further inquiry to an extension of our knowledge, I shall feel that the objects of my desire and endeavour in addressing you have been fully attained.

**AVERAGE COMPOSITION, PER CENT. AND PER TON, OF VARIOUS KINDS OF
AGRICULTURAL PRODUCE, &c.**

	PER CENT.					LBS. PER TON.			
	Total Dry Matter.	Total Mineral Matter (ash).	Phosphoric Acid, reckoned as Phosphate of Lime.	Potash.	Nitrogen.	Total Dry Matter.	Total Mineral Matter (ash).	Phosphoric Acid, reckoned as Phosphate of Lime.	Potash.
1. Linseed Cake	88.0	7.00	4.92	1.65	4.75	1971	156.8	110.2	37.0
2. Cotton Seed Cake	89.0	8.00	7.00	3.12	6.50	1994	179.2	156.8	70.0
3. Rape Cake	89.0	8.00	5.75	1.76	5.00	1994	179.2	128.8	39.4
4. Linseed	90.0	4.00	3.38	1.37	3.80	2016	89.6	75.7	30.7
5. Beans	84.0	3.00	2.20	1.27	4.00	1882	67.2	49.3	28.4
6. Peas	84.5	2.40	1.84	0.96	3.40	1893	53.8	41.2	21.5
7. Tares	84.0	2.00	1.63	0.66	4.20	1882	44.8	36.5	14.8
8. Lentils	88.0	3.00	1.89	0.96	4.30	1971	67.2	42.3	21.5
9. Malt Dust	94.0	8.50	5.23	2.12	4.20	2106	190.4	117.1	47.5
10. Locust Beans	85.0	1.75			1.25	1904	39.2		
11. Indian Meal	88.0	1.30	1.13	0.35	1.80	1971	29.1	25.3	7.8
12. Wheat	85.0	1.70	1.87	0.50	1.80	1904	38.1	42.0	11.2
13. Barley	84.0	2.20	1.35	0.55	1.65	1882	49.3	30.2	12.3
14. Malt	95.0	2.60	1.60	0.65	1.70	2128	58.2	35.8	14.6
15. Oats	86.0	2.85	1.17	0.50	2.00	1926	63.8	26.2	11.2
16. Fine Pollard	86.0	5.60	6.44	1.46	2.60	1926	125.4	144.2	32.7
17. Coarse Pollard	86.0	6.20	7.52	1.49	2.58	1926	138.9	168.4	33.4
18. Bran	86.0	6.60	7.95	1.45	2.55	1926	147.8	178.1	32.5
19. Clover Hay	84.0	7.50	1.25	1.30	2.50	1882	168.0	28.0	29.1
20. Meadow Hay	84.0	6.00	0.88	1.50	1.50	1882	134.4	19.7	33.6
21. Bean Straw	82.5	5.55	0.90	1.11	0.90	1848	124.3	20.2	24.9
22. Pea Straw	82.0	5.95	0.85	0.89		1837	133.3	19.0	19.9
23. Wheat Straw	84.0	5.00	0.55	0.65	0.60	1882	112.0	12.3	14.6
24. Barley Straw	85.0	4.50	0.37	0.63	0.50	1904	100.8	8.3	14.1
25. Oat Straw	83.0	5.50	0.48	0.93	0.60	1859	123.2	10.7	20.8
26. Mangold Wurzel	12.5	1.00	0.09	0.25	0.25	280	22.4	2.0	5.6
27. Swedish Turnips	11.0	0.60	0.13	0.18	0.22	246	13.4	2.9	4.0
28. Common Turnips	8.0	0.68	0.11	0.29	0.18	179	15.2	2.5	6.5
29. Potatoes	24.0	1.00	0.32	0.43	0.35	537	22.4	7.2	9.6
30. Carrots	13.5	0.70	0.13	0.23	0.20	302	15.7	2.9	5.1
31. Parsnips	15.0	1.00	0.42	0.36	0.22	336	22.4	9.4	8.1

SEWAGE OF TOWNS.

THIRD REPORT

OF

THE COMMISSION

APPOINTED

TO INQUIRE INTO THE BEST MODE

OF

DISTRIBUTING THE SEWAGE OF TOWNS,

AND

**APPLYING IT TO BENEFICIAL AND
PROFITABLE USES;**

WITH APPENDICES 1, 2, AND 3.



LONDON:

**PRINTED BY GEORGE E. EYRE AND WILLIAM SPOTTISWOODE,
PRINTERS TO THE QUEEN'S MOST EXCELLENT MAJESTY.**

1865.

CONTENTS

REPORT OF THE COMMISSION - - - - -	Page 3
SECOND REPORT OF THE EXPERIMENTS MADE BY ORDER OF THE COMMISSION AT RUGBY :—	
Introductory Observations - - - - -	5
I.—Experiments on the application of sewage to meadow land; quantities of sewage applied, and of green produce obtained -	6
II.—Experiments on the application of sewage to Italian rye grass -	16
III.—Experiments with fattening oxen - - - - -	19
IV.—Experiments with milking cows - - - - -	21
V.—Composition of the Rugby sewage - - - - -	27
VI.—Estimated average composition of the Metropolitan sewage -	37
VII.—Composition of the drainage water (Rugby) - - - - -	45
VIII.—Chemical composition of the unsewaged and sewaged grass -	51
IX.—Effects of sewage on the mixed herbage of meadow land in developing the more freely growing at the expense of the less freely growing plants - - - - -	56
X.—Composition of the milk yielded from the unsewaged and sewaged grass - - - - -	58
XI.—Experiments on the application of sewage to oats - - - - -	60
XII.—Miscellaneous results obtained in 1864 - - - - -	62
XIII.—Concluding observations; general considerations on the agri- cultural utilisation of town sewage - - - - -	66
Summary - - - - -	72
APPENDIX No. 1.—TABLES I.—XLIV. inclusive - - - - -	81
APPENDIX No. 2.—Notes on the Edinburgh Sewage Meadows -	198
APPENDIX No. 3.—Notes on the Croydon Sewage Meadows -	202

THIRD REPORT OF THE COMMISSION.

TO THE LORDS COMMISSIONERS OF HER MAJESTY'S TREASURY.

MAY IT PLEASE YOUR LORDSHIPS,

WE, the undersigned, whom Her Majesty's Commission, bearing date 5th January 1857, appointed to "inquire into the " best mode of distributing the Sewage of Towns, and applying " it to beneficial and profitable uses," have now again, according to our instructions, the honour of reporting to your Lordships our further progress in the matter committed to us for inquiry.

Since the date of our last Report (August 1861) we have, through a committee of our number, consisting of Mr. Lawes and Professor Way, continued at Rugby the experiments which were undertaken in 1861 on the application of sewage to land. The report of that committee, which we append, contains the results for the three years 1862-4.

Your Lordships will observe that these experiments have not been confined to the application of sewage in different quantities to land, but have extended to the consumption, by cattle, of the produce so obtained, and to the production of meat and milk, and have been accompanied by a careful record of the quantities and market-value of the products, and by numerous analyses of the sewage before and after irrigation, as also of the grass and of the milk.

It appears to us that these experiments have solved many of the difficulties which have hitherto attached to the question of the agricultural application of sewage, and that they leave no reasonable doubt of the practicability and advantage of so employing the sewage of towns.

We have also continued to give our best attention to all kindred experiments and inquiries which have been going on elsewhere.

As the results of our labours, extending over eight years, we have confidence in submitting to your Lordships the following conclusions:—

1. The right way to dispose of town sewage is to apply it continuously to land, and it is only by such application that the pollution of rivers can be avoided.
2. The financial results of a continuous application of sewage to land differ under different local circumstances; first, because in some places irrigation can be effected by gravity, while in other places more or less pumping must be employed; secondly, because heavy soils (which in

given localities may alone be available for the purpose) are less fit than light soils for continuous irrigation by sewage.

3. Where local circumstances are favourable, and undue expenditure is avoided, towns may derive profit, more or less considerable, from applying their sewage in agriculture. Under opposite circumstances, there may not be a balance of profit; but even in such cases a rate in aid, required to cover any loss, needs not be of large amount.

Finally, on the basis of the above conclusions, we further beg leave to express to your Lordships that, in our judgment, the following two principles are established for legislative application:—

First, that, wherever rivers are polluted by a discharge of town sewage into them, the towns may reasonably be required to desist from causing that public nuisance:

Second, that where town-populations are injured or endangered in health by a retention of cesspool-matter among them, the towns may reasonably be required to provide a system of sewers for its removal.

And should the law, as it stands, be found insufficient to enable towns to take land for sewage-application, it would, in our opinion, be expedient that the legislature should give them powers for that purpose.

(Signed) ESSEX.
ROBERT RAWLINSON.
J. THOMAS WAY.
J. B. LAWES.
JOHN SIMON.

8, Richmond Terrace, Whitehall,
March 1865.

SECOND REPORT of EXPERIMENTS on the APPLICATION of
TOWN SEWAGE to GRASS LAND, conducted at RUGBY, by
ORDER of the ROYAL SEWAGE COMMISSION.

SEASONS 1861, 1862, and 1863.

IN the Second Report of the Commission, presented to both Houses of Parliament in 1862, an account was given of the results obtained in the First Season (1861) of Experiments on the application of town sewage to grass land, which were undertaken by order of the Royal Sewage Commission, and conducted in the neighbourhood of Rugby, where, arrangements being made for the distribution of the sewage of the town over a considerable area of adjacent land, the conditions were considered well adapted for the purposes of the inquiry.

As stated in the preliminary Report above referred to, the Commission, guided by the information acquired in the course of their investigation of the then existing experience on the subject, which had led them to visit almost every locality where town sewage was applied in any way to the purposes of agriculture, had come to the conclusion that to obtain the largest amount and value of produce at the least proportionate cost for distribution, dilute town sewage should be applied to the growth of succulent crops, and that it was best adapted for grass. It was decided, therefore, to confine attention, at any rate in the first instance, to grass alone.

In arranging the experiments, it was considered that the object was to provide such information as might be taken as the basis of arrangements for the application of the sewage of towns, in the manner the most advantageous to both urban and rural interests.

To this end it was sought to determine :—

1. The amount and the composition of the produce, in relation—to the amount of water supplied to the land by irrigation, to the amount of manurial constituents so applied, and to the amount of population contributing the manurial constituents to the water.
2. The most profitable method of utilising the produce; that is, whether it should be used in the green state or as hay; whether for the production of milk or of meat; and whether it should be consumed alone or in conjunction with other food.

In the experiments of the first year, three portions of land, of about five acres each, were operated upon, and each of these was divided into four plots, to be treated, respectively, as follows :

Plot 1. To be unsewaged.

Plot 2. To be irrigated with sewage at the rate of 3,000 tons per acre per annum.

Plot 3. To be sewaged at the rate of 6,000 tons per acre per annum.

Plot 4. To be sewaged at the rate of 9,000 tons per acre per annum.

The produce of one such set of four plots was to be given, in the green state, to fattening oxen; that of the second (also in the green state) to milking cows; and that of the third was to be made into hay.

As explained in the former Report, owing to deficient supply, but little sewage was applied to the portion of land devoted to the production of hay; and, since the first season, the five acres in question have each year been sublet.

The results obtained in the first year's experiments on the other portions of land were given in detail in the previous Report; but it was admitted that the experience of one year only could be taken as little more than initiative on many points; it being obviously essential to determine the effects of the continued application, and the influence of seasons of different characters, before safe deductions could be drawn in regard to some of the most important economical questions at issue.

The results of two more seasons (1862 and 1863) are now at command; and it is proposed to call attention chiefly to the difference of result obtained in the different seasons, and to the average result over the three seasons, making but few comments on those of each separate season. The full details will, however, be given for reference, in the tabular form, in Appendix, No. 1., p. 81, et seq.

In addition to the experiments above referred to, which are a continuation of those already reported, by the kindness of Mr. Campbell some results obtained on the application of sewage to Italian rye-grass and to oats are also given.

For the convenience of reference and comparison, the numerical results will, as far as possible, be arranged in the same form, and the subject considered in the same order, as in the former Report.

I. Quantities of Sewage applied, and of Green Produce obtained.

In the first season (1861) the application of sewage did not commence until March in the one field, and April in the other; but, as it was considered that any scheme for the general application of town sewage to agricultural purposes must of necessity *be based on the fact of a daily supply the year round*, which

must be dealt with in winter when of comparatively little value as well as in summer when of more, the amounts of 3,000, 6,000, and 9,000 tons, respectively, were, in the second and third seasons, distributed over the entire year, and the quantities supplied from November 1st of one year to October 31st of the next were taken as those to which the increase of crop was due.

The detailed records relating to the application of the sewage are given for reference in Tables I. and II. pp. 81-89. Appendix, No. 1. Of these, Table I., which now follows, is a convenient summary.

TABLE I.—Quantities of Sewage applied per Acre, on each Plot, in each Month, in each of the Three Seasons.

	SEWAGE PER ACRE.					
	FIVE-ACRE FIELD.			TEN-ACRE FIELD (half).		
	Plot 2.	Plot 3.	Plot 4.	Plot 2.	Plot 3.	Plot 4.

1st SEASON, 1861; March—October, inclusive.

	Tons.	Tons.	Tons.	Tons.	Tons.	Tons.
March - - -	632'1	1045'1	1444'2
April - - -	279'9	666'4	1177'0	563'0	1145'9	1376'9
May - - -	75'8	96'5	97'7	18'3	64'1	116'8
June - - -	78'8	223'3	577'2	392'3
July - - -	531'7	430'2	654'1	512'0	392'2	995'7
August - - -	130'6	580'2	787'3	225'9	316'3	596'1
September -	143'1	703'3	614'7	34'0	517'7	381'8
October - -	201'7	678'2	800'7	34'0	367'7	455'8
Totals - - -	2073'7	4423'2	6152'9	1367'2	2993'9	4226'4
Rate per annum -	5110'4	6634'8	9229'1	2378'1	4906'7	7346'4

2d SEASON, 1862; November 1861—October 1862, inclusive.

November 1861 -	513'1	499'8	745'0	116'3	341'5	451'4
December „ -	126'6	429'8	537'6	71'5	143'5	277'8
January 1862 -	323'0	457'6	563'3	77'4	189'2	232'7
February „ -	169'2	575'0	751'6	227'5	167'2	339'1
March „ -	235'4	455'8	478'2	109'9	374'0	524'7
April „ -	211'8	503'0	720'1	506'5	903'5	876'8
May „ -	281'8	581'4	763'0	269'6	495'1	1201'5
June „ -	77'1	164'7	292'7	173'0	327'0	416'0
July „ -	589'1	846'4	1223'7	595'5	1255'9	2017'0
August „ -	309'1	476'9	1442'0	397'5	1000'8	1222'6
September „ -	79'4	623'6	628'6	177'6	396'3	776'9
October „ -	285'2	386'5	730'9	235'2	485'5	517'8
Totals -	2991'8	5991'5	8986'7	2999'5	5599'5	9004'2

TABLE I.—*continued.*

Quantities of Sewage applied per Acre, on each Plot, in each Month,
in each of the Three Seasons.

	SEWAGE PER ACRE.					
	FIVE-ACRE FIELD.			TEN-ACRE FIELD (half).		
	Plot 2.	Plot 3.	Plot 4.	Plot 2.	Plot 3.	Plot 4.
3d SEASON, 1863; November 1862—October 1863, inclusive.						
ber 1863 -	201'9	513'0	806'8	288'2	597'6	771'3
ber " -	114'2	315'2	500'1	151'6	623'5	894'6
ry 1863 -	351'2	469'3	580'2	463'6	436'0	840'3
ry " -	194'6	550'6	655'4	200'9	550'9	1013'6
" -	462'7	774'4	1375'8	203'9	390'8	397'0
" -	234'2	550'9	640'5	203'9	395'8	592'5
" -	90'4	579'3	553'3	39'8	582'9	748'3
" -	896'6	540'9	462'5	385'4	974'8	940'1
" -	100'9	426'4	980'6	492'4	344'8	779'6
" -	..	409'5	964'2	150'0	..	717'0
ber,, -	275'1	617'3	680'3	..	640'9	495'1
r " -	57'1	232'8	819'0	391'3	459'8	797'1
Totals -	2998'9	5999'6	9000'7	3001'0	5999'8	8996'5

SUMMARY. 1861, 1862, and 1863.

Season; to } 31, 1861	2073'7	4423'2	6152'9	1387'2	2803'9	4226'4
Season; to } 31, 1862	2991'8	5994'5	8986'7	2999'5	5999'5	9004'3
Season; to } 31, 1863	2998'9	5999'6	9000'7	3001'0	5999'8	8996'5
Totals -	8064'4	16417'3	24140'3	7387'7	14803'2	22229'2

It is seen that when calculated over the entire season the supply of sewage was in each case very nearly at the rate intended, but it varied considerably from month to month. Regularity in this respect was, indeed, sought to be attained within certain limits, but it was necessarily subject to various controlling circumstances. Thus, the sewage was, as a rule, applied on the same day for an entire day at a time, in order the better to secure its distribution over the whole, and the rate of flow, and consequently the day's supply, varied considerably. Then, again, throughout the summer months the stage of growth of the crop influenced the time of application, which was regulated so as to ensure, as far as possible, the beneficial action of the total quantity applied in each case. Occasionally, too, the supply was temporarily stopped, owing to derangements in the machinery, after which there was unfortunately no control.

The amounts of green grass obtained from the respective plots, each of the three seasons, are shown in Tables II. and III. (pp. 9 and 10); Table II. giving the amounts obtained in each month, and Table III. those in each successive crop. For further details, see Appendix Tables, III. and IV., pp. 90-96.

TABLE II.—Amount of Green Grass obtained during each separate Month.

	GREEN GRASS PER ACRE.											
	FIVE-ACRE FIELD.						TEN-ACRE FIELD (half).					
	Without Sewage.			With Sewage.			Without Sewage.			With Sewage.		
	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.
FIRST SEASON, 1861.												
May	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.	tona. cwt. gra. lbs.
June	3 2 3 27	4 17 0 14	7 19 1 23	8 14 3 8	4 10 1 1	4 11 2 8	5 4 1 7	8 16 1 7	2 5 0 0	2 5 0 0	2 5 0 0	2 5 0 0
July	3 1 0 8	2 8 0 15	4 6 0 26	6 14 0 13	0 8 1 18	5 5 1 2	5 8 0 19	1 16 3 24	0 8 1 18	5 5 1 2	5 8 0 19	1 16 3 24
Aug.	- - -	3 16 2 23	4 12 2 8	5 19 2 24	1 6 3 9	0 10 2 28	4 12 1 1	5 4 2 7	1 6 3 9	0 10 2 28	4 12 1 1	5 4 2 7
Sept.	2 2 2 8	1 3 0 12	6 5 1 7	5 0 0 5	2 3 2 7	0 11 0 13	1 15 0 3	4 16 2 0	2 3 2 7	0 11 0 13	1 15 0 3	4 16 2 0
Oct.	0 19 0 18	2 11 3 0	0 8 0 24	0 16 0 23	0 9 0 8	3 19 1 15	4 0 3 7	3 6 0 2	0 9 0 8	3 19 1 15	4 0 3 7	3 6 0 2
Nov.	- - -	- - -	3 9 1 6	4 2 0 5	- - -	- - -	- - -	- - -	- - -	- - -	- - -	- - -
Dec.	- - -	- - -	- - -	- - -	- - -	- - -	- - -	- - -	- - -	- - -	- - -	- - -
Totals	9 5 3 5	14 16 3 8	27 1 0 10	32 16 3 8	8 18 0 15	15 16 3 2	23 15 2 12	26 13 3 12	8 18 0 15	15 16 3 2	23 15 2 12	26 13 3 12
SECOND SEASON, 1862.												
May	- - -	0 19 1 19	9 15 3 9	8 7 3 6	0 17 1 16	1 14 0 5	11 13 2 7	8 14 0 21	- - -	- - -	- - -	- - -
June	0 8 1 26	5 10 0 10	5 16 0 26	5 15 1 15	7 10 2 11	15 0 0 3	2 7 2 20	1 15 1 7	0 8 1 26	5 10 0 10	5 16 0 26	5 15 1 15
July	3 16 0 23	8 2 3 3	- - -	4 19 1 19	2 11 2 0	0 8 2 24	6 9 3 6	8 16 1 2	3 16 0 23	8 2 3 3	- - -	4 19 1 19
Aug.	2 8 2 10	4 16 2 16	9 12 0 25	2 0 2 24	0 11 3 0	6 12 1 3	- - -	2 15 3 8	2 8 2 10	4 16 2 16	9 12 0 25	2 0 2 24
Sept.	- - -	4 0 1 10	6 12 2 10	9 15 0 0	3 8 0 17	0 16 0 8	9 16 1 22	6 2 2 13	- - -	- - -	- - -	- - -
Oct.	1 10 0 7	4 8 3 16	2 13 1 5	1 11 1 14	- - -	3 0 0 5	1 14 3 15	3 8 0 26	1 10 0 7	4 8 3 16	2 13 1 5	1 11 1 14
Nov.	- - -	- - -	- - -	- - -	1 10 3 9	- - -	- - -	- - -	- - -	- - -	- - -	- - -
Totals	8 3 1 10	37 18 0 18	34 10 0 19	32 9 2 22	16 10 0 25	27 11 0 20	32 2 1 14	31 12 1 26	8 3 1 10	37 18 0 18	34 10 0 19	32 9 2 22
THIRD SEASON, 1863.												
April	- - -	- - -	- - -	3 16 3 12	- - -	- - -	- - -	3 14 3 4	- - -	- - -	- - -	- - -
May	- - -	4 9 1 25	13 4 3 18	6 0 0 15	- - -	- - -	9 0 0 24	4 7 2 27	- - -	- - -	- - -	- - -
June	- - -	6 4 3 3	3 9 2 20	10 13 2 0	3 7 3 25	13 1 3 22	5 16 0 22	11 13 2 21	- - -	- - -	- - -	- - -
July	2 18 3 14	6 7 3 9	5 17 2 2	- - -	2 19 3 27	5 13 0 6	4 15 0 13	0 9 3 11	2 18 3 14	6 7 3 9	5 17 2 2	- - -
Aug.	- - -	- - -	4 2 3 5	7 15 3 16	- - -	- - -	5 14 3 0	7 16 2 14	- - -	- - -	- - -	- - -
Sept.	0 19 3 27	2 7 1 27	6 13 0 23	7 14 0 2	1 12 3 23	6 1 0 2	1 5 1 12	0 13 3 4	0 19 3 27	2 7 1 27	6 13 0 23	7 14 0 2
Oct.	- - -	2 12 1 15	2 5 3 5	0 14 1 20	- - -	- - -	3 9 3 20	5 12 2 3	- - -	- - -	- - -	- - -
Nov.	- - -	0 3 0 16	0 5 2 10	0 5 2 24	- - -	0 9 1 6	0 10 0 5	0 11 2 4	- - -	- - -	- - -	- - -
Totals	4 13 3 13	23 5 0 11	34 18 1 27	37 0 2 5	8 0 3 19	25 1 1 8	30 11 2 13	34 19 1	4 13 3 13	23 5 0 11	34 18 1 27	37 0 2 5

TABLE III.—Amounts of Green Grass obtained in each successive Crop.

	GREEN GRASS PER ACRE.																															
	FIVE-ACRE FIELD.										TEN-ACRE FIELD (half).																					
	Without Sewage.		With Sewage.								Without Sewage.		With Sewage.																			
			Plot 1.		Plot 2.		Plot 3.		Plot 4.				Plot 1.		Plot 2.		Plot 3.		Plot 4.													
FIRST SEASON, 1861.																																
1st Crop	6	4	0	7	7	5	1	1	10	7	0	24	13	5	1	23	4	18	3	19	9	13	5	3	2	11	1	1	4			
2d Crop	3	1	2	26	4	3	1	25	7	8	1	0	9	15	0	11	3	19	1	24	3	11	0	0	5	18	2	17	7	1	2	6
3d Crop	-	-	-	-	3	4	0	11	5	16	1	8	5	14	0	26	-	-	-	-	3	2	3	16	5	3	1	0	8	2	2	2
4th Crop	-	-	-	-	0	3	3	27	3	9	1	6	4	2	0	5	-	-	-	-	0	4	0	2	0	7	3	21	0	8	2	0
Totals	9	5	3	5	14	16	3	8	27	1	0	10	32	16	3	8	8	18	0	15	15	16	3	2	23	15	2	12	26	13	3	12
SECOND SEASON, 1862.																																
1st Crop	6	13	1	3	14	12	1	4	15	12	0	7	14	3	0	21	10	19	1	27	13	14	0	8	14	1	0	27	8	14	0	21
2d Crop	1	10	0	7	8	16	3	26	9	13	0	25	7	0	0	15	3	19	3	17	7	0	3	27	6	9	3	6	10	11	2	9
3d Crop	-	-	-	-	3	18	1	9	8	1	0	5	9	15	0	0	1	10	3	9	3	7	3	2	9	16	1	22	8	18	1	20
4th Crop	-	-	-	-	0	10	2	7	1	4	3	10	1	11	1	14	-	-	-	-	0	8	1	11	1	14	3	15	3	2	0	5
5th Crop	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Totals	8	3	1	10	27	18	0	15	34	10	0	19	32	9	2	22	16	10	0	25	27	11	0	20	32	2	1	14	31	12	1	20
THIRD SEASON, 1863.																																
1st Crop	3	18	3	14	10	14	1	0	12	4	3	18	9	16	3	27	6	7	3	24	13	1	3	32	12	9	0	7	8	2	1	27
2d Crop	0	19	3	27	6	7	3	9	9	7	0	22	10	13	3	0	1	12	3	22	5	13	0	6	7	2	1	24	11	12	3	21
3d Crop	-	-	-	-	4	2	3	24	7	2	3	1	6	18	2	5	-	-	-	-	6	1	0	2	7	0	0	12	8	6	1	26
4th Crop	-	-	-	-	0	16	3	18	5	13	0	4	8	11	1	12	-	-	-	-	0	9	1	6	3	9	3	20	6	5	1	3
5th Crop	-	-	-	-	0	3	0	16	0	5	2	10	0	14	1	20	-	-	-	-	-	-	-	-	0	10	0	5	0	11	2	0
6th Crop	-	-	-	-	-	-	-	-	-	-	-	-	0	5	2	24	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Totals	4	18	3	13	22	5	0	11	34	18	1	27	37	0	2	5	8	0	3	19	25	5	1	8	30	11	2	12	34	19	1	21
SUMMARY—1861, 1862, and 1863.																																
1861	9	5	3	5	14	16	3	8	27	1	0	10	32	16	3	8	8	18	0	15	15	16	3	2	23	15	2	12	26	13	3	12
1862	8	3	1	10	27	18	0	15	34	10	0	19	32	9	2	22	16	10	0	25	27	11	0	20	32	2	1	14	31	12	1	20
1863	4	18	3	13	22	5	0	11	34	18	1	27	37	0	2	5	8	0	3	19	25	5	1	8	30	11	2	12	34	19	1	21
Average	7	9	1	9	21	13	1	12	33	3	1	0	34	2	1	12	11	3	0	10	22	17	3	1	26	9	3	13	31	1	3	13

Table II. shows that the crops obtained before the end of May were very much larger in the second and third seasons than in the first season, in which case no sewage had been applied during the winter months. The much greater luxuriance of growth in the early season was, indeed, remarkable after the winter applications; and the crops were invariably in the most forward condition where the largest quantities of sewage had been applied. There is, of course, a great advantage in getting an early cut of green food, and a given weight will be worth more quite early in the season than some weeks later. Still, it is not to be supposed that the same amount of increase of produce will be obtained for a given amount of sewage applied during winter, as during the periods of active growth.

It is also to be observed that the crops obtained late in the season, in September and afterwards for example, were always considerably heavier with sewage than without it, heavier with 6,000 tons than with 3,000, and generally heavier with 9,000 than with 6,000.

Thus, not only was the total amount of produce obtainable per acre very much increased by the application of sewage, but the period during which an abundance of green food was available was extended considerably, both at the beginning and the end of the season, and it was the more so the larger the quantity of sewage applied, almost up to the highest amount adopted in the experiments, namely, 9,000 tons per acre.

Table III. shows that in only one instance, that of the 10-acre field in the wet and cold season of 1862, was there any third cut at all without sewage, whilst, with sewage, four or more cuttings were always obtained, and the later crops were pretty uniformly the larger the larger the quantity of sewage applied.

Leaving the question of the amounts of produce obtained during each separate month, or at each successive cutting, Table IV. shows the total amount of produce on each plot, in each of the three seasons, and also the amount of increase for every 1,000 tons of sewage applied; and produce and increase are each recorded both as green grass, and calculated as hay. The means of the results obtained in the two fields are also given.

TABLE IV.—Total Amounts of Produce, and the Increase for 1,000 Tons of Sewage applied, both reckoned as Green Grass, and calculated as Hay.

SEASONS 1861, 1862, and 1863.

	Five-acre Field.				Ten-acre Field (half).				Mean of the two Fields.			
	Without Sewage.		With Sewage.		Without Sewage.		With Sewage.		Without Sewage.		With Sewage.	
	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.
Produce per Acre, per Annum, as Green Grass.												
1861	9 5 3 5	14 16 3 8	27 1 0 10	82 16 3 8	11 3 0 10	22 17 3 1	23 15 3 12	23 13 3 12	9 1 3 24	15 6 3 5	24 18 1 11	29 15 1 10
1862	8 3 1 10	27 18 0 18	34 10 0 19	32 9 0 20	16 10 0 20	27 11 0 20	32 2 1 14	31 13 1 20	12 6 3 1	27 14 2 19	33 6 1 3	32 1 0 7
1863	4 18 3 13	22 5 0 11	34 13 1 27	37 0 2 5	8 0 3 19	25 5 1 8	30 11 2 12	34 10 1 21	6 9 3 16	23 15 0 24	32 15 0 6	35 19 3 27
Average	7 9 1 9	21 13 1 12	32 3 1 0	34 2 1 12	11 3 0 10	22 17 3 1	28 9 3 13	31 1 3 18	9 6 0 24	22 5 2 7	31 6 3 6	32 12 0 15
Produce per Acre, per Annum, calculated as Hay.*												
1861	2 19 0 17	4 3 1 24	6 3 0 0	7 5 0 27	2 11 1 24	3 15 2 20	4 9 1 6	5 5 1 4	2 15 1 7	3 19 2 8	5 6 0 17	6 5 1 2
1862	2 10 1 23	6 8 2 13	5 19 2 16	6 2 0 11	4 13 3 4	5 17 3 18	5 14 1 5	6 10 1 15	3 12 0 14	6 3 1 1	5 16 3 24	6 6 0 27
1863	3 2 0 10	5 4 0 22	6 2 1 1	7 1 0 13	3 7 2 27	5 0 3 16	6 4 1 25	6 12 3 2	2 14 3 19	5 2 2 6	6 3 1 13	6 16 3 23
Average	2 10 2 7	5 5 1 20	6 1 2 15	6 16 0 17	3 11 0 0	4 18 0 18	5 9 1 3	6 2 3 7	3 0 3 4	5 1 3 5	5 15 1 23	6 9 1 26
Increase, reckoned as Green Grass, for each 1,000 Tons of Sewage applied.												
1861	..	2 10 1 7	4 0 1 9	3 16 2 6	..	4 10 3 23	4 18 3 23	4 4 0 20	..	3 10 2 15	4 9 2 16	4 0 1 13
1862	..	6 11 3 23	4 7 3 16	2 14 0 13	..	3 13 2 20	2 12 0 4	1 13 2 7	..	6 2 3 8	3 9 3 24	3 5 3 10
1863	..	5 15 1 22	4 19 3 21	3 11 1 4	..	5 14 3 1	3 15 0 13	2 19 3 10	..	5 15 0 12	4 7 2 3	3 5 2 7
Average	..	5 2 0 27	4 9 1 15	3 7 1 8	..	4 16 0 15	3 15 1 13	2 19 0 23	..	4 19 0 21	4 2 1 14	3 3 1 1
Increase, calculated as Hay, for each 1,000 Tons of Sewage applied.*												
1861	..	0 12 3 20	0 14 1 21	0 13 3 27	..	0 17 2 8	0 13 2 6	0 12 3 3	..	0 15 1 0	0 14 0 0	0 13 1 15
1862	..	1 6 0 14	0 11 2 5	0 7 3 25	..	0 8 0 4	0 3 1 19	0 4 0 7	..	0 17 0 9	0 7 1 28	0 6 0 2
1863	..	1 0 2 23	0 13 1 13	0 11 0 6	..	0 11 0 6	0 9 1 23	0 7 0 25	..	0 15 3 15	0 11 1 18	0 9 0 13
Average	..	0 19 3 19	0 13 0 13	0 10 3 27	..	0 12 0 25	0 8 3 7	0 8 0 3	..	0 16 0 8	0 10 3 24	0 9 3 1

In the first and third years (1861 and 1863) there was, in both fields, more produce, whether reckoned as green grass or as hay, with each increased amount of sewage applied. In the wet and cold season of 1862, however, although there was considerably more green produce per acre from 6,000 tons of sewage than from 3,000, yet, owing to its more succulent condition, the amounts with the larger quantities of sewage represented even a few cwts. less of hay. On the other hand, with 9,000 tons of sewage, the produce of green grass was, in this wet season, in both fields, less than with 6,000 tons, but calculated as hay it was slightly more. In no case, however, is the increase with the larger amounts of sewage in proportion to the increased application. This point is well illustrated in the two lower sections of Table IV., which give, not the produce per acre, but the increase for each 1,000 tons of sewage applied, reckoned respectively as green grass and as hay.

It is obvious that the proportion of the produce to be reckoned as increase due to the sewage applied must depend very much on the yield of the unmanured land with which the produce of the sewaged land is to be compared. It is necessary, therefore, to bear in mind the quality and condition of the land upon which the experiments were made, when estimating and judging of the amounts of increase yielded for a given amount of sewage applied. Both fields were fattening pastures. It may safely be concluded, therefore, that their natural or unmanured produce would be higher than that of the average of such land as would be likely to be devoted to the growth of grass by means of sewage on the large scale. On the other hand, there is no reason to suppose that under the influence of a liberal supply of sewage, the produce would be in a corresponding degree higher on land of high natural yield than on land which, from its suitable physical qualities, might, with sewage, yield heavy crops, but without it, very light and poor ones. Then, again, a large proportion of the grasses of a good fattening pasture will yield less produce under the influence of sewage than others associated with them, and it is not until the application has been continued for some years that the more freely growing grasses become so far dominant as to secure the maximum result for a given amount of sewage that soil and season will admit of. For these reasons it seems probable that the amounts of *increase* obtained for a given amount of sewage, applied in these experiments, are more likely to be below than above those which may be anticipated on the continued application of sewage, over large areas of land, selected, prepared, and properly seeded for the purpose.

As between the two fields, it should be stated that, to the five-acre field no sewage had been applied during the season preceding the commencement of the experiments; one crop of hay had been taken from it, and it had afterwards been eaten down by sheep. To the ten-acre field, however, sewage had been applied in indefinite quantity for nearly 12 months; two crops of hay had been taken from the land in the previous year, and it had been kept *very closely grazed down* by stock, almost up to the time of

commencing the experiment. Thus, though the land of this field was undoubtedly of higher natural quality, and was probably also in a higher condition, so far as the influence of recent manuring was concerned, yet, owing to its herbage being so much more closely grazed down, it was in that respect in a less favourable condition for the first year's crop, and accordingly, in the first season, gave, without sewage, less produce than the other, though in succeeding seasons it gave much more. Indeed, whilst in the five-acre field the produce without sewage diminished from year to year, being even less in the second year than in the first, notwithstanding the much larger amount of rain, it was in the ten-acre field so very much larger in the second year than in the first, and so very much larger in both the second and third years in that field than in the other, that it was thought, until full inquiry had been made, that there must be some error either in the measurement of the land or in the records. None was, however, found; and the difference in the character and composition of the two soils, which subsequent examination showed, satisfactorily accounted for the great difference in their natural yield. For further information on this point see p. 63.

These few remarks on the character and condition of the land in the two fields will serve as some explanation of the proportionally much greater difference in the amounts of increase over the natural produce from a given amount of sewage, than in the amounts of total produce per acre, where the same amounts of sewage are applied in the different fields, or in different seasons. A few comments only need be made on the results themselves as recorded in the Table.

In the first season the sewage was not applied experimentally until March in the five-acre, and April in the ten-acre field, and hence the amounts of increase of produce yielded had to be reckoned as due to comparatively small quantities of sewage applied. Taking the average result of the two fields, the increase obtained for 1,000 tons of sewage applied, when reckoned as green grass, was rather more when the two larger than when the smallest quantity of sewage was applied per acre, but reckoned at one uniform condition of dryness as hay, it was slightly less with each increased amount of sewage applied.

In the wet and cold season of 1862, which was, of course, favourable for the unsewaged land, and by comparison the less appropriate the greater quantity of sewage applied, the amount of increase for 1,000 tons of sewage, whether reckoned as green grass or as hay, diminished considerably with each increase of sewage applied per acre. And notwithstanding the amounts of total produce per acre with equal quantities of sewage were not very different in the two fields, the amounts reckoned as increase for 1,000 tons of sewage applied were very much less in the ten-acre than in the five-acre field, owing to the much larger produce without sewage in the former.

In 1863 again, though a much warmer and more genial season for the action of sewage, there was still, though in a much less

degree than in 1862, a diminishing proportion of increase for 1,000 tons of sewage, the larger the quantity applied per acre. There was also, owing chiefly to the much larger produce without sewage, much less to be reckoned as increase for each 1,000 tons of sewage applied in the ten-acre than in the five-acre field.

It is worthy of remark, that although the produce per acre without sewage is so much the greater in the ten-acre field than in the other, it is with equal, but especially with the larger amounts of sewage, pretty uniformly the greater in the five-acre field. This result was doubtless partly due to its being better fitted, from its porosity, for sewage irrigation, but partly also to the fact, that whilst it was comparatively flat, allowing the sewage to pass over it more slowly and so to be better absorbed, the ten-acre field was in high ridges, and steeply inclined, rendering it difficult to prevent the water running over it too quickly. This point will be illustrated further on by reference to the comparative composition of the drainage water from the two fields.

Taking the average results of the three years, and the two fields, we have, with sewage applied at the rate of 3,000 tons per acre per annum a produce per acre of a little over $22\frac{1}{2}$ tons of green grass, equal rather more than 5 tons of hay; with 6,000 tons of sewage rather more than $30\frac{1}{2}$ tons of green grass, equal rather more than $5\frac{1}{2}$ tons of hay; and with 9,000 tons of sewage rather more than $32\frac{1}{2}$ tons of green grass, equal about $6\frac{1}{2}$ tons of hay.

The largest quantities of produce reached were those obtained with the largest quantities of sewage (9,000 tons per acre per annum), and in the third year of the experiments, amounting in the five-acre field to 37 tons of green grass, equal rather more than 7 tons of hay, and in the ten-acre field to nearly 35 tons of green grass, equal nearly 6 tons 13 cwts. of hay.

The average increase of green grass over the natural produce for 1,000 tons of sewage applied was, with 3,000 tons of sewage per acre nearly 5 tons, with 6,000 tons of sewage rather more than 4 tons, and with 9,000 tons not quite $3\frac{1}{2}$ tons. Reckoned as hay, the average increase for 1,000 tons of sewage was, with 3,000 tons of sewage per acre 16 cwts., with 6,000 tons nearly 11 cwts., and with 9,000 tons $9\frac{1}{2}$ cwts. As, however, these average results relating to increase include those of the ten-acre field, where, owing to the very high natural produce, the amount reckoned as increase due to sewage was comparatively small, it is probable that results equal at any rate to those of the five-acre field may be expected in the average of cases elsewhere; and where, as may frequently happen, a soil which yields a very small natural produce, may, nevertheless, owing to its physical qualities, be well adapted for the application of sewage and give large amounts of produce per acre under its influence, the amounts of increase for a given amount of sewage applied may be even considerably higher than those obtained in the five-acre field.

The general result is, that there was much more total produce per acre with 6,000 tons of sewage than with 3,000, and more still

with 9,000; but that the increase for a given amount of sewage applied was less with 9,000 tons than with 6,000, and less with 6,000 than with 3,000.

The increase in the amount of produce with each increase in the quantity of sewage applied appears proportionally greater when reckoned as green grass than as hay. This is due to the much greater succulence, and, therefore, less proportion of dry substance in the more highly sewaged and heavier crops. The question arises, whether, with a less proportion of dry substance in the sewaged grass, a given weight of that dry substance will have a greater or a less value as food for stock than an equal weight from the less succulent unsewaged grass? This point will be fully considered in subsequent Sections of the Report.

II. *Experiments with Italian Rye-grass.*

In April 1863 arrangements were made with Mr. Campbell for gauging the sewage applied, and weighing and sampling the produce obtained, in a field of Italian rye-grass, and also for trying the feeding qualities of the grass. From the field in question, a crop of tares, which had been manured with farm-yard dung, had been carried off in the spring of 1862. The land was then cleaned, again manured with stable and farm-yard dung, and sown down with rye-grass in September (1862); and, at the time of commencing the experiment in the following Spring, there was a promising and tolerably even crop.

Three plots of about an acre each were set apart; plot 1 to be unsewaged; plot 2 to receive sewage at the rate of 3,000; and plot 3 at the rate of 6,000 tons per acre per annum. So meagre was the flow, however, that up to the end of October (1863) only 787 tons had been applied to plot 2, and 1,522½ tons to plot 3, instead of 1,512 and 3,057 tons, respectively, that were required according to the rates intended, reckoning from the date of the first application in the Spring.

The particulars of the amounts of sewage applied, and of the amounts of produce and increase obtained, reckoned both as green grass and in the condition of dryness of hay, are given in Table V., p. 17. Further details will be found in Appendix, Table V., pp. 97-100.

TABLE V.—Amounts of Sewage applied, and of Produce and Increase obtained, in Experiments on Italian Rye-grass.

SEASON 1863.

	Without Sewage.	With Sewage.	
	Plot 1.	Plot 2.	Plot 3.
Sewage applied per Acre.			
April - - -	..	<i>Tons.</i> ..	<i>Tons.</i> 48·0
May - - -	..	152·1	237·6
June - - -	..	178·0	354·7
July - - -	..	218·1	403·1
August - - -	..	120·8	163·4
September - -	..	60·1	219·9
October - - -	..	58·1	75·0
Total - - -	..	787·2	1522·6

Rye-grass obtained per Acre, during each separate Month.

	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>
April - - -	3 4 1 21	4 3 1 23	3 15 3 15
May - - -	1 17 2 23	- - - -	1 11 1 7
June - - -	8 18 0 22	7 1 1 27	6 9 3 13
July - - -	- - - -	5 16 3 21	6 10 2 21
August - - -	2 0 1 13	1 14 1 13	3 0 2 10
September - -	0 9 2 7	0 13 1 6	1 12 0 11
October - - -	0 6 0 9	1 5 3 16	2 3 0 12
	16 16 0 19	20 15 1 21	25 3 2 5

Rye-grass obtained per Acre, in each successive Crop.

	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>
1st Crop - - -	5 2 0 16	4 3 1 23	3 15 3 15
2d Crop - - -	8 18 0 2	7 1 1 27	5 14 0 3
3d Crop - - -	2 0 1 13	5 16 3 21	7 7 1 13
4th Crop - - -	0 9 2 7	1 14 1 13	4 2 2 23
5th Crop - - -	0 6 0 9	0 13 1 6	2 0 1 23
6th Crop - - -	- - - -	1 5 3 16	2 3 0 12
	16 10 0 19	20 15 1 21	25 3 2 5

Summary of Produce per Acre.

	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>
Green Grass - -	16 16 0 19	20 15 1 21	25 3 2 5
Reckoned as Hay*	4 18 3 8	5 5 0 16	5 12 2 11

Increase of Produce per Acre.

	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>
As Green Grass -	..	3 19 1 2	8 7 1 14
Reckoned as Hay*	..	0 6 1 8	0 13 3 3

Increase for each 1,000 Tons of Sewage applied.

	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>	<i>Tons cuts. grs. lbs.</i>
As Green Grass -	..	5 0 2 25	5 9 3 17
Reckoned as Hay*	..	0 8 0 4	0 9 0 5

* The amount of hay to which the grass is equivalent is calculated by raising the amount of the experimentally determined perfectly dry or solid substance in the grass, in the proportion of from 84 to 100, on the assumption that the hay would contain 84 per cent. of dry substance and 16 per cent. moisture.

When the experiment with rye-grass was determined upon early in April, the grass was so far forward that it was found necessary to take a first cutting without sewage, before the

water-runs could be properly adapted for the separate irrigation of the respective plots, and hence but little sewage was applied before the end of April, and that little only on plot 3. The effect of the sewage was, therefore, as the second and third divisions of the Table show, to increase the produce chiefly during the later months and later crops of the season, and it did so very much in proportion to the amounts applied.

The total produce per acre was, without sewage (though otherwise pretty well manured), rather more than $16\frac{1}{4}$ tons, with 787 tons of sewage rather more than $20\frac{1}{4}$ tons, and with 1,522 $\frac{1}{2}$ tons of sewage nearly $25\frac{1}{4}$ tons of green grass; or, reckoned at a uniform condition of dryness as hay, the amounts were equivalent to 4 tons $18\frac{1}{4}$ cwts., 5 tons $5\frac{1}{4}$ cwts., and 5 tons $12\frac{1}{4}$ cwts. respectively. The increase of produce per acre was, therefore, nearly 4 tons of green grass due to the smaller, and about 8 tons $7\frac{1}{4}$ cwts. due to the larger application of sewage; though the increase in real dry substance represented only $6\frac{1}{4}$ cwts., and $13\frac{3}{4}$ cwts. of hay, respectively.

The increase reckoned for 1,000 tons of sewage in each case was, with the smaller quantity applied, 5 tons $0\frac{1}{4}$ cwts., and with the larger quantity, nearly 5 tons 10 cwts. of green grass; but the increase of real dry substance represented only 8 cwts., and 9 cwts. of hay, respectively. The increase in real dry or solid substance was, therefore, very small; but it will be seen further on that, at any rate in the case of the meadow grass (and it is probably the same with the rye-grass), a given amount of the dry substance of the sewaged produce was more productive of milk, and even slightly more of increase, than an equal amount of the dry substance of the unsewaged.

The general result is, that there was as much or more increase of green produce for 1,000 tons of sewage with the rye-grass than in most of the cases in the same season with the meadow-grass, where so very much larger quantities of sewage were applied, though the increase of dry substance reckoned as hay was generally the higher with the meadow-grass. That is to say, the comparatively large amounts of sewage applied to the meadow-grass gave, on the average, a larger amount of increase in dry or solid substance, for a given quantity of sewage, than the much smaller amounts applied to the rye-grass. It is also to be observed that there was a larger amount of increase, both of green grass and of dry substance reckoned as hay, for a given quantity of sewage on plot 3 with the larger, than on plot 2 with the smaller amount applied to the rye-grass. The facts point to the conclusion that, for the season in question, the larger quantity applied was below that required to yield the maximum increase for a given amount of sewage. It is obvious, however, that it may be advantageous even to pass this point; for, within certain limits, it will be economical to reduce the area and cost of distribution at the expense of a certain sacrifice of sewage.

It is to be regretted that the plant of rye-grass was so much injured by frost during the winter of 1863-4 (and it was the more

To face page 19.

TABLE VI.—Summary of the Results of

Plots, &c.	Consumed per Head per Day.				Consumed per 1,000 weight per		calculated sewage applied.
	Fresh Food.		Dry Substance of Food.		Fresh Food.		Value of Increase in Live-weight from the increased Produce from 1,000 tons Sewage.
	Grass.	Oilcake.	In Grass.	In Oilcake.	Grass.	Oilcake.	Including the Oilcake consumed (if any) -
							Reckoning of Oilcake
Season							
1. Unsewaged	Lbs. 89'8	Lbs. ..	Lbs. 23'7	Lbs. ..	Lbs. 576	Lbs. ..	£ s. d.
2. Sewaged	105'2	.	21'3	..	668
3. Sewaged							..
4. Sewaged							..
Season 1862 ; Reckoning							
1. Unsewaged	105'4	3'5	23'6	3'1	584	19'7	..
2. Sewaged	126'1	3'5	20'9	3'1	704	19'9	3 8
3. Sewaged							2 6
4. Sewaged							1 8
Season 1862 ; Reckoning							
1. Unsewaged	100'9	3'7	23'9	3'3	535	19'5	..
2. Sewaged	123'2	3'7	20'9	3'3	668	19'7	2 12
3. Sewaged							1 15
4. Sewaged							1 1

* The value of the increase in live-weight, "exclusive of oilcake," is reckoned by weight, "exclusive of oilcake," by deducting from the gross increase with oilcake a very much larger proportion of the cost was chargeable against the manure, and
† This period includes the first three weeks during which the two oken, otherwise
‡ This period excludes the first month of the experiment, during three weeks of

re the sewage had been the most liberally applied), that it necessary to plough it up, otherwise the experiment would been continued through the season of 1864. Instead of this, was sown over the three plots without any further manure, was obvious to the eye during growth that the crop was heavier where sewage had been applied to the rye-grass than it had not, and heavier where the larger than where the quantity had been applied.

III. *Experiments with fattening Oxen.*

in 1861, so in 1862, 10 oxen were purchased and tied up in to consume the grass from the five acre field; two to be unsewaged, and the remaining eight on sewaged grass; the to be cut, as ready, indiscriminately from the three sewaged. In 1861, the animals had grass alone for the first 16 out 20 weeks of the whole experiment, and they had oilcake in (four pounds per head per day) only during the concluding weeks. The object was to try grass alone in the first season, the result was very unfavourable. In 1862, oilcake was given in addition to the grass from the commencement, in quantity which, fed over the whole period of nearly 23 weeks, amounted to 3½ lbs. per head per day.

The detailed records of the experiments in 1861 were given in appendix to the last Report, and those of the experiments of will be found in Tables VI. to VIII., pp. 101-108, in the Appendix to the present Report. Table VI. (opposite) summarises, and together at one view, the results obtained in 1861 on grass and those in 1862, on grass with oilcake in addition. The division of the Table relates to the experiments of 1861; middle one to the whole period of experiment of 1862, giving the first four weeks during three of which the oxen fed on unsewaged grass, had (for want of supply) sewaged and the lower division represents the results exclusive of that month. Inasmuch as both lots increased very much more in the first month than afterwards, the rate of increase on the whole would appear comparatively small if that period were excluded; whilst, including it, included also the period of three weeks during which the two oxen had sewaged grass. It was better, therefore, to give the results both ways. The comparison between the effects of the unsewaged and the sewaged grass is, however, much the same whichever period be adopted.

In both years a greater weight of the fresh sewaged grass was consumed per head per day, and per 1,000 lbs. live-weight per day than of the less succulent unsewaged grass; but the dry solid substance contained in the larger amount of sewaged consumed was less than that in the unsewaged. Again, as in 1861, grass was given alone, more of the sewaged than of the unsewaged, reckoned in the green or fresh state, required to produce 100 lbs. increase in live-weight; though

the amount of dry substance contained in the sewaged grass so required was only about four fifths as much as that in the unsewaged grass. But when, as in 1862, a fair allowance of oilcake was given in addition, very much less both of fresh food and of dry or solid substance of food were required to produce 100 lbs. increase in live-weight than in 1861 with grass alone, and considerably less of the dry or solid substance of the more succulent sewaged than of the drier unsewaged grass was required.

It is also observable, that, reckoned in the green state, about the same amount both of the unsewaged and sewaged grass was consumed per 1,000 lbs. live-weight per week in 1862 with oilcake in addition, as in 1861 with grass alone; but the dry substance supplied in the grass consumed in 1862, with oilcake, was, both with the unsewaged and the sewaged grass, less than in 1861 without it.

The result in 1861, when cut grass was given alone, was extremely unsatisfactory, the amount of food required to produce a given amount of increase being unusually large, and the rate of increase on a given weight of animal within a given time unusually small. But when, in 1862, oilcake was given with the grass, and especially when given with the sewaged grass, very much better results were obtained. Indeed, in 1862, the rate of increase per 1,000 lbs. live-weight per week (if taken over the whole period of nearly 23 weeks) was, both with unsewaged and with sewaged grass (and oilcake in addition), about equal to the average obtained with animals of fair quality fed on good fattening food; but the food consumed for the production of 100 lbs. of increase, even in the case of the sewaged grass, contained more dry or solid substance than is usually required when oxen are liberally fed on oilcake, hay-chaff, and roots, and with the unsewaged grass considerably more. It should be borne in mind, however, that the experiment with unsewaged grass was on two animals only, whilst that with the sewaged was on eight, giving, therefore, a much more trustworthy average; and the results given in Appendix, Table VII. p. 107, show that one of the two oxen on unsewaged grass gave less increase than any of those on the sewaged, whilst the other gave considerably more than the average increase of the latter.

The general result is that the sewaged grass cut green and given to oxen tied up under cover, gave, when supplemented with a fair allowance of oilcake, a good average rate of increase in relation to the weights of the animals within a given time, and also a moderate rate of increase in relation to the amount of dry or solid substance provided in the food consumed.

It remains to say a few words on the last ten columns of Table VI.

It is seen that, by the aid of sewage, the time which an acre of land would provide food for an ox was increased three or more fold, varying according to the amount of sewage employed. Taking into account, however, the large amounts of oilcake consumed with the produce of each acre in 1862, it results

that (excepting on plot 2, where the produce was very much larger than in 1861) a given area would support considerably less stock in the cold and wet season of 1862 than in the more genial one of 1861.

The amount of increase in live-weight yielded from the produce of an acre was also increased several fold by means of sewage; about three-fold with the highest amount of sewage when the grass was consumed alone, and nearly four-fold in 1862, when oilcake was given in addition, in much about the same proportion to a given amount of the unsewaged and the sewaged grass.

It was shown in the last Report how very small was the gross money value of the increase in live-weight obtained from the consumption of the produce of an acre, or of the increased produce from 1,000 tons of sewage, when the grass was consumed alone, and the results then referred to are given in the upper division of the Table to compare with those in the lower divisions relating to 1862, when oilcake was also used. The result is seen to be, that the money value of the increase in live-weight from the produce of an acre of sewaged land, or from the produce of 1,000 tons of sewage, was very much greater in 1862, when oilcake was given, than in the corresponding cases in 1861 without it. The money return per acre was also from three to four times as great with sewage as without it, and although it is greater where 9,000 or 6,000 than where only 3,000 tons of sewage were applied, yet the return calculated, not per acre, but for each 1,000 tons of sewage, is, in each case, the less the greater the amount applied.

The next section of the Report will show that a very much higher money value was obtained both from an acre of land, and from a given amount of sewage, when the sewaged grass was employed for the production of milk instead of meat. But it may be mentioned that at Croydon, although the land is there more liberally sewaged than was the case in any of the Rugby experiments, satisfactory results have been obtained with fattening stock fed on the land. The practice there is, to irrigate for three or four days and nights together two or three times for each crop, and when the grass has got a sufficient head, to stop the application and turn the stock upon the land, where they remain until the grass is closely eaten down; they are then removed, the land is re-irrigated, and so on.

IV. *Experiments with Milking Cows.*

By the kindness of Mr. Campbell, experiments were made with his cows each year, 1861, 1862, and 1863, on the milk-yielding qualities of the grass.

In 1861, 12 of Mr. Campbell's cows were carefully selected and set apart to be fed on grass alone, 2 on unsewaged and 10 on sewaged grass, and the experiment was so conducted over a period of 16 weeks. It was afterwards continued for 4 weeks longer, with an allowance of oilcake as well as grass.

In 1862, 3 cows were selected to receive oilcake and unsewaged, and 12 oilcake and sewaged grass, and the experiment was continued for 24 weeks.

In 1863, 20 recently calved cows were selected, 5 to be fed on unsewaged meadow grass, 10 on sewaged meadow grass, and 5 on Italian rye grass. The design was to give each lot grass alone for the first 12 weeks, and afterwards a certain amount of oilcake in addition.

The detailed records of the experiments with cows in 1861 were given in the Appendix to the last Report; and those of the experiments in 1862 and 1863 will be found in Tables IX. to XIV., pp. 109-162, in the Appendix to the present Report. The results of all the experiments are given, in a condensed form, at one view, in Table VII. opposite.

Leaving the more detailed Tables for reference, to supply any further illustrations or explanations that may be needed, it will be sufficient to make a few comments on the main facts brought to view in the last-mentioned summary Table.

Reviewing the results of the experiments in which sewaged was tried against unsewaged meadow grass, it is observable that, excepting in the first season (1861), the cows required more both per head per day, and per 1,000 lbs. live weight per week, of the fresh or green sewaged than of the unsewaged grass; yet, the yield of milk, both per head and per 1,000 lbs. live weight, was, without exception, the greater with the unsewaged grass. The increase in live weight was also somewhat the greater on the unsewaged grass in 1861 and 1862, but the contrary was the case in 1863.

Reckoned in the fresh or green state in which it was cut and carted, there was, in fact, in every case but one (and then the quantities were equal), considerably less of the unsewaged than of the sewaged grass required to be consumed for the production of one gallon of milk. It should be remarked, however, that the unsewaged grass was generally cut in a much riper and less succulent condition, and therefore contained a considerably higher per-centage of dry or solid substance than the sewaged. It may be also here mentioned that in 1863 the cows having professedly unsewaged meadow grass, in default of a sufficient supply of it, had necessarily for a considerable part of each of the periods of 12 weeks unsewaged rye-grass.

Weight for weight, in the fresh or green state in which the grass was cut, weighed, and given to the cows, the unsewaged grass has, therefore, proved to be far more productive than the sewaged. But when the comparison is made, not between the amounts of grass reckoned in the fresh state, but between the amounts of dry or solid matter which the different descriptions of grass supplied, the result is that, in only one instance was there more, and in the others there was either an equal amount or even less of dry or solid substance of sewaged than of unsewaged grass required for the production of a given amount of milk.

The general result in regard to these points was, then, that in both milk and increase, but, especially milk, a given weight of

TABLE VII.—SUMMARY OF THE RESULTS OF THE EXPERIMENT WITH RYE GRASS (Unsewaged and Sewaged),

PLOTS, &c.	Consumed per Acre.†		Results calculated per 1,000 tons Sewage applied.†							
	Fresh Food.		Value of the Milk from the Produce of each Acre at 8d. per Gallon.		Amount of Milk from the increased Produce from 1,000 tons of Sewage.		Value of the Milk from the increased Produce from 1,000 tons of Sewage at 8d. per Gallon.			
	Grass.	Oil-cake.	Including the Oilcake consumed (if any).	Exclusive of Oilcake.‡	Including the Oilcake consumed (if any).	Exclusive of Oilcake.‡	Including the Oilcake consumed (if any).	Exclusive of Oilcake.‡		
1. Unsewaged	Lbs.	Lbs.	s.	£ s. d.	£ s. d.	Gallons.	Gallons.	£ s. d.	£ s. d.	
2. Sewaged	150·2	19 14 3	
3. Sewaged	124·0	19 0 6	..	179½	..	5 19 10	
4. Sewaged			27 6 11	..	178	..	5 18 8	
			32 0 10	..	151½	..	5 0 11	
1. Unsewaged	127·2	3·5	..	22 4 5	20 8 10	
2. Sewaged	30 13 8	27 16 10	84½	74	2 16 5	2 9 4	
3. Sewaged	140·2	3·6	..	35 15 2	32 8 11	67½	60	2 5 2	2 0 0	
4. Sewaged			..	35 4 1	31 18 10	43½	38½	1 8 10	1 5 7	
1. Unsewaged	99·1	13 8 0	
2. Sewaged	33 19 5	..	205½	..	6 17 2	
3. Sewaged	142·9	46 16 7	..	167½	..	5 11 5	
4. Sewaged			51 9 5	..	126½	..	4 4 7	
1. Unsewaged	90·5	3·9	..	14 16 3	14 7 6	
2. Sewaged	26 0 0	25 4 2	111½	109½	3 14 7	3 12 3	
3. Sewaged	172·0	3·7	..	35 16 9	34 14 11	105½	101½	3 10 1	3 7 11	
4. Sewaged			..	39 7 9	38 3 10	81½	79½	2 14 7	2 12 11	
Grass and Oilcake.										
1. Unsewaged	95·4	1·7	..	13 19 7	13 16 0	
2. Sewaged	29 12 4	29 3 9	156½	153½	5 4 3	5 2 7	
3. Sewaged	157·4	1·8	..	40 16 6	40 4 7	134½	132½	4 9 6	4 8 1	
4. Sewaged			..	44 17 5	44 4 4	103	101½	3 8 8	3 7 7	
1. Unsewaged	159·3	24 0 0	
2. Sewaged	29 13 3	..	215½	..	7 3 11	
3. Sewaged			35 19 0	..	235½	..	7 16 11	
1. Unsewaged	121·3	3·3	..	28 5 6	27 11 4	
2. Sewaged	34 18 11	34 1 5	254½	247½	8 9 6	8 5 3	
3. Sewaged			..	42 7 1	41 5 11	277½	270½	9 4 11	9 0 3	
Rye Grass and Oilcake.										
1. Unsewaged	142·5	1·5	..	25 12 2	25 6 10	
2. Sewaged	31 13 0	31 6 5	230½	227½	7 13 5	7 11 11	
3. Sewaged			..	38 7 3	37 19 3	251½	248½	8 7 5	8 5 9	

* In the preparation of this Table the animals received any other than regular throughout the seasons as the point of fact the allowance, at parallel.

† In the experiments with meadow increase per 1,000 tons of sewage, in the two fields.

‡ The value of the milk, "exclusive of oilcake," and the amount of milk, "exclusive of oilcake," by deducting from the gross amount of milk with oilcake and cotton-cake, were used, a very much larger proportion of the cost was chargeable against the milk.



animal was more productive when fed on unsewaged than on sewaged grass, and that a given weight of fresh unsewaged grass was more productive than an equal weight of fresh sewaged grass; but that a given weight of dry or solid substance supplied in sewaged grass was more productive than an equal weight supplied in unsewaged.

A careful consideration of the results leads to the conclusion that there was some considerable variation in the quality of the grass in the three different seasons. It was obviously very inferior in the wet and cold season of 1862. There is also evidence of a considerably diminished productiveness of a given weight both of green sewaged grass and of the dry substance of sewaged grass towards the end of the season; though part of the falling off which the figures show is doubtless attributable to the changing condition of the cows themselves as the season advanced.

The experiments do not afford the means of strictly comparing the productive qualities of rye grass with those of meadow grass, or of sewaged with those of unsewaged rye grass. Thus, as already alluded to, the cows professedly fed on unsewaged meadow grass in 1863 had, during a considerable part of the experimental period, unsewaged rye grass; whilst those fed on rye grass had indiscriminately the unsewaged and the sewaged. The indication is, however, that somewhat more of the dry substance of the sewaged rye grass than of the sewaged meadow grass was required to produce a given result; though the difference is less during the later than the earlier period of the season. It is probable, indeed, that sewaged Italian rye grass deteriorates less towards the end of the season than sewaged meadow grass.

It remains to indicate, approximately, the increased yield of saleable produce, and the money value of that produce, from an acre of land, and from 1,000 tons of sewage, according to season, and to the amount of sewage applied.

The last ten columns of Table VII. refer to these points. In explanation of the figures there given it should be stated, that the estimates of the amount, and value, of the milk yielded per acre, are, in the case of each plot, based upon the total amounts of grass obtained per acre throughout the season, and upon the average rate of consumption and yield of milk during each separate period, on unsewaged grass in the case of plot 1., and on the mixed sewaged grass in that of plots 2, 3, and 4; and the estimates are framed so as to show, as far as practicable, the amount and value, both inclusive and exclusive of oil-cake when it was given, as will be better understood by reference to the columns in the Table, and the foot note relating thereto.

It is obvious that such estimates can only be approximations to the truth. But such they are, and considered as such only, they are little likely to mislead any acquainted with practical agriculture, and with the limits within which such calculations are and are not of general application.

Referring first to the experiments with meadow grass, the result (excluding the case of 1862, when the unsewaged crop in the ten-

acre field was so very large) was, that the produce of an acre without sewage was competent to feed one cow for from 19 to 23 weeks, varying according to the season, or whether the grass were consumed alone or with oil-cake in addition. The same area was, with the aid of 3,000 tons of sewage, rendered capable of providing keep for two to two and a half cows for the same period of time; or, as represented in the Table, for one cow from two to two and a half times as long, with 6,000 tons of sewage for three to three and a half times as long, and with 9,000 tons from three and a half to four times as long.

Represented in quantity of milk, in 1861, when the sewage was not applied until the Spring, the produce per acre was, without sewage, 321½ gallons, and with the different amounts of sewage 570½, 820½, and 961½ gallons, respectively. Reckoned according to the rate of consumption of grass and of the yield of milk during the first 12 weeks, or most favourable period, of the grass season of 1863, when, as in 1861, the grass was consumed alone, but unlike 1861 the sewage had been applied throughout the winter months, and when the cows being mostly newly calved were also in their most favourable condition, the estimated yield of milk reckoned upon the total produce of grass per acre was, without sewage 402 gallons, and with the different amounts of sewage 1,019, 1,404½, and 1,544 gallons, respectively; or, so far as the sewaged plots were concerned, from one-half to two-thirds more per acre than in 1861 reckoned according to the rates of consumption and yield of milk over the whole of that season.

With the aid of the large quantities of oilcake stated in the Table, the yield of milk per acre in 1862, when the season was very favourable for the unsewaged but comparatively unfavourable for the sewaged land, was, without sewage 666½ gallons, and with the different amounts of sewage 920½, 1,072½, and 1,056 gallons; and, according to the rates of consumption and yield of milk when oilcake was given during the latter half of the season of 1863, the yield of milk per acre calculated upon the total produce of grass throughout the season in each case was, without sewage 444½ gallons, and with the different amounts of sewage 780, 1,075, and 1,181½ gallons respectively.

The general result is, that, on the sewaged plots, the yield of milk was at a less rate per acre with oilcake during the latter or inferior part of the season of 1863, than without it during the earlier or more favourable portion of the same season; but it was at much the same rates per acre during the latter or inferior part of the comparatively favourable season of 1863, as during the entire period of the unfavourable season of 1862. Lastly, on this point, the yield of milk per acre over the entire season of 1863, half without and half with oilcake, was higher than that of 1862 with oilcake throughout, and considerably so on the more highly sewaged plots.

So far as may be judged from the limited experience which these results record, it would appear probable that with an average supply of about 5,000 tons of sewage per acre per annum to meadow

and, and with cows taken indiscriminately at various periods after milking, an average of not less than 1,000 gallons of milk per acre might be expected; or more than this when cows are taken at their best, and the season and other circumstances are more than usually favourable.

In the case of the experiments with rye-grass much less sewage was applied than is above assumed, and the results relate to the experience of one season only, which was not only a comparatively favourable one for the action of sewage, but, being the first year of growth after sowing down, was also favourable so far as the condition of the crop was concerned. The indication is, however, that with Italian rye-grass a larger yield of milk per acre may be obtained, for the application of a given amount of sewage, than with meadow grass. But with Italian rye-grass the land has to be periodically broken up, during which time less sewage per acre, if any, can be utilized, and hence, for the distribution of a given amount of sewage, the expense of laying down much larger area would be necessary so far as this crop were introduced. On the other hand, the advantage of the practice would be, that other crops, for which the direct application of sewage is less appropriate than to grass, would be intermediately grown, either relying upon the residue of sewage manuring remaining in the broken up land, or by means of the solid manure derived from the consumption of the sewaged grass. In either case, therefore, such produce would be obtained indirectly by means of sewage.

Bearing in mind the varying conditions of the several experiments, the differences which the Table shows in the estimated value of the milk yielded from the produce of each acre ostensibly receiving the same amount of sewage will be intelligible, and it will not be necessary to call attention to the figures in detail. The results taken as a whole lead to the conclusion that the gross money return per acre, reckoned in milk at 8d. per gallon, might be estimated at certainly not less than 30l. to 35l. with an application of about 5,000 tons of sewage per acre per annum.

No allowance can be made for the value of the increase in weight in the case of cows, for at the end of their milking period, even though some may gain in weight considerably, they are certainly, on the average, of less money value than at the beginning, so that a deduction rather than an addition should be made on the score of the difference in value of the animals themselves, at the end as compared with the beginning of their milking period.

The last four columns of Table VII. show the estimated yield of milk, and the value of the milk at 8d. per gallon, from the increased produce of 1,000 tons of sewage in the different experiments; and the estimates are given both inclusive and exclusive of oilcake (if any) as before referred to.

In 1861, when the sewage was not applied until the Spring, and less had therefore to be reckoned as contributing to the increase obtained, nearly 180 gallons of milk are estimated to be

obtained for each 1,000 tons of sewage, when the amount actually applied had not exceeded 3,000 tons, and the rate per annum not 5,000. Again, reckoned according to the yield of milk for a given amount of grass consumed during the first 12 weeks, or the most favourable half, of the season of 1863, the increased yield of milk for each 1,000 tons of sewage was 205½ gallons with 3,000, 167 gallons with 6,000, and only 126½ gallons with 9,000 tons of sewage per acre per annum. According to the results obtained with rye grass during the same 12 weeks of the season of 1863, the yield was even somewhat better than in the most favourable case with the meadow grass, and it is the better the larger the quantity of sewage, which, however, was not applied until the Spring, and, owing to deficiency of supply, little exceeded 1,500 tons per acre.

Owing to the very large produce without sewage in 1862, and to the unfavourable character of the season for the action of sewage, the increased yield of milk estimated as due to 1,000 tons of sewage, even with the addition of a considerable amount of oilcake, was very small; much smaller, indeed, than in the more favourable seasons without the oilcake. It was also smaller with oilcake during the latter half of the season of 1863, than without it during the former half, when the quality of the grass, and the condition of the cows, were both so much more favourable. Even taking the whole of the comparatively favourable season of 1863, half without and half with oilcake, but when winter supply of sewage had to be reckoned against the produce, the estimated increased yield of milk for each 1,000 tons of sewage applied was considerably less than in 1861 without oilcake, but when no winter supply of sewage had to be reckoned against the yield.

With rye-grass, unlike the meadow grass, the yield of milk for a given amount of sewage was greater during the latter part of the season with oilcake, than during the earlier or more favourable part without it. As before observed, it would appear that rye-grass does not deteriorate so much as meadow grass as the season advances.

In 1861, when no winter supply of sewage had to be reckoned against the increase, and the grass was consumed without oilcake, the estimated value of the milk obtained for each 1,000 tons of sewage applied was between 5*l.* and 6*l.* In 1863, according to the yield of milk for a given amount of grass during the first half of the season, without oilcake, the value for each 1,000 tons of sewage was, with 3,000 tons more, but with 6,000 and 9,000 tons less, than in 1861; and, reckoned according to the yield during the latter half of the season, it was, even with oilcake, little more than half as much as according to that during the earlier half; and in the unfavourable season of 1862, also with oilcake, it was even less still. Taking the whole of the comparatively favourable season of 1863, half without and half with oilcake, the return for each 1,000 tons of sewage, reckoned in milk at 8*d.* per gallon, was rather over 5*l.* when only 3,000 tons were applied, one seventh less when 6,000 tons, and one third less when 9,000 tons were employed.

With the rye-grass the estimated money return, for a given amount of sewage applied, was considerably higher than with the meadow grass; but it must be borne in mind that there was no winter supply of sewage to reckon against the produce, and that the results relate both to the first and most productive year of the growth of the crop, and to a favourable season for the application of sewage.

Excepting in the cases of the rye grass, and of the meadow grass in 1861, in neither of which had there been any winter supply, there was a very marked diminution in the money return for a given amount of sewage where the largest quantities were applied. The practical question suggests itself—what is approximately the limit of maximum yield for a given amount of sewage which is attainable without so far increasing the area, and consequently the cost of distribution, as to more than counter-balance the increased return? Special reference will be made to this point further on. But it may be here observed that, so far as these results give the means of judging, it would appear that an average of about 5*l.* increased value of milk, reckoned at 8*d.* per gallon, may be anticipated from the application of each 1,000 tons of sewage when the amount applied does not exceed about 5,000 tons per acre, per annum. This would be equivalent to a gross value of increased produce of milk of rather more than 1*d.* for every ton of sewage applied.

V. *Composition of the Rugby Sewage Water.*

From the commencement of the Rugby experiments, samples of sewage have been collected for analysis at short intervals in each of the two fields. The plan was, to take about a quart from the gauge tank in the field, holding about 3½ tons, at intervals of about two hours for as many days in the week as the sewage was applied; and at the end of the week, after well shaking the mixture, a one or two gallon sample was sent to Professor Way for analysis.

In 1861, twelve such samples from each field, representing the supply for the seven months of April to October inclusive, were taken and submitted to analysis.

The season of 1862 is reckoned to include the months from November 1861 to October 1862 inclusive. During each month from November 1861 to April 1862 inclusive only one sample was taken from each field, during May two, during June one, and during July, August, September, and October, two were taken; thus making a total of 17 samples from each field to represent the sewage supplied during the season.

In like manner the season of 1863 was reckoned to include the period from November 1862 to October 1863 inclusive. From November 1862 to March 1863 inclusive two samples of sewage were taken in each month in each field; in April one only in each field, and in May two in the 5-acre, and one only in the 10-acre field. From this date, namely, during June, July, August, September, and October, a different plan of collection

was adopted. During two weeks in each of these months (as a rule the first and third) a sample was taken every two hours from the gauge-tank in whichever field the sewage was being applied, and at the end of the week the samples from the two fields were well mixed, and a portion of the mixture sent off for analysis, as representing the sewage of that week, without distinction as to the field in which it was collected.

The results of each of the 24 analyses of the sewage representing the season ending October 1861 were given in the Appendix (and a summary of them was given in Table VI.) in the previous Report; those of the 34 for the season ending October 1862 are given in Tables XV. to XVII., and those of the 35 relating to the season of 1863 ending October 1863, in Tables XVIII. to XXI., pp. 163-169, in Appendix No. 1. in the present Report.

Leaving the details for reference it will be sufficient to give in this place the summary view of the composition of the sewage which Tables VIII. and IX. (pp. 29 and 30) present; in the former of which is given the average grains per gallon, and in the latter the average lbs. per 1,000 tons, of the different constituents, in the sewage from each field, in each of the three seasons, respectively.

TABLE VIII.—Average Composition of the Rugby Sewage, supplied in each Field, in 1861, 1862, and 1863
GRAINS PER GALLON.

CONSTITUENTS.	Season 1861: April to October.		Season 1862: November 1861 to October 1862.		Season 1863: November 1862 to October 1863.		General Means of			
	5-acre Field; 12 Samples.	10-acre Field; 12 Samples.	5-acre Field; 17 Samples.	10-acre Field; 17 Samples.	5-acre Field; 23 (1) Samples.	10-acre Field; 23 (2) Samples.	24 Samples; April to October 1861.	24 Samples; November 1861 to Oc- tober 1862.	25 (1) Samples; November 1862 to Oc- tober 1863.	25 Samples; November 1861 to Oc- tober 1863.
Organic matter	10.26	10.30	8.48	7.98	8.35	8.29	10.28	8.20	7.93	8.63
	16.75	11.57	16.71	16.96	27.35	25.99	14.16	16.84	24.03	18.85
	27.01	21.87	25.13	24.94	35.70	34.28	24.44	25.04	31.96	27.48
Inorganic matter	36.83	35.85	35.00	35.83	39.57	38.77	36.34	34.43	36.50	35.81
	16.18	12.55	21.31	20.42	39.41	34.93	14.36	20.86	34.45	24.30
	53.00	48.40	56.31	54.25	78.98	73.70	50.70	55.28	71.25	60.11
Total in solution	47.06	46.15	43.43	41.81	47.02	47.06	46.03	42.63	44.73	44.44
Total in suspension	32.93	24.13	38.02	37.38	66.76	60.92	28.63	37.70	58.43	43.16
Total solid matter	80.01	70.27	81.44	79.19	114.08	107.98	75.14	80.33	103.20	87.60
Ammonia	4.99	4.98	4.46	4.51	5.83	5.69	4.98	4.46	5.23	4.89
	1.65	1.18	1.45	1.43	2.08	1.96	1.41	1.47	1.86	1.00
	6.64	6.16	5.90	5.99	7.91	7.67	6.39	5.93	7.08	6.49

(1) 13 Samples exclusively from the 5-acre Field, and 10 of mixtures from the 5-acre and the 10-acre Fields.
(2) 12 Samples exclusively from the 10-acre Field, and 10 of mixtures from the 5-acre and the 10-acre Fields.
(3) 13 Samples exclusively from the 5-acre Field, 13 from the 10-acre Field, and 10 of mixtures from the two fields.

TABLE IX.—Average Composition of the Rugby Sewage, supplied in each Field, in 1861, 1862, and 1863.

LBS. PER 1,000 TONS.

CONSTITUENTS.	Season 1861: April to October.		Season 1862: November 1861 to October 1862.		Season 1863: November 1861 to October 1863.		General Means of			
	10-acre Field; 13 Samples.		5-acre Field; 17 Samples.		5-acre Field; 23 (1) Samples.		24 Samples; April to October 1861.		34 Samples; November 1862 to Oc- tober 1863.	
	10-acre Field; 23 (2) Samples.		10-acre Field; 17 Samples.		10-acre Field; 23 (3) Samples.		34 Samples; November 1862 to Oc- tober 1863.		34 Samples; November 1862 to Oc- tober 1863.	
Organic matter	In solution	328.3	290.5	265.4	297.3	265.3	339.0	263.4	263.4	276.3
	In suspension	556.0	554.7	542.7	875.2	831.7	453.1	538.9	538.9	603.2
	Total	884.3	845.2	798.1	1142.4	1097.0	792.1	801.3	1022.4	879.4
Inorganic matter	In solution	1178.2	1130.0	1062.6	1366.3	1340.6	1163.9	1101.4	1177.6	1148.9
	In suspension	517.8	631.9	653.4	1261.1	1117.8	459.5	667.5	1102.4	777.6
	Total	1696.0	1861.9	1716.0	2627.3	2458.4	1623.4	1768.9	2280.0	1926.5
Total in solution		1506.5	1389.5	1335.0	1533.4	1505.9	1497.9	1363.8	1431.0	1422.1
Total in suspension		1053.3	1216.6	1196.1	2136.3	1949.5	913.6	1206.4	1871.4	1380.8
Total solid matter		2559.3	2606.1	2534.1	3669.7	3455.4	2404.5	2570.2	3302.4	2802.9
Ammonia	In solution	199.7	142.4	144.3	186.6	182.1	189.4	143.4	167.0	156.5
	In suspension	82.8	46.4	47.4	66.5	63.4	45.1	47.0	59.5	51.2
	Total	212.5	188.8	191.7	253.1	245.5	204.5	190.4	226.5	207.7

(1) 13 Samples exclusively from the 5-acre Field, and 10 of mixtures from the 5-acre and the 10-acre Fields.
 (2) 12 Samples exclusively from the 10-acre Field, and 10 of mixtures from the 5-acre and the 10-acre Fields.
 (3) 13 Samples exclusively from the 5-acre Field, 13 from the 10-acre Field, and 10 of mixtures from the two fields.

As the ammonia contributes about three fourths of the estimated money value of the constituents of town sewage, and as its amount is the best index to that of the associated constituents valuable as manure, and also to the relation of population to quantity of sewage, its proportion is of great importance to consider.

Tables VIII. and IX. show that, according to the mean of all the analyses for the entire season in each case, there was comparatively little difference in the amounts either of ammonia, or of total matters in solution, in the sewage from the two fields. The difference in the amounts of suspended matter was, however, more considerable.

Comparing the composition of the sewage of one season with that of another, as indicated by the mean result of all the analyses relating to each, it is seen that the amounts of ammonia, and total matter in solution, are less in the comparatively wet season of 1861-2, than in either of the others, whilst the ammonia is the highest in the dry season of 1862-3. As no samples were analysed during the winter of 1860-61 when the sewage was probably weaker than during the remaining portions of the season, it might be concluded that the mean of the analyses for 1861 would give too high an average composition; but, on the other hand, there were fewer analyses during the months of August, September, and October, when the sewage was comparatively strong, than during the five preceding months when it averaged much weaker, which, of course, would tend to reduce the mean. It is probable, therefore, that the mean of the analyses for 1861 gives a pretty fair indication of the composition of the sewage for that season. With regard to 1862, there were fewer analyses during the months when the sewage averaged the strongest, and hence, the mean of the results for that season probably indicate somewhat too low an average composition. Lastly, in the season of 1863, there were again fewer analyses relating to the periods when the sewage was strongest, and hence the mean of the results for that season also probably indicates somewhat too low a composition. There can be little doubt, however, that the sewage of the wet season of 1862 was the weakest, that of 1861 somewhat stronger, and that of the dry season of 1863 stronger still.

It is obvious that the variations in the composition of sewage must depend chiefly upon its state of dilution, and that the state of dilution must be very much influenced by the rainfall. An inspection of the Tables in the Appendix, above referred to, will show how extremely different was the composition at one time compared with another, notwithstanding the general agreement of the average results as between one field and the other, or between one season and another. The great variation in the composition according to circumstances is strikingly illustrated in the following Table X.; and it is the more remarkable when it is borne in mind that every sample analysed was a mixture of portions taken every few hours throughout the day for several days together:

TABLE X.—Showing the highest, lowest, and average Amounts of Ammonia, and total solid matter, in mixed Samples of Sewage, collected at different Periods in each of the three Seasons.

		Ammonia.		Total solid matter.	
		Grains per Gallon.	lbs. per 1,000 Tons.	Grains per Gallon.	lbs. per 1,000 Tons.
1861	Highest - -	15·64	500·5	216·5	6928
	Lowest - -	2·99	95·7	37·6	1203
	Mean of 24 analyses	6·39	204·5	75·1	2465
1861-2	Highest - -	11·38	364·2	129·3	4138
	Lowest - -	2·55	81·6	50·5	1616
	Mean of 34 analyses	5·95	190·4	80·3	2570
1862-3	Highest - -	12·81	409·9	269·9	8637
	Lowest - -	3·14	100·5	62·2	1989
	Mean of 35 analyses	7·08	226·5	103·2	3302

Thus, the amount of ammonia, which to such a great extent rules the estimated money value of the sewage, varied at different times during the 31 months to which the samples refer, from about $2\frac{1}{2}$ to about $15\frac{1}{2}$ grains per gallon, or from $81\frac{1}{2}$ to $500\frac{1}{2}$ lbs. per 1,000 tons; and the total solid matter varied from about $37\frac{1}{2}$ to about 270 grains per gallon, or from 1,203 to 8,637 lbs. per 1,000 tons. It will be obvious from these results how valueless for the purposes of determining the average composition of the sewage of any locality—indeed, how utterly misleading—must be the analyses of samples taken without due regard to the circumstances by which its composition is so materially affected.

Although the sewage of 1862-3 was considerably richer in valuable constituents than that of the wetter season of 1861-2, yet the mean of the whole 93 analyses relating to the sewage of the three seasons indicates a composition closely agreeing, in all essential points, with that adopted in the previous report according to the results then obtained relating to the season of 1861 alone. For reasons explained above, however, the real average composition of the sewage of the period was probably somewhat higher than is indicated by the direct numerical mean of the 93 analyses. The latter gives—

	Grains per gallon.	Lbs. per 1,000 tons.
Total solid matter - -	87·6	2,803
Ammonia - -	6·5	208

From a careful consideration of the circumstances alluded to it is concluded that the average sewage of the 31 months would more nearly contain as follows—

	Grains per gallon.	Lbs. per 1,000 tons.
Total solid matter - -	92·5	2,960
Ammonia - -	7·0	224

Assuming this to represent the average composition of the Rugby sewage during the period in question, 1,000 tons may be estimated to contain nitrogen reckoned as ammonia equivalent to that con-

tributed in the mixed excrements, and associated matters, of between 17 and 18 persons of a mixed population of both sexes and all ages in a year, or to that in between 11 and 12 cwts. of Peruvian guano.* In other words, about 1,700 tons of such sewage would contain nitrogen reckoned as ammonia equal to that in 1 ton of Peruvian guano. Yet it has been seen that the increase of grass obtained by the use of 1,000 tons of this sewage did not, under the most favourable circumstances, exceed that which would correspond to about 26 cwts. of hay, and was on the average much less. To this question of the amount of produce obtained for a given amount of manurial constituents applied further reference will be made.

Before leaving the subject of the composition of the Rugby sewage, attention should be directed to one or two other points.

In Table XI. are given the results of the analyses of samples of sewage collected in each of the two fields, every two hours from 7 a.m. to 5 p.m., on April 16, 1861. It is seen that the amounts of both total solid matter and ammonia were the least in the samples collected early in the morning, the greatest during the middle of the day, and diminished towards the evening. As, however, the samples were not taken at the outfall as the sewage came from the town at the respective hours, but after it had been pumped from the main tank in which it was being collected and mixed throughout the day as it was produced, the variations in composition at the different periods are not so striking as have been observed when samples have been collected directly from the outfall at different periods of the day.

TABLE XI.

SHOWING the COMPOSITION of SAMPLES of the RUGBY SEWAGE collected at different hours of the day, on April 16, 1861.

				Per Gallon.			
				Organic matter.	Inorganic matter.	Total solid matter.	Ammonia.
				Grains.	Grains.	Grains.	Grains.
7 A.M.	{ Five-acre Field	-	-	14·30	38·50	52·80	2·74
	{ Ten-acre Field	-	-	14·40	38·90	53·30	2·89
9 A.M.	{ Five-acre Field	-	-	20·50	39·80	60·30	4·71
	{ Ten-acre Field	-	-	20·20	47·20	67·40	3·51
11 A.M.	{ Five-acre Field	-	-	21·90	38·00	59·90	5·14
	{ Ten-acre Field	-	-	22·70	39·60	62·30	5·39
1 P.M.	{ Five-acre Field	-	-	20·10	36·40	56·50	4·82
	{ Ten-acre Field	-	-	25·20	41·60	66·80	5·74
3 P.M.	{ Five-acre Field	-	-	21·10	34·10	55·20	4·55
	{ Ten-acre Field	-	-	21·50	35·30	56·80	4·69
5 P.M.	{ Five-acre Field	-	-	22·00	35·80	57·80	4·32
	{ Ten-acre Field	-	-	20·30	37·50	57·80	4·37

* The rainfall of the period of the experiments, and, therefore, the dilution of the sewage, was, however, less than the average, according to which it is estimated that, with the present arrangements, 1,000 tons would represent the excretal matters of scarcely 17 average individuals, and the ammonia of scarcely 11 cwts. of Peruvian guano. (See pp. 44-45, and 76.)

Another point to which reference should be made is as to the amounts of phosphoric acid and potass in the sewage, and the relation of these to the ammonia (or nitrogen) associated with them; for it is obviously important to consider whether or not the mineral or incombustible constituents of sewage exist in it in sufficient proportion to the ammonia or nitrogen for the requirements of the crops to be grown; and as the phosphoric acid and potass (the one or the other or both according to circumstances) are, perhaps, the mineral constituents the most likely to be deficient relatively to the nitrogen, the proportion of them to the latter in the sewage, and in various crops, may appropriately be referred to in illustration of the point in question.

Table XII. shows the number of grains per gallon, of ammonia, phosphoric acid, and potass, and the relation of the phosphoric acid and the potass to one of nitrogen, in the few cases only in which the phosphoric acid and potass were determined in the Rugby sewage.

TABLE XII.

Particulars of the Samples.		Per Gallon.			Proportion to 1 Nitrogen.	
When collected.	Where collected.	Ammonia.	Phosphoric Acid.	Potass.	Phosphoric Acid.	Potass.
		Grains.	Grains.	Grains.		
April 1-6, 1861	- - Five-acre field	2.09	0.64	0.86	0.26	0.35
" 1-6, "	- - Ten-acre field	4.69	0.70	0.61	0.18	0.16
" 20, - May 4, 1861	- - Five-acre field	5.08	1.09	1.37	0.26	0.33
" 20, - " 4, "	- - Ten-acre field	5.64	1.23	1.20	0.23	0.23
Nov. 4-6, 1861	- - Five-acre field	10.91	2.14	4.95	0.24	0.55
" 5, "	- - Ten-acre field	11.38	2.39	4.63	0.21	0.49
Dec. 17-18, "	- - Five-acre field	9.30	1.53	3.23	0.20	0.43
" 17, "	- - Ten-acre field	7.07	1.15	3.40	0.18	0.54
July 6-11, 1861	- - Another field	8.66	3.12	3.84	0.44	0.54
" 13-18 " -	- - Another field	8.78	2.80	3.90	0.39	0.54
	Average	7.51	1.63	2.81	0.27	0.42

It is seen that when the sewage was poor in ammonia it was also poor in phosphoric acid and potass, and when rich in ammonia rich in phosphoric acid and potass. In the case of neither constituent, however, is the relation to one of nitrogen the same in all the samples; though the correspondence is perhaps quite as close as could be expected when the circumstances which rule the amount of each are taken into consideration.

By far the larger proportion of the ammonia (or nitrogen) exists in solution in the sewage, so that if the sample taken did not contain its fair proportion of suspended or sedimentary matter, the amount of ammonia would not be very materially affected thereby. Of the phosphoric acid, on the other hand, a much larger proportion exists in the suspended matter, so that if the agitation in the main tank were incomplete, the pumps worked sluggishly, or the mixing in the gauge-tank in the field before taking the sample were insufficient, the proportion of suspended matter in the sample analysed might be too small, or under other conditions it might be too large. Hence, although the phosphoric acid must bear a pretty constant proportion to the nitrogen in sewage, it

great care be not taken in the sampling, analysis may show considerable variation in the amount, and in the proportion to the nitrogen, in samples taken at different times. It was, in fact, the case, that in the instances among those to which Table XII. refers in which the amount of phosphoric acid, and its proportion to the nitrogen, were the highest, there also was the largest amount of sedimentary matter in the sewage.

Again, the potass exists exclusively in the solution, so that if its proportion to the ammonia were constant in sewage, any deviation from the exactly due proportion of sedimentary matter in the sample taken would comparatively little affect the indications of analysis in regard to it. But whilst the phosphoric acid of sewage may be said to be derived exclusively from food-refuse and excretal matters, and therefore necessarily to bear within comparatively narrow limits a pretty uniform relation to the nitrogen, the amount of potass will vary very much according to locality, and be considerably greater where the streets or roads are constructed of potass minerals (as granite) than elsewhere.

According to the average result obtained upon the analysis of ten different samples of the Rugby sewage there were, as is shown in the bottom line of Table XII., 0·27 parts of phosphoric acid, and 0·42 parts of potass, to 1 of nitrogen in the sewage; or, in other words, for 100 of nitrogen in the sewage there were 27 parts of phosphoric acid, and 42 parts of potass. The question arises—what are the proportions of the phosphoric acid and potass to the nitrogen in different crops? These vary considerably for the same description of crop according to circumstances, but the figures given in the following Table (XIII.) may be taken to represent approximately the average proportions.

TABLE XIII.

	Proportion to 1 Nitrogen.					
	Phosphoric Acid.			Potass.		
	In Corn, Roots, &c.	In Straw, Leaves, &c.	In Total Produce.	In Corn, Roots, &c.	In Straw, Leaves, &c.	In Total Produce.
Wheat	0·48	0·42	0·46	0·28	1·08	0·57
Barley	0·40	0·34	0·38	0·34	1·20	0·60
Oats	0·28	0·37	0·30	0·25	1·55	0·65
Meadow Hay	0·27	1·00
Clover Hay	0·23	0·52
Beans	0·25	0·46	0·30	0·33	1·23	0·50
Mangolds	0·17	1·00
Swedes	0·27	0·16	0·21	0·82	0·44	0·63
Common Turnips	0·28	0·18	0·23	1·60	0·71	1·17
Potatoes	0·42	1·23

Thus, according to this Table, meadow hay contains on the average 0.27 parts of phosphoric acid to 1 of nitrogen, and according to the average result obtained on the analysis of the ten samples of the Rugby sewage it contained exactly the same proportion. Of potass, on the other hand, whilst meadow hay contains 1 part, the sewage only contained 0.42 parts to 1 of nitrogen.

According to these figures, if on the application of sewage to meadow land the whole of the nitrogen supplied were recovered in the increase of produce, it is obvious that there would be associated with it in the manure almost exactly the amount of phosphoric acid, but less than half the amount of potass, required by the crop. But in practice considerably less nitrogen is recovered in the increase of crop than is supplied in the manure employed to produce it. Then, again, the dry or solid substance of sewage grass, as it is generally cut, contains a considerably higher per-centage of nitrogen than that of ordinary meadow grass as cut for hay; whilst, from the results of direct experiments made on the point, it is probable that the proportion of phosphoric acid to 1 of nitrogen is somewhat lower, and that of potass somewhat higher, in sewage grass than in ordinary meadow hay. It follows, that if the relation of the phosphoric acid and potass to the nitrogen in sewage be fairly represented by the average result of the few analyses given on the point, it would contain more phosphoric acid, though perhaps not so much potass as could be turned to the account of growth under the influence of the amount of nitrogen at the same time supplied.

Of phosphoric acid at any rate there would probably be an accumulation within the soil, rather than an exhaustion of it, by the use of sewage to grass land. Still, agricultural experience shows that an apparently excessive supply of phosphoric acid is frequently useful in giving a favourable development, or tendency of growth, to a plant; and in this way it is possible that the application of phosphatic manures in conjunction with sewage might be advantageous. As above stated, the proportion of the potass to the nitrogen in town sewage would vary considerably according to locality; and where there was no other source of it than food-refuse and the excretal matters of man and animals, it would be more likely than the phosphoric acid to be in relative defect in case of the constant application of the sewage to grass land.

In corn crops, such as wheat or barley, the proportion of phosphoric acid to nitrogen is much higher than in meadow hay, and much higher also than was found in the Rugby sewage. The average proportion in the sewage was, however, not deficient compared with that of the phosphoric acid in these crops to the amount of nitrogen which in common practice is required to be supplied in manure, to yield one of nitrogen in the form of *increased produce*. Of potass, the proportion to 1 of nitrogen in

these crops is, in the grain, which alone is generally sold off the farm, considerably less than was found on the average in the Rugby sewage. In fact, if town sewage were used on any comprehensive scale to corn crops grown in rotation, phosphoric acid would be more likely to become deficient than potass in the majority of soils; but if phosphatic manures were employed for other crops of the course, they would not need to be supplemented to the sewage for corn. It is, indeed, well known that phosphatic manures are in practice much more used, and are much more effective, for root than for corn crops, yet, as the Table shows, the proportion of phosphoric acid to 1 of nitrogen is lower in the root than in the corn crops.

Without further comments on the figures given in Tables XII. and XIII., it may be stated, in general terms, that a careful consideration of the subject leads to the conclusion that potass would be more likely than phosphoric acid to become deficient where town sewage was applied constantly to meadow land, whilst phosphoric acid would be more likely to become deficient than potass where it was applied to the ordinary crops of rotation.

VI. *Estimated Average Composition of the Metropolitan Sewage.*

It will be well to offer a few observations here on the evidence at command relating to the average composition of the Metropolitan sewage, and to the estimated money value, according to trade prices, of its manurial constituents.

In our former Report attention was called to the fact that, so far as it was then sampled and analysed, the Rugby sewage showed an average composition agreeing very closely with that calculated for the Metropolitan sewage, according to the analyses by Dr. Letheby of samples, taken at noon and midnight respectively, from 10 different sewers; and, as above stated, the further results now at command indicate pretty nearly the same average composition for the Rugby sewage as that formerly adopted. Thus, whilst calculated according to Dr. Letheby's analyses the Metropolitan sewage contained 6·66 grains of ammonia per gallon, the Rugby sewage was then estimated to contain 6·65 grains, and taking into the calculation all the results now obtained the direct mean of the 93 analyses gives 6·5 grains, and the calculated average about 7 grains.

The amount of ammonia will still, for convenience, be taken as the gauge of comparative value; not, of course, that other constituents are not equally important, but, as already referred to, as the amount of ammonia contributes a very large proportion of the money value as estimated according to composition, and is a pretty sure indication of the approximate value of the associated constituents, it comes to be a very safe index to the approximate value of the whole, and at the same time the discussion is much simplified.

Baron Liebig, adopting as his basis an analysis of Dorset Square sewage by Mr. Way, which showed nearly 18 grains of

ammonia per gallon, estimates the constituents in sewage of that composition to be worth $1\frac{1}{2}d.$ per ton, but that the value will be raised to about $4d.$ if to each ton be added $1\frac{1}{2}$ lb. of superphosphate of lime. But in his report Professor Way stated in reference to the analysis in question, together with another given at the same time, that although the results showed that there was great manurial value in sewage, yet they could not be taken as in any way affording a measure of that value; and he has since given it as his opinion that the sample of sewage, upon the analysis of which Baron Liebig relied, was undoubtedly very much stronger than the average of Metropolitan sewage. Indeed, Baron Liebig himself admitted that more certain data as to the average composition of sewage were wanting. He said, "In the calculation of the value of sewer water there is one factor doubtful, viz., the absolute amount of phosphoric acid, ammonia, and potass, which a ton of the said water contains." Moreover, his proposal to add a certain quantity of superphosphate of lime to the sewage has for its object to assimilate the proportion of phosphoric acid to the ammonia or nitrogen in the sewage to that found in Peruvian guano, which, of course, is a standard having no necessary connexion with the proportions required by soils or crops. For these reasons Baron Liebig's estimates are obviously irrelevant.*

In the Report of Messrs. Hofmann and Witt to the Main Drainage Referees, they record $8\cdot21$ grains of ammonia per gallon as the amount they found in a mixture of samples of sewage collected from the Savoy Street sewer every hour during 24 in dry weather; and the Referees had ascertained by gauging that the rate of flow from the sewers under such circumstances averaged as closely as could be expected that which would be due to water supply exclusive of rainfall. In fact, the Referees concluded that the mixed sample in question represented pretty closely the average normal sewage without rainfall of that sewer, and further, that the sewage from that sewer might be taken as fairly representing the Metropolitan sewage under the conditions stated.

With $8\cdot21$ grains of ammonia per gallon (and the associated matters), Messrs. Hofmann and Witt estimated the constituents in a ton of such sewage to be worth $2\frac{1}{10}d.$, and taking (in accordance with the information supplied to them by the Referees) the total amount of the Metropolitan sewage without rainfall at 95 million gallons a day, or about 158 million tons per annum, they estimated the total annual value of the constituents in the Metropolitan sewage to be 1,385,540*l.*

To control this estimate founded on the amount and composition of the sewage, Messrs. Hofmann and Witt calculated the value of the constituents annually voided in the forms of urine

* In a recent letter, published some time after the above was in type, Baron Liebig adopts $7\cdot2$ instead of 18 grains, as the average amount of ammonia in a gallon of the Metropolitan Sewage, inclusive of rainfall, &c. It will be seen, by the sequel, that even after this very liberal amendment of his estimate of about a year and a half ago, he is doubtless still too high; and his estimate of value of constituents is so in a corresponding degree.

and fæces by the entire Metropolitan population, assuming it to number 2,600,000 persons. In this way they arrived at a value of 1,444,177*l.*, which they considered satisfactory confirmation of the estimate deduced from the quantity and composition of the sewage. But in their confirmatory estimate they adopt, for the composition and value of the liquid and solid voidings of each individual of a mixed population of both sexes and all ages, the amounts fixed for adult males,* assuming that other matters reaching the sewers would probably make up the difference. There can be little doubt that this was making far too liberal an allowance for the other matters than human voidings, which contribute to the value of the Metropolitan sewage.

The object of the Referees appears to have been to obtain an estimate of the total annual value of the sewage, and with such very great variations as occur in both the amount and composition of the sewage when mixed with different amounts of rainfall, it is obvious that the method they adopted of collecting for analysis samples of sewage as far as possible free from rainfall, and then calculating the composition and value of the estimated total amount of dry weather sewage according to the results of the analysis of such samples, was the only admissible one unless they had taken samples almost the year round. Their estimate of the value per ton of the dry weather sewage, or sewage without rainfall, was, in fact, only a step in the process of calculating the total annual value of the Metropolitan sewage, and they did not give any estimate of the value per ton in the average condition of dilution with rainfall in which it would have to be utilized.

It is variously estimated that the normal sewage, or sewage without rainfall, is on the average the year round mixed with from two thirds to an equal bulk of subsoil water and rain. Assuming it to be so diluted by an addition of two thirds to its own bulk, this, adopting Messrs. Hofmann and Witt's analysis and valuation for the sewage without rainfall, would reduce the average amount of ammonia in the total sewage with rainfall to 4.93 grains per gallon, and it would in like manner reduce the value from $2\frac{1}{10}$ *d.* for a ton of sewage without rainfall to $1\frac{1}{4}$ *d.* for a ton inclusive of rainfall.

It is obviously important to consider whether the sample of the Savoy Street sewage analysed by Messrs. Hofmann and Witt was more probably of above or below the average composition and value of the Metropolitan sewage without rainfall? Some judgment may be formed on this point by a careful consideration of the quantity and estimated value of the constituents contributed to the sewage by each individual of a mixed population of both sexes and all ages, taken in connexion with the amount of water through which the constituents are supposed to be on the average distributed.

According to the data of Messrs. Hofmann and Witt the amounts of total solid matter, and ammonia, and the value of

* Messrs. Hofmann and Witt even add something to the actual experimental results obtained with Hessian soldiers to adapt them, as they say, to "John Bull"!

the constituents, in the annual total voidings of an adult male are as follows:—

Adult males, per head per annum.				
		Total solid matter.	Ammonia.	Value.
		lbs.	lbs.	s. d.
Urine	-	61	15·8	10 0½
Fæces	-	34	2·3	1 8½
Total	-	95	18·1	11 9½

Dr. Thudichum, whose estimate is more recent, and whose experimental data on the point are much more comprehensive than those relied upon by Messrs. Hofmann and Witt, gives for the urine alone of adult males the following:—

Adult males, per head per annum.				
		Total solid matter.	Ammonia.	Value.
		lbs.	lbs.	s. d.
Urine alone	-	47	15·9	10 3½

It will be observed that so far as the important items of ammonia and value are concerned, the estimates of Messrs. Hofmann and Witt for the urine of an adult male agree very closely with those of Dr. Thudichum, founded on much more comprehensive analytical data. Dr. Thudichum does not, however, give any estimate of the amount, composition, and value of the fæces. With regard to the urine, he considers that the voidings of 2,800,000 persons of a mixed population of both sexes and all ages, may be taken as equivalent to those of 2,000,000 adult males. If, therefore, we take the mean of the estimates of Dr. Thudichum and Messrs. Hofmann and Witt with regard to the urine, and those of Messrs. Hofmann and Witt with regard to the fæces, of an adult male, and reduce both in the proportion of from 2·8 to 2 according to Dr. Thudichum's basis of calculation, we shall, provided the estimates of the two authorities be correct, arrive at amounts approximately applicable to an average individual of a mixed population of both sexes and all ages. The following results are so obtained:—

Average of both sexes and all ages, per head per annum.				
		Ammonia.	Value.	
		lbs.	s. d.	
Urine	-	11·32	7 3	
Fæces	-	1·64	1 2½	
Total	-	12·96	8 5½	

Here then, founded upon the estimates of these authorities, we have nearly 13 lbs. of ammonia and nearly 8s. 6d. of value, to represent the annual mixed voidings of an average individual of a mixed population of both sexes and all ages. In our last Report 10 lbs. of ammonia only (and the value would

be less in a corresponding degree) were taken to represent the mixed excrements of an average individual. This was the estimate of Messrs. Lawes and Gilbert, founded on very comprehensive data, relating both to the amounts of constituents consumed in the food, and voided in the urine and fæces, of persons of different ages and both sexes. But as this estimate was made nearly 10 years ago, since which time much more evidence has been published relating to the amount and composition of human excremental matters, it has been thought desirable to collate, for the purposes of this Report, as far as possible the whole of the information at command up to the present time bearing upon these points.* The result of very much labour expended in this way indicates 12·6 lbs. of ammonia as the amount annually yielded by each average individual, when the calculation is based upon determinations or estimates of the amounts of nitrogen or ammonia-yielding matters voided by persons of different sexes and ages. But when the estimate is founded upon the recorded amounts of fresh urine and fæces voided by individuals of the different classes, and upon the average composition of urine and fæces respectively, the amount of ammonia indicated as the average per head, per annum, is 12·7.

It is admitted, however, by authorities on the subject, that the experimental data relating to males other than in the prime of life, and especially to females of all ages, are as yet inadequate for the basis of really trustworthy average estimates. Indeed, a careful consideration of the circumstances of the majority of the cases contributing to the averages among those divisions of the population in relation to which the evidence is the most plentiful, and of the relative character of the results where it is the most deficient, leads to the conclusion that the estimate of 12·6 lbs. of ammonia per head per annum, arrived at as above described, is in all probability too high.

Then, again, even assuming the approximate correctness of the estimates of Messrs. Hofmann and Witt and Dr. Thudichum of the amount of ammonia yielded by the urine of an average adult male (and those of Dr. Thudichum are based upon a considerable portion of the same data as the estimate of 12·6 given above), it is difficult to suppose that the urine of two such adults should be equalled by so small a proportion as 2·8 persons of a mixed population of both sexes and all ages; and hence we think that the estimate of nearly 13 lbs. of ammonia as deduced from their joint data regarding adult males, and Dr. Thudichum's measure of reduction, also involves elements of error on the side of excess.

* For nearly the whole, if not the whole, of the data upon which the new estimates are based, see "On the Sewage of London," by J. B. Lawes, F.R.S., *Journal of the Society of Arts*, March 9, 1855; "The Composition of the Urine in Health and Disease," by E. A. Parkes, M.D., 1860; "On an improved Mode of collecting Excrementitious Matter, with a view to its Application to the Benefit of Agriculture, &c.," by J. L. W. Thudichum, M.D., F.C.S., *Journal of the Society of Arts*, May 15, 1863; and "On the Elimination of Urea and Urinary Water, in relation to the Period of the Day, Season, Exertion, Food, &c. &c.," by Edward Smith, M.D., J.L.B., F.R.S., *Philosophical Transactions*, Vol. CLL, p. 747.

Lastly, the estimate of Messrs. Lawes and Gilbert of the amount of nitrogen in the food of an average individual, which was founded upon the calculation of 86 different dietaries arranged in 15 classes, according to sex, age, activity of mode of life, and other circumstances, showed an amount equivalent to rather less than 12·2 lbs. of ammonia, from which, of course, a deduction has to be made for the nitrogen retained in the body, and for loss in various ways.

Upon the whole, therefore, it is concluded that the amount of ammonia contributed to the sewer-water by each individual of a mixed population of both sexes and all ages, is pretty certainly more than 10 lbs. per annum as formerly assumed, but probably less than 12 lbs.

But it is not the human excretal matters of the resident population that alone contribute to the value of the sewage of the Metropolitan area. To these must be added those contributed by the population daily visiting the Metropolis from beyond its own limits (less the amount from those daily leaving it), the fractional part of the excretal matters of horses, cows, dogs, and other animals, of the refuse from slaughter-houses, of soot, and of the matters derived from the abrasion of the streets, which does not reach the land as stable dung, street sweepings, or in some other form, and also the refuse matters from certain manufacturing processes. It is very difficult to estimate the amount and value of the constituents reaching the sewers from these sources. They are doubtless large in the aggregate; but a little reflection leads to the conclusion that they are very small per head of the population, and that they must bear a very small proportion to those of the human excretal matters of the resident population. Indeed, so far as existing information bears upon the point, it would appear probable that not more than 12½ lbs. of ammonia are contributed annually to the sewers from all sources, per head, of the population. This, including the associated matters, would, according to trade prices, represent a value of 8s. 4d. per annum for the manurial constituents of the sewage for each average individual of the population.

The question arises—through how much water are these 12½ lbs. of ammonia (with the associated constituents), or, reckoned in value, through how much is this 8s. 4d. or 100 pence worth of manurial matter on the average distributed?

The dry weather sewage, or sewage without rainfall, of the Metropolis is variously estimated at from 5 to 7 cubic feet per head per day, equal respectively 31½ and 43½ gallons per head per day, or 50½ and 71 tons per head per annum. According to the information furnished by the Referees to Messrs. Hofmann and Witt, it averaged about half way between these two extremes, namely, about 36½ gallons per head per day, equal about 59 tons per head per annum. We shall therefore probably be not far *wrong* if we take 60 tons per head per annum as the average *amount of the normal or dry weather sewage*. It is further *variously estimated* that by subsoil water and rainfall the bulk

Of the fluid will be increased by from two thirds to an equal volume.* Adopting the lower of these suppositions, which if too low will allow something for the occasional escape of storm-water, we have the 100 pence worth of constituents distributed through 100 tons of fluid, giving to it a value of one penny per ton according to the estimated market value of its manurial constituents.

What the real average dilution of the sewage the year round is, or will be, will doubtless be more certainly ascertained before long, when the main drainage system is completed, and the works have been a sufficient time in operation to render the gaugings safely available for the purposes of average estimates. In the meantime the following Table will be useful, showing the grains of ammonia per gallon, and the estimated value of the constituents per ton of sewage, on the assumption that the amount of ammonia contributed to the sewage from all sources will be at the rate of $12\frac{1}{2}$ lbs. per head per annum of the population, and on the alternative assumptions, that the amount of fluid will be 60, 70, 80, 90, and 100 tons per head per annum. For comparison with these estimates there are given those of Messrs. Hofmann and Witt, according to their direct analysis of the dry weather sewage, and also the calculated amounts according to their analysis, supposing a dilution of 100 tons per head per annum with subsoil water and rainfall, instead of 60 tons without them, which their mixed sample of the Savoy Street sewer was supposed to represent.

TABLE XI.—Grains of Ammonia per Gallon, and estimated Value of the constituents in one ton, of Sewage at different dilutions, supposing $12\frac{1}{2}$ lbs. of ammonia per head per annum, from all sources.

	Ammonia per Gallon.	Estimated Value per Ton.
	Grains.	d.
If 60 tons of fluid per head per annum - - -	6.51	1.67
70 " " " - - -	5.58	1.43
80 " " " - - -	4.88	1.25
90 " " " - - -	4.34	1.11
100 " " " - - -	3.91	1.00
Hofmann and Witt—mixed sample of dry weather sewage from Savoy Street sewer - - -	8.21	2.11
The same if diluted with two-thirds its volume of sub- soil water and rain-fall - - - -	4.93	1.27

Thus, assuming the amount of ammonia contributed to the sewage from all sources to be $12\frac{1}{2}$ lbs. per head of the population, per annum, the dry weather sewage reckoned at 60 tons per head per annum would contain $6\frac{1}{2}$ grains of ammonia per gallon; and, according to the currently adopted modes of computation, taking dry and portable manures as the standard, the total manurial

* Mr. Bazalgette informs us that the quantities of subsoil water and rainfall are too variable and irregular to enable him to give us any average.

constituents in one ton of such sewage will be worth $1\frac{1}{2}$ penny. Or, reckoning the average sewage, including subsoil water and rainfall, at 100 tons per head per annum, it will contain scarcely 4 grains of ammonia per gallon, and the manurial constituents in one ton of it will be worth just one penny.

If the above estimates for the dry weather sewage be correct, it is obvious that the sample analysed by Messrs. Hofmann and Witt was about one fourth stronger than the average Metropolitan sewage without rainfall. Their sample was, indeed, collected in a manner which gives it a better claim to the character of an average sample than any other of which the circumstances of collection, and the composition as determined by analysis, have hitherto been recorded. Still, it cannot be surprising to find that their results are probably in error to the extent above stated, when it is borne in mind that their sample was taken during 24 hours only, from one sewer only, and from one contributed to by a much more dense population than the average of equal areas in the Metropolitan district.

The approximate correctness of the above estimates of the average composition of the Metropolitan sewage, founded upon a consideration of the amount of manurial matters contributing to it and the amount of water through which they are distributed, is confirmed in a very striking and satisfactory manner by the fact, that the average amount of ammonia in the Rugby sewage, as deduced from the results of the direct analysis of 93 samples, representing the flow of 31 months, is almost exactly that arrived at by calculation, based upon a knowledge of the amount of population, and the average amount of water contributing to the sewage, and upon the assumption that $12\frac{1}{2}$ lbs. of ammonia will be contributed to it per head per annum. Thus, based upon the actual water supply over the three years 1859, 1860 and 1861, and upon the average rainfall over the seven years from 1855–1861 inclusive, it is estimated that there is at Rugby an average of about 60 tons of sewage per head per annum; and as Table XI. shows, if $12\frac{1}{2}$ lbs. of ammonia be distributed through that amount of fluid it will contain $6\frac{1}{2}$ grains of ammonia per gallon, and will have a value, calculated according to the trade prices of its manurial constituents, of $1\frac{1}{2}$ penny per ton. But the 31 months, to the sewage of which the analyses refer, were drier than usual, and hence the average amount of ammonia per gallon amounted to 7 grains, which, still reckoning $12\frac{1}{2}$ lbs. of ammonia per head per annum, is equivalent to only $55\frac{1}{2}$ tons of fluid per head per annum, which is probably more nearly the amount than 60 tons for the comparatively dry period.

Assuming the sewage of Rugby to amount to 60 tons, and that of the Metropolis to 100 tons per head per annum, including subsoil water and rainfall, it results that the latter will have less than two thirds the strength and value of the former. In fact, whilst 1,000 tons of the average sewage of Rugby *would represent between 16 and 17 individuals of a mixed population of both sexes and all ages, the same quantity of that of*

the Metropolis would represent only about 10 such individuals; and whilst, reckoned according to the amount of constituents they respectively contain, a ton of the average Rugby sewage would be worth 1½ penny, a ton of the Metropolitan sewage would be worth only 1 penny.

It may safely be affirmed that there is, as yet, no record of the analysis of any sample of sewage the circumstances of the collection of which justify the assumption that it fairly represents the average Metropolitan sewage, either with or without subsoil water and rainfall. Compared with any analytical results hitherto published, there can be little doubt that those arrived at by the synthetic method above described are much more trustworthy. It is to be hoped, however, that when the main drainage system comes to be in full operation, competent persons will be appointed to superintend the gauging, sampling, and analysis, with a view to providing data which may serve to determine satisfactorily the approximate average composition of the Metropolitan sewage as it will have to be dealt with in any plan of utilization.

For the convenience of those interested in estimates of the composition and value of town sewage, it may be mentioned that if a value of 8d. be put upon every lb. of ammonia, or if for each grain of ammonia per gallon a value of one farthing per ton be given to the sewage, the result will in either case agree almost exactly with that obtained by the elaborate method of giving the currently adopted market values to the several constituents, taking dry and portable manures as the standard.

The subject of the value realized, or realizable, by the agricultural utilisation of sewage in various ways is, of course, quite distinct from that of the estimated money value, according to the trade prices of the constituents, and is considered in other Sections of the Report.

VII. *Composition of the Drainage Water (Rugby).*

It is obvious that in attempting to determine experimentally the effects of applying to the land given amounts of sewage of known composition, there are other data essential to a right judgment of the results than those provided in the records of the increased amounts of produce yielded. It is necessary to consider,—

1. To what extent the sewage is deprived of its manurial or putrescible constituents in its passage over and through the land?
2. Whether the sewaged land is left in a higher or a lower condition after the removal of the crop?

To gain some information in reference to the second of these points it was decided that during the season of 1864 the produce of each plot should be carefully weighed, sampled, and analysed, without any further application of sewage, and that the soil of each plot should be submitted to such chemical examination as time and other circumstances would permit. The results of this part

of the inquiry are recorded in Section XII., p. 62, et seq. It may, however, be here observed, in passing, that barley (unmanured) was grown upon the three plots devoted to the experiments on the application of sewage to rye-grass in 1863, and that the crop, during growth, was obviously heavier where sewage had been applied than where it had not, and heavier also where the larger than where the smaller quantity was employed.

In order to determine how far the sewage-water was deprived of its manurial, or putrescible, constituents in its passage over and through the land, samples of the drainage water were collected simultaneously with those of the sewage in each field (commencing May 1862 and ending October 1863), and sent to Professor Way for analysis. In all 62 analyses of drainage water (or rather partial analyses corresponding in detail with those of the sewage) have thus been made. A few other special analyses, in much more detail, have been made of the sewage and drainage of the season of 1864.

The detailed results of the 62 analyses above referred to will be found in Tables XXII.-XXVII., pp. 170-175, in the Appendix. The following Table XV. gives a summary of the results, showing, in parallel columns, the average composition of the sewage and the drainage water, the first division referring to the samples collected during the months of May to October inclusive, 1862, and the second to those obtained from November 1862 to October 1863, both inclusive.

TABLE XV.—Average Composition of the Sewage and Drainage Water collected at Rugby in the Seasons of 1862 and 1863.

Grains per Gallon.

		Five-acre Field.		Ten-acre Field.		The Two Fields.	
		Sewage water.	Drainage water.	Sewage water.	Drainage water.	Sewage water.	Drainage water.
Season 1862.—May to October, both inclusive.							
Substances in Solution	Organic -	7.83	7.18	7.60	7.63	7.71	7.56
	Inorganic -	34.49	34.50	32.38	37.10	33.44	36.01
	Total -	42.32	41.68	39.98	44.73	41.15	43.57
Substances in Suspension	Organic -	14.69	1.40	17.14	1.39	15.92	1.39
	Inorganic -	25.67	1.81	24.80	3.74	25.28	2.92
	Total -	40.36	3.21	42.03	5.13	41.20	4.31
Total organic matter -		22.52	8.58	24.74	9.23	23.63	8.95
Total inorganic matter		60.16	36.31	57.27	40.84	58.72	38.93
Total solid matter		82.68	44.89	82.01	50.06	82.35	47.88
Ammonia	In solution -	4.13	0.80	4.26	1.85	4.20	1.41
	In suspension -	1.87	0.24	1.52	0.33	1.41	0.29
	Total -	6.00	1.04	5.78	2.18	5.64	1.70

	Five-acre Field.		Ten-acre Field.		The Two Fields.	
	Sewage water.	Drainage water.	Sewage water.	Drainage water.	Sewage water.	Drainage water.

Season 1863.—November 1862 to October 1863, both inclusive.

Substances in solution	Organic -	8.35	7.46	8.30	7.98	8.32	7.73
	Inorganic -	39.67	38.66	38.77	41.35	39.18	39.98
	Total -	47.92	46.01	47.07	49.33	47.50	47.71
Substances in suspension	Organic -	27.35	1.41	25.99	3.29	26.69	2.37
	Inorganic -	39.41	2.14	34.93	3.98	37.22	3.06
	Total -	66.76	3.55	60.92	7.22	63.91	5.43
Total organic matter -		35.70	8.87	34.29	11.27	35.01	10.10
Total inorganic matter		78.98	40.69	73.70	45.28	76.40	43.04
Total solid matter		114.68	49.56	107.99	56.55	111.41	53.14
Ammonia	In solution -	5.83	0.69	5.69	1.85	5.76	1.23
	In suspension -	2.03	0.15	1.98	0.31	2.03	0.23
	Total -	7.91	0.84	7.67	2.16	7.79	1.51

It is seen that of the matter in suspension nearly the whole, both organic and inorganic, has been separated from the sewage in its passage over and through the land; the drainage water containing but little of either, and probably a considerable part of that which it did contain was not contributed by the sewage, but was derived from the soil itself.

Of matter in solution, on the other hand, a gallon of drainage water contained sometimes more and sometimes less, but on the average much about the same amount both organic and inorganic as a gallon of the sewage. Here, again, there can be little doubt that a considerable portion of the matters found in the drainage water had their source in the soil itself—that there had, in fact, been an interchange; the sewage giving up to the soil valuable manurial constituents, and the fluid in its turn taking up substances from the soil for which the latter had less power of retention.

The character of the interchange of matters in solution as the sewage passed through the soil will be better understood on an inspection of Table XVI. (overleaf); but before referring to its details attention should be directed to one or two other points brought to light in Table XV.

As might be expected there was a larger portion of almost every constituent, or class of the constituents, enumerated, in the drainage from the high ridged and steeply sloping 10-acre field, over and through which the fluid passed the more rapidly, than in that from the almost level 5-acre field. This result is not only the most remarkable in degree, but the most important to remark, in the case of the ammonia. Thus, whilst a gallon of the drainage water from the 5-acre field contained, in 1862 only from one fifth to one sixth, and in 1863 little more than one tenth as much ammonia

as a gallon of the sewage, a gallon of the drainage from the 10-acre field contained, in 1862 more than one third, and in 1863 more than one fourth as much as an equal volume of the sewage. It is obvious, therefore, that the retention by the soil of the valuable manurial matters of the sewage was much less perfect in the case of the high ridged and steeply sloping 10-acre than in that of the flatter 5-acre field.

As above referred to, Table XVI., which now follows, shows more in detail the changes in the composition of the fluid in its passage through the soil.

TABLE XVI.—Detailed Composition of Samples of Sewage and Drainage Water collected at Rugby in the Summer of 1864.

Constituents.		Grains per Gallon.			
		Collected July 6-11.		Collected July 13-18.	
		Sewage.	Drainage.	Sewage.	Drainage.
In solution	Inorganic matter:—				
	Oxide of Iron, &c.	Traces.	—	1·25	0·25
	Lime	8·45	10·25	8·23	10·08
	Magnesia	1·76	1·69	1·80	1·69
	Soda (1)	5·46	0·33	5·24	2·30
	Chloride of Sodium (1)	6·82	9·73	8·53	9·21
	Chloride of Potassium (1)	6·08	1·50	6·17	2·84
	Sulphuric acid	4·39	6·55	4·01	6·75
	Phosphoric acid	1·23	0·44	1·66	0·32
	Carbonic acid	8·83	6·13	7·42	7·01
	Silica	1·80	0·80	1·00	0·80
	Total	44·87	37·52	45·31	40·75
In suspension	Organic matter	11·20	7·80	10·00	7·05
	Total matter in solution	56·07	45·32	55·31	47·80
	Inorganic matter:—				
	Oxide of Iron and Alumina	4·87	..	6·80	..
	Lime	4·43	..	3·75	..
	Magnesia	0·66	..	0·25	..
	Carbonic acid	3·25	..	2·17	..
	Phosphoric acid	1·84	..	1·14	..
	Silica, sand, &c.	31·00	..	39·30	..
	Total	46·39	..	52·91	..
	Organic matter	40·40	..	33·40	..
	Total matter in suspension	86·79	..	86·31	..
Total inorganic matter		91·26	37·52	96·22	40·75
Total organic matter (2)		51·60	7·80	43·40	7·05
Total solid matter		142·86	45·32	140·62	47·80
(1) Containing	Potass	3·84	0·04	3·90	1·48
	Soda	9·07	5·54	9·76	7·17
	Chlorine	7·03	6·61	8·10	6·70
(2) Containing	Ammonia	5·74	0·93	6·36	0·93
	In solution	2·92	..	2·42	..
	In suspension	2·82	..	3·94	0·93
	Total	8·66	0·93	8·78	0·93
Nitric acid in solution = Ammonia	(2) 1·33	..	(2) 1·41

(2) 4·227 Nitric acid = 1·096 Nitrogen = 1·331 Ammonia.

(1) 4·463 " " = 1·163 " " = 1·411 " "

77

The samples of sewage and drainage to which Table XVI. refers were not collected in either of the fields formerly under experiment, but in a meadow in the occupation of Mr. Campbell, where sewage had been pretty liberally applied for the last two or three seasons, but in which the application had been suspended for some weeks until within a few days of commencing to take the samples. The plan of collection, both for sewage and drainage, was, to take of the former about a gallon, and of the latter about half a gallon, 8 or 10 times during the 10 or 12 working hours of the day; at the end of the day after well shaking to take a gallon from each mixture, and to repeat this for six consecutive days until six gallons of each was obtained; when, from each, after well shaking, a two gallon sample was taken, and sent to Professor Way for analysis. The sewage flowed from the main into open runs for distribution over the land, but for the purpose of collecting the samples, the stand pipe was each time affixed, and the water allowed to flow through it for ten minutes before the sample was taken. The drainage was taken direct from the main cross pipe drain at the lower end of the field. Both sewage and drainage were allowed to run an hour or two each morning before taking the first sample.

During, and for sometime previous to, the collection of these samples, the weather was unusually dry, and the land was itself so dry, and in many places cracked, that it was feared a good deal of the sewage would find its way too directly to the drains. Judging from the results, however, which show a less amount of matter in solution (both organic and inorganic) in a gallon of drainage in proportion to that in an equal volume of sewage than in the average of the cases relating to the other fields and previous seasons, it would appear that the soil had done its work of absorption at any rate as well in the cases to which the more detailed analysis refer as in the majority of the others. It would, obviously, be desirable to have samples taken for analysis under very different conditions of the weather and of the land; and the plan was, to have such samples taken and analysed; but up to the time of writing the drought has continued, and it is doubtful whether results relating to wetter weather can be available for embodiment in the present report.

In judging of the results of either Table XV. or XVI., it will of course be borne in mind, that the quantity of any constituent in a gallon of the drainage water compared with that in a gallon of the sewage, by no means directly indicates the proportion supplied by the latter which has not been taken up by the soil. We had not the means of gauging the amount of fluid passing off as drainage water; but excepting when the land is already saturated, it must obviously be considerably less than that passed on to it as sewage. In fact, a gallon of drainage water will represent much more than a gallon of sewage, and hence the amount of any constituent of the sewage found in a gallon of the drainage must have been derived from more than a gallon of the former. The non-retention of valuable manurial matters by the soil was, therefore,

not so great as would at first sight appear from an inspection of the comparative composition of equal volumes of sewage and of drainage water.

It is satisfactory to observe that, of the inorganic matters in solution in the sewage, by far the larger proportion of those constituents which, by the removal of the produce from the land, are the most likely to become deficient relatively to others, is retained by the soil. Thus, smaller proportions of both the potass and the phosphoric acid coming on to the land in the sewage passed off in the drainage than of any other constituents. Of the bases, soda was also retained by the soil to a considerable extent, magnesia in a less degree, and lime less still. Indeed, of lime, there was more in a gallon of drainage than in a gallon of sewage. Of sulphuric acid, again, there was considerably more in the drainage than in an equal volume of the sewage. Lastly, of soluble silica a considerable portion passed off in the drainage.

Of inorganic matter in suspension, the quantity in the drainage water was so small, and it was so obviously derived from the soil, that it was considered quite unnecessary to submit it to analysis. It may be concluded, indeed, that practically the whole of the suspended matter of the sewage, both inorganic and organic, would be retained by the soil. It will be observed that a considerable proportion of the phosphoric acid of the sewage was in the suspended matter; and as there was none in that of the drainage, a much larger proportion of the total amount of that constituent of the sewage was retained by the soil than appears from the figures relating to the phosphoric acid in solution alone.

Of organic matter in solution a very considerable quantity was found in the drainage water; though, compared with the amount in the sewage, not quite so much in the two cases to which Table XVI. refers as on the average of the large number of cases to which Table XV. relates. There can be little doubt, however, that the soluble organic matter of the drainage was very different in character to that in the sewage. That set down as soluble organic matter in the sewage contained a very much larger proportion of nitrogen as ammonia, or ammonia-yielding matter, than that in the drainage. It is probable too, that, during periods of active vegetation, a notable portion of the soluble organic matter of drainage will frequently be derived from vegetable matter within the soil, and not directly from the sewage.

An important point to remark, which the more detailed analyses recorded in Table XVI. discloses, is, that whilst the sewage did not contain an appreciable amount of nitric acid, the drainage contained more nitrogen in that form than as ammonia; and adding the amount of ammonia to which the nitrogen in the nitric acid is equivalent to that determined as such, it would appear that the soil had not retained so large a proportion of that important manurial constituent of the sewage as might have been judged if only the more partial analyses, the average results of which are recorded in Table XV., had been made.

The amounts of potass, phosphoric acid, ammonia, and nitric acid, found in the drainage water, clearly show that the sewage was not perfectly deprived of its valuable manurial matters in its passage through the soil; and the amounts of total soluble matter, and especially of soluble organic matter, show that it was by no means perfectly purified. There is, indeed, a limit, depending upon the physical and chemical characters of the soil, and upon the amount and composition of the fluid passed through it, to the power which a soil possesses of removing substances from solution, or of preventing those already absorbed from being dissolved, in water passing through it; and so far as the soluble organic matters of the drainage are derived from vegetable matter within the soil, it is a question whether there will not always be a considerable amount in that passing from land covered with a luxuriant vegetation. So far, however, as the nitrogen of the drainage exists in the form of nitric acid, it is a pretty satisfactory indication that the organic matter has to a great extent already passed the stage of deleterious putrescence.

In the experiments under consideration the arrangements were not such as to allow of the water drained from one portion of the land being passed over another, otherwise it would no doubt have been more completely both utilized and purified. At Beddington, near Croydon, where a tract of about 250 acres is already laid down for sewage irrigation by gravitation and open runs (and the area is in course of enlargement), a great portion of the water does duty twice, and sometimes three times; and from the results of some analyses of the sewage and of the drainage water, which have kindly been communicated by Mr. Latham, the Engineer to the Croydon Local Board of Health, it appears (see Appendix, No. 3, p. 203.) that the water eventually passes from the land in a state of much greater purity than was the case in the Rugby experiments. In fact, before the present arrangements were in force, the Croydon Board had to meet numerous law suits on account of the pollution of the river by the sewage; but so efficiently is the sewage now purified that those having the right of fishing in the river have found it worth while to fix gratings to prevent the fish going up the main outfall from the sewage irrigated land.

It is clear that in any extensive scheme for the irrigation of grass land by sewage, the arrangements should be such as to allow of the water being passed over or through the land more than once, so that both the utilisation and the purification may be as complete as possible.

VIII. *Composition of the Unsewaged and Sewaged Grass.*

It has been seen that, reckoned in the fresh or green state, a greater weight of sewaged than unsewaged grass, was required to yield a given amount of milk or increase in live

weight; but that less of the dry or solid substance of the sewage grass was required to produce a given amount of milk or increase. It was further found that especially in the case of the sewage grass, it required less, both of green grass and of dry substance of grass, to yield a given return in milk during the earlier than the later portions of the season, and also less in one season than in another. It is obviously important, therefore, to ascertain the difference in the proportion of dry or solid substance, and the difference in the composition of the solid substance itself, of the grass grown without and with sewage, with smaller and with larger quantities of sewage, at different periods of the season, and in different seasons.

The mode of taking and preparing samples for analysis was sufficiently described in our former Report; and the general results of the analyses of the produce of the first season (1862) were also there discussed; the details being given in Tables XXXIX., XLV., in the Appendix to that Report.

The detailed results of the analysis of the produce of 1862 and 1863 are given in Tables XXVIII.-XL., pp. 176-193, Appendix No. 1. to the present Report; but the following summary Tables XVII. and XVIII. bring together, side by side, the mean results, both as to proportion of dry substance and composition of dry substance, for each of the three seasons over which the experiments have been conducted, and these will be sufficient to indicate the chief points of interest.

The general indications of the further results now adduced are strictly accordant with those formerly reported; the only new point illustrated being the difference in the composition of the grass in one season compared with another.

Comparing first the composition of the grass produced under different conditions, in one and the same season, it is seen that there is in each season a very great difference, both in the proportion of the dry substance and in the composition of that dry substance, according to the varying circumstances of growth. With scarcely an exception in either season, the proportion of dry or solid substance in the grass as cut, weighed, and given to the animals, was considerably lower in the sewage grass than in the unsewaged grass, and generally the lower the larger the quantity of sewage employed. There was also, pretty uniformly, a diminished proportion of dry substance in each successive cutting as the season advanced.

It will be readily understood that the proportion of dry or solid substance in the grass depends upon the stage of growth, the proportion of leaf and stem, and the condition of the weather at the time of cutting. The grass grown without sewage was for the most part cut at a later stage of growth, and showed more tendency to form stem and seed than that grown with it, and *the greater the quantity of sewage the greater was the product*

TABLE XVII.—Per-centages of Dry Substance in the Unsewaged and the Sewaged Grass.

SEASONS 1861, 1862, and 1863.

		Five-acre Field.					Ten-acre Field.				
		Un-sewaged, Plot 1.	Sewaged.			Mean.	Un-sewaged, Plot 1.	Sewaged.			Mean.
			Plot 2.	Plot 3.	Plot 4.			Plot 2.	Plot 3.	Plot 4.	
Meadow Grass. First Season, 1861.											
1st Crop	-	27.9	30.5	26.9	27.7	28.3	22.0	23.3	21.4	18.4	21.3
2d Crop	-	24.4	19.8	14.2	13.3	17.9	26.9	17.1	15.1	16.1	18.8
3d Crop	-	..	13.4	13.7	12.9	13.3	..	12.6	7.3	14.4	11.4
4th Crop	-	15.4	9.6	12.5	..	16.9	15.1	17.8	16.6
Mean	-	26.2	21.2	17.6	15.9	..	24.5	17.5	14.7	16.7	..
Meadow Grass. Second Season, 1862.											
1st Crop	-	26.7	22.8	14.4	15.3	19.8	26.9	19.5	13.5	13.1	18.3
2d Crop	-	22.8	14.3	16.4	19.4	18.2	17.9	16.2	19.0	16.7	17.5
3d Crop	-	..	18.2	12.9	14.2	15.1	..	14.5	14.4	15.8	14.9
4th Crop	-	*33.8	33.8
Mean	-	24.8	18.4	14.6	16.3	..	22.4	16.4	15.6	15.2	..
Meadow Grass. Third Season, 1863.											
1st Crop	-	26.1	21.5	17.6	16.3	22.9	30.8	18.6	20.0	14.6	23.3
2d Crop	-	34.4	18.5	14.9	17.8	21.4	18.2	17.7	16.3	18.3	17.8
3d Crop	-	..	17.7	10.9	17.6	15.4	..	12.4	14.6	15.2	14.1
4th Crop	-	..	15.8	13.0	12.3	13.7	13.9	13.6	13.8
5th Crop	-	15.3	15.3
Mean	-	35.3	18.4	14.1	15.9	..	29.0	16.2	16.2	15.6	..
Italian Rye Grass, 1863.											
		Un-sewaged, Plot 1.	Sewaged.		Mean.						
			Plot 2.	Plot 3.							
1st Crop	-	21.3*			21.3						
2d Crop	-	23.7	18.8	17.5	20.0						
3d Crop	-	36.4	27.5	18.3	27.4						
4th Crop	-	33.2	13.8	18.9	22.0						
5th Crop	-	19.9	16.8	17.8	18.2						
6th Crop	-	..	18.6	20.4	19.5						
		28.3	19.1	18.6	..						

* All three plots were unsewaged until after the first cutting.

TABLE XVIII.—Mean Composition (per cent.) of the Dry Substance of the Grass, without and with Sewage, and in each successive Crop.

In Seasons 1861, 1862, and 1863.

	Without and with Sewage.				Each successive Crop.					
	Un-sewaged, Plot 1.	Sewaged.			1st Crop.	2d Crop.	3d Crop.	4th Crop.	5th Crop.	6th Cr
		Plot 2.	Plot 3.	Plot 4.						

Meadow Grass—First Season, 1861.

Number of analyses giving the means	5	7	9	9	11	9	7	5		
Nitrogenous substance ($N \times 6.3$)	13.08	18.67	18.92	19.78	10.33	18.07	23.76	28.25		
Fatty matter (ether extract)	3.21	3.54	3.53	3.44	3.01	3.60	3.65	3.84		
Woody fibre	28.80	29.34	30.15	29.13	30.80	28.45	28.50	28.60		
Other non-nitrogenous substances	45.66	37.09	35.94	35.92	47.79	38.28	30.84	24.57		
Mineral matter (ash)	9.25	11.36	11.46	11.73	8.07	11.60	13.25	14.74		
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00		

Meadow Grass—Second Season, 1862.

Number of analyses giving the means	4	6	6	7	11	9	6	1		
Nitrogenous substance ($N \times 6.3$)	9.49	15.65	15.70	16.83	11.65	12.70	20.44	18.22		
Fatty matter (ether extract)	2.93	3.81	3.64	3.85	2.82	3.72	4.34	4.42		
Woody fibre	29.80	29.20	29.18	29.86	32.42	29.01	26.69	24.86		
Other non-nitrogenous substances	47.84	40.70	40.50	39.01	44.40	43.87	36.00	38.70		
Mineral matter (ash)	9.94	10.64	10.98	10.45	8.71	10.70	12.53	13.80		
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00		

Meadow Grass—Third Season, 1863.

Number of analyses giving the means	4	7	8	9	8	8	6	5	1	
Nitrogenous substance ($N \times 6.3$)	10.31	18.43	20.49	22.38	15.05	16.64	19.88	26.12	32.19	
Fatty matter (ether extract)	4.08	5.04	4.68	4.83	4.79	4.31	5.00	4.93	5.07	
Woody fibre	28.64	26.08	25.62	25.40	26.55	28.07	26.08	23.33	20.51	
Other non-nitrogenous substances	47.29	39.48	37.82	35.79	44.32	39.68	37.05	33.26	29.62	
Mineral matter (ash)	9.68	10.97	11.39	11.60	9.29	11.30	11.99	12.36	12.61	
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

Italian Rye-grass, 1863.

Number of analyses giving the means	5	5	5		1	3	3	3	3	
Nitrogenous substance ($N \times 6.3$)	12.44	18.78	18.11		12.51	12.88	11.07	16.70	20.91	24
Fatty matter (ether extract)	3.53	4.45	3.85		3.61	3.01	3.28	4.31	4.49	5
Woody fibre	24.68	25.51	25.55		17.79	26.95	28.79	27.16	22.64	22
Other non-nitrogenous substances	49.21	39.76	41.47		56.59	48.28	48.88	40.87	38.34	33
Mineral matter (ash)	10.14	11.50	11.02		9.50	8.90	7.98	10.96	13.62	14
	100.00	100.00	100.00		100.00	100.00	100.00	100.00	100.00	100

of succulent leaf; though, even with sewage, the tendency to run to seed is much greater in hot and dry than in cold and wet seasons. Then again, the earlier crops of the season are not only grown under much more favourable maturing conditions, but, from their much greater abundance, they are generally cleared more slowly, and are therefore liable to be in a more advanced stage when cut; whilst the later ones are not only produced under less favourable maturing circumstances, but are frequently much more affected in their condition by unfavourable weather at the time of cutting.

With Italian rye grass, as well as with meadow grass, the herbage in the condition in which it was cut was found to be much more succulent when grown with sewage than without it. In the case of the rye grass, however (though it is true only small quantities of sewage were applied, and the results relate to only one season), the diminution in the proportion of dry substance as the season advanced was somewhat less marked.

The general result is, that there was a less proportion of dry or solid substance in the sewaged grass, as cut, than in the unsewaged; but that a given amount of dry substance in the sewaged was more productive of milk and increase than an equal amount in the unsewaged grass.

The question arises,—was there any difference in the composition of the dry or solid substance of the unsewaged and the sewaged grass which may account for the higher food-qualities of that of the sewaged?

Table XVIII. shows that the most remarkable difference was in the proportion of the nitrogenous constituents, the per-centage of which was in each season much higher in the solid matter of the sewaged than in that of the unsewaged grass, and also the higher the greater the quantity of sewage applied. The proportion of green and impure fatty or waxy matter was also somewhat, but in a less degree than the nitrogenous substance, the greater in the sewaged grass. The comparatively indigestible woody fibre, judging from the results of 1862 and 1863, when the sewaged crops were cut in a younger and more favourable condition than in 1861, will probably average less in sewaged than in unsewaged grass. But the mineral matter, like the nitrogenous and green fatty or waxy matters, is in the larger amount in the sewaged grass, and like them also, a relatively large amount of it is generally indicative of a more unripe and succulent condition.

That the higher milk-yielding qualities of the solid matter of the sewaged grass do not depend simply on its higher per-centage of nitrogenous constituents is evident from the fact that the solid matter of the later crops of the season, which, weight for weight, had much less value as food than that of the earlier, nevertheless contained a very much higher proportion of nitrogenous substance. Indeed, there was generally more than twice as much nitrogenous

substance in a given amount of the solid matter of the last than of the first crop of the season.

It would appear that the higher qualities of the solid matter of the sewage grass, and of the grass grown in the earlier and more genial periods of the season, were due rather to a favourable condition of maturation, and, therefore, of digestibility and assimilability, of the constituents. That the condition of maturation or elaboration of the constituents had much to do with the quality of the grass is evident from the fact that the produce of the warmer seasons of 1861 and 1863 was much more productive than that of the cold and wet season of 1862. And that a comparatively high per-centage of nitrogenous substance is only advantageous when accompanied with a favourable condition of maturation, may be gathered from the fact that with the higher per-centage of nitrogen in the produce grown in the more favourable seasons of 1861 and 1863 there was higher feeding quality, whilst with the higher per-centage of nitrogen in the produce grown in the later and colder periods of the seasons, there was lower feeding quality.

Italian rye grass seems to be subject to very similar variations in composition, by the application of sewage, and at different periods of the season, as meadow grass; but as the amounts of sewage applied to it were comparatively small, and the results relate to one season only, it can scarcely be judged with certainty, whether or not the changes in composition would, under comparable circumstances, be much the same in degree as well as kind, with the two descriptions of herbage. The feeding results seem to indicate that the Italian rye grass deteriorated somewhat less than the meadow grass as the season advanced, but the difference in chemical composition offers no very obvious explanation of the fact.

IX. *Effects of Sewage on the mixed herbage of grass land in developing the more freely growing, at the expense of the less freely growing plants.*

It is well known that active manures of any kind, when applied to the mixed herbage of grass land, develop certain more freely-growing plants to the partial, or in some cases, the entire exclusion of others. Irrigation, whether by sewage or otherwise, produces very similar effects.

On careful inquiry, and by the aid of samples obtained from some of the most important sewage meadows in the neighbourhood of Edinburgh, it is found that wherever the application has been continued for a considerable number of years, the produce consists almost exclusively of rough meadow grass (*poa trivialis*), common couch grass (*triticum repens*), and in a smaller proportion of rye grass (*lolium perenne*), or rough cock's-foot (*dactylis glomerata*), or both; the chief weed being crow-foot (*ranunculus*), of various

species. In four out of five reports from as many different sewage farmers, the poa is said to stand first, and the couch grass second in degree of prominence. The poa also seems to stand first in estimation as a sewage grass; whilst the common couch is also much valued. Indeed, Mr. Thomson, of Roseburn, informs us that he has actually transplanted this weed of our corn fields from his arable land to lay down for sewage meadow, and that the result has been quite satisfactory; he also informs us that when he has sown as many as 15 or 20 different kinds, most of them have gradually died out, and after some years only a few suitable to the land and the treatment remained. (See notes on the Edinburgh Meadows, Appendix, p. 198 et seq.)

At Rugby similar effects, but at present in a less degree, have been produced. The following observations on the character of the herbage in the two fields are founded upon the records of a careful examination made in August 1862, since which time, however, further change has doubtless taken place on the sewaged plots.

The portion of land left unsewaged by the Commission in the five-acre field had received less sewage previously than that in the ten-acre field, and showed somewhat greater complexity of herbage.

In the five-acre field the most prominent grasses on the unsewaged portion were woolly soft-grass (*holcus lanatus*), common bent grass (*agrostis vulgaris*), rough meadow grass (*poa trivialis*), hard fescue (*festuca duriuscula*), rough cock's-foot (*dactylis glomerata*), and rye grass (*lolium perenne*), with a number of others in much smaller proportion. The herbage also comprised several species of the Leguminous family, besides a number of weeds, of which the most prominent were ribwort (*plantago lanceolata*), milfoil (*achillea millefolium*), sorrel dock (*rumex acetosa*), and dandelion (*taraxacum dens-leonis*). In the ten-acre, as in the five-acre field, the cock's-foot, woolly soft grass, rye grass, and hard fescue, were among the most prominent of the grasses without sewage, whilst the rough meadow grass, and others, were less prominent than in the five-acre field. The amount of Leguminous herbage was also less than in the five-acre field, whilst crow-foot was extremely abundant.

In the sewaged herbage of both fields the cock's-foot and woolly soft grass were by far the most abundant, the rye grass coming next, and perhaps the rough meadow grass or the hard fescue next, others being more reduced. In both fields the Leguminous herbage was much reduced in proportion under the influence of sewage, whilst in the five-acre field the sorrel-dock, and in the ten-acre the crow-foot, were the most prominent weeds.

In the sewage meadows near Croydon, the cock's-foot and rye grass appear to be the predominating grasses.

The general effect of sewage irrigation on the mixed herbage of meadow land may be stated to be, to develop the Gramineous

herbage chiefly, nearly to exclude the Leguminous, and to reduce the prevalence of miscellaneous or weedy plants, but much to encourage individual species. It also, at the expense of the rest, encourages a few free-growing grasses, among which, according to locality and other circumstances, the rough meadow grass, couch grass, rough cock's-foot, woolly soft grass, and rye grass, have been observed to be very prominent. The result is an almost exclusively Gramineous, and very simple herbage. But, as the produce of sewage irrigated meadows is generally either cut or grazed in a very young and succulent condition, the tendency which the great luxuriance of a few very free-growing grasses has to give a coarse and stemmy later growth is not an objection, as it is in the case of meadows left for hay. Indeed, as has been already shown, when the produce is given to animals in a green and succulent state, a given weight of the dry or solid substance of the more simple sewaged grass is more productive than an equal weight of that of the more complex unsewaged produce.

X. Composition of the milk yielded from the unsewaged and from the sewaged grass.

Once a week during the greater part of the season of 1861, the morning and evening milk of the cows fed on unsewaged grass was mixed together, and a gallon sample taken. Samples of the milk from the sewaged grass were taken in the same way. In 1862 similar samples were collected, but then only once a month. In 1863 none were taken. In all cases the samples were, as soon as taken, put into bottles filled up to the corks, sealed down, and sent off the same evening by railway to Professor Way for analysis.

In 1861 there were in all 13 samples of milk from unsewaged, and 15 from the sewaged grass, and in 1862, six from the unsewaged and six from the sewaged grass so collected and submitted to analysis. The results of each of the 28 analyses of the milk of 1861 were given in Table XLVI. in the Appendix to the previous Report, and those of the 12 made of the milk of 1862 are given in Table XLI., p. 194, in the Appendix to the present Report.

In the following summary Table, XIX., the results of the whole 40 analyses of milk are so classified as to bring to view the chief points of interest.

TABLE XIX.—Mean per-centage Composition of the Milk produced from Unsewaged and Sewaged Grass, alone or with Oilcake in addition, and in different Seasons.

SEASONS 1861 and 1862.

	SEASON 1861.				SEASON 1862.		
	Cows fed on						
	Grass alone.		Grass and Oilcake.		Sewaged Rye-grass and Clover, and Oilkake.	Grass and Oilcake.	
	Un- sewaged.	Sewaged.	Un- sewaged.	Sewaged.		Un- sewaged.	Sewaged.
	Number of Analyses giving the Means. }	9.	10.	4.	4.	1.	6.
Casein	3.246	3.241	3.353	3.423	3.125	3.513	3.467
Butter	3.604	3.430	3.657	3.707	3.473	3.834	3.559
Sugar of Milk, &c. .	4.405	4.218	4.561	4.699	4.700	4.502	4.440
Mineral matter .	0.753	0.776	0.740	0.771	0.753	0.753	0.771
Total solid matter	12.008	11.665	12.310	12.599	12.050	12.603	12.257
Water	87.992	88.335	87.690	87.410	87.950	87.398	87.743
	100.000	100.000	100.000	100.000	100.000	100.000	100.000

The average results of the numerous analyses of the milk of the season of 1861 showed a somewhat lower proportion of each of the constituents—casein, butter, sugar, &c.—and also of total solid substance, but a slightly higher proportion of mineral or saline matter in that from the sewaged than in that from the unsewaged grass, when, for a period of 16 weeks, the cows were fed on grass alone. But when, for a period of four weeks at the end of the season, oilcake was given in addition to the grass, the milk from the sewaged grass contained rather more instead of less of casein, butter, sugar, &c., and total solid matter, than from the unsewaged; and both kinds of grass, although at that end of the season, gave, with oilcake in addition, milk containing more of each of the constituents mentioned than the grass of the earlier and more genial periods of the season when it was given alone.

In 1862 the season was very cold and wet, and the yield of milk, with cake given in addition to the grass throughout the season, was little if any better than during the period of the more favourable season of 1861, when the cows had grass alone. But both from unsewaged and from sewaged grass the milk of 1862, when oilcake was given, was somewhat richer than that of 1861 without it; and as during the longer period of 1861 when grass was given alone, so now in 1862 when cake was given throughout, milk from the sewaged grass contained less solid matter, and was in fact somewhat less rich, than that from the unsewaged. This result of an entire season is, of course, more reliable than that obtained with oilcake during the concluding four weeks only of the season of 1861, which showed as above observed a rather

richer milk from oilcake and sewaged than from oilcake and unsewaged grass. Upon the whole it would appear probable that, under otherwise comparable conditions, unsewaged grass will give a slightly richer milk than sewaged, whether given alone or with other food in addition.

The general result is, that a given weight of fresh unsewaged grass, supplying as it did much more solid matter, gave more milk than an equal weight of the fresh sewaged grass; that a given amount of the dry or solid substance of the more succulent sewaged grass gave considerably more milk than an equal quantity of that of the unsewaged; that the addition of oilcake, whether to unsewaged or to sewaged grass, increased the richness of the milk; but that the milk from the sewaged grass (whether given alone or with oilcake) was somewhat less rich than that from the unsewaged.

XI. *Experiments on the application of Sewage to Oats in 1863.*

For reasons that have been already explained, the Commission did not think it desirable to undertake a systematic series of experiments with any other crops than grass. Indeed, so to have extended their inquiry, would have required much more ample funds than were at their disposal for experiments on the agricultural utilisation of sewage. By the kindness of Mr. J. A. Campbell, however, they are enabled to record the results of an experiment on the application of sewage to oats.

In the spring of 1863, in a field from which a crop of clover had been carried off in 1861, and in 1862 a crop of wheat, Mr. Campbell was about to give the then growing crop of oats a top dressing of nitrate of soda. Instead of this, four plots of about an acre each, were set apart and treated as follows:—

Plot 1. Left unmanured.

Plot 2. Sewaged at the rate of 135½ tons per acre.

Plot 3. Sewaged at the rate of 510 tons per acre.

Plot 4. Top-dressed with 1½ cwt. of nitrate of soda.

The applications of the sewage, and of the nitrate, were made much later in the season than was desirable. The sewage was applied from April 28 to May 16 inclusive; the two acres requiring, with the hindrance of gauging by means of a barrel, 16 days for the application by hose and jet of the small quantities stated. The nitrate of soda was sown broadcast, partly on April 24, and partly on May 4.

For several weeks from the time of sowing there was very little rain, so that the plant top-dressed with nitrate of soda was obviously injured by the application for some time, the foliage being much "burnt." The sewage, on the other hand, being applied during dry weather, and the application followed by a very unusually dry period, during which spring corn and even wheat crops were reputed over a considerable range of country *to be suffering for want of rain*, produced, as might be expected, *very marked effects*. Owing, too, to the small amount of rain,

the sewage was of more than the average concentration of that of Rugby, and probably about double the average strength of that of the Metropolis (including rain, &c.) The following Table (XX.) shows the results obtained.

TABLE XX.—Results of Experiments on the application of Sewage to Oats.

Rugby 1863.

Plots.	Quantities per Acre.						Particulars of Quality.		
	Manures.	Dressed Corn.	Offal Corn.	Straw.	Increase by Manure.		Weight per Bushel of dressed Corn.	Offal Corn to 100 dressed.	Total Corn to 100 Straw.
					Corn.	Straw.			
1	Unmanured - -	Bush. Pks.	Lbs.	Cwts.	Lbs.	Cwts.	Lbs.		
2	135½ tons sewage - -	53 2½	85	42½	44	3.5	53.3
3	510 tons sewage - -	69 1½	212	53	658	10½	43	7.1	53.9
4	1½ cwt. Nitrate of Soda	66 2½	302	61	565	18½	42	10.8	45.4
		54 0½	131	45½	-11	3	44	5.5	40.6

Thus, under the conditions of season described, there was with the nitrate of soda even rather less corn, and only about 3 cwts. more straw, than without manure, and the smaller quantity of sewage gave more increase of corn than the larger, though the latter gave considerably the most straw. Both the sewaged crops were, indeed, too luxuriant to bear up against the heavy rains of June, and the one with the largest amount of sewage was very much laid, and hence the deficient yield of corn in proportion to straw. The last three columns show, by the deficient weight per bushel of the dressed corn, the large proportion of offal corn, and the low proportion of corn to straw, where the largest quantity of sewage was employed, that the defective result as to corn in its case was due to over rather than to under luxuriance. In fact, the usual complaint when sewage has been applied to growing corn crops has been of over production of straw and deficient proportion of corn—that is to say, of a tendency of growth which is as unfavourable in the case of corn as it is favourable in that of grass.

There was, however, a very high gross money return per ton of sewage applied, at any rate where the smaller quantity only was employed. Thus, reckoning oats at 3s. per bushel, and oat straw at 20s. per load, the gross value of the increased produce from one ton of sewage was—

With 135½ tons of sewage per acre - 5½d. per ton.

With 510 tons of sewage per acre - 1½d. per ton.

Here, then, with a small quantity of sewage of nearly double the average strength of that of the Metropolis, applied during a period of very dry weather, which was followed by a season of very unusual productiveness—the harvest of 1863 being the best for many years past—the gross value of the increased produce amounted to more than 5d. per ton of sewage employed, or to

nearly three times the market value of the constituents of the sewage supposing them to have been extracted and dried.

The smaller amount of sewage applied was equivalent in water to something under an additional $1\frac{1}{4}$ inch of rain at the critical period of growth; and the larger amount was equal to about five inches, which would at that period have been a very great excess and of itself caused rank and over luxuriant growth on any soil in such condition as the unmanured produce showed the one in question to have been. It is indeed difficult to say how much of the actual result obtained was due to the manurial constituents, and how much to the water of the sewage. At any rate, whether considered with regard to the amount of manurial constituent supplied, or the amount of water, an average of 500 tons of sewage per acre to arable land otherwise treated in the ordinary way would most probably be found more than appropriate to the average of soils and seasons, and would most certainly be more than appropriate for heavy lands and for wet seasons; nor even in dry ones, when sewage would be worth a maximum value for some crops by virtue of its water, if applied at the proper time would more than this amount be required the year round; though it is possible that the demand might be as much beyond the supply for a short period, as the supply would undoubtedly be beyond the demand for very much the greater part of the year so far as arable land was concerned.

XII. *Miscellaneous Results obtained in 1864.*

It has been already stated (Section VII. p. 45) that in order to ascertain whether the meadow land experimented on in 1861, 1862, and 1863 were left in a higher or in a lower condition by the application of sewage, and the removal of the produce during those three years, it was decided that the produce of each plot should, in 1864, be carefully weighed, sampled, and analysed without any further application of sewage; and also that the soil of the respective plots should be so far submitted to chemical examination as time and other circumstances would allow. The results of this part of the inquiry will form the subject of the present Section.

Owing to the extraordinary drought of the season of 1864, it was, as will be readily understood, as unfavourable as it possibly could be for meadow land without either sewage or other manure. Indeed, from only one plot was any second cutting taken, and then only a few cwts. of green grass were obtained. In all other cases the after-grass came forward so late in the season that it was thought better to feed it off than to cut it.

The following Table (XXI.) shows the amounts of green grass obtained from each plot; and some particulars of the feeding of the remainder of the produce will be given further on.

TABLE XXI.—Produce of Green Grass obtained per Acre in 1864, without Sewage.

Plots.	Treatment in 1861, 1862, and 1863.	Five-acre Field.		Ten-acre Field.		Mean of the Two Fields.
		Dates of Cutting.	Quantity.	Dates of Cutting.	Quantity.	
First Crop.						
			tons. cwt. grs. lbs.		tons. cwt. grs. lbs.	tons. cwt. grs. lbs.
1	Unsewaged	June 18—20	1 14 3 26	June 16 and 17	3 1 2 0	2 8 0 27
2	3,000 tons sewage per acre, per ann.	June 4—9	2 17 2 20	June 8—13	5 12 1 15	4 5 0 33
3	6,000 tons sewage per acre, per ann.	May 24—30	5 12 0 18	June 1—7	7 4 0 9	6 8 0 133
4	9,000 tons sewage per acre, per ann.	May 19—23	5 9 0 17	May 26—June 1	6 11 0 11	6 0 0 14
Second Crop.						
4	9,000 tons sewage per acre, per ann.	Aug. 24	0 4 3 3	..	- - - -	- - - -

Small as were the amounts of produce on all the plots, it is nevertheless clear that there was much more growth where sewage had been applied in the preceding years than where it had not; and there was more where 6,000 than where 3,000 tons had been applied, even though the crop was in the former case cut some days earlier at the most active period of growth; and from the indications there would doubtless have been more still where 9,000 tons had been annually applied, but for the still earlier dates of cutting.

The evidence so far is, then, that the land was left in the higher condition of productiveness the larger the quantities of sewage applied, and of produce removed, in previous seasons; and although a second cutting was taken in only one instance, and when feeding off the after-grass the plots were not separated so as to afford exact evidence on the point, it may be stated that in both fields the amount of feed was obviously much the greater the greater the quantity of sewage previously applied. Indeed, it is concluded that in each field the plot 4 gave as much after-feed as plots 2 and 3 together.

In the ten-acre field the after-grass of the 4½ to 5 experimental acres kept 8 heifers, of about 7½ cwt. live-weight each, for 11 days, from Nov. 11 to Nov. 22; 104 sheep of about 160 lbs. average live-weight for 14 days, from Nov. 10 (morning) to Nov. 23 (evening); and 102 of the same sheep for 7 days from Dec. 7 to Dec. 14. The five-acre field, where the growth was not so good, kept 32 lambs, averaging about 90 lbs. live-weight, for 35 days from Nov. 18 to Dec. 23.

Further evidence of the effects of the unexhausted residue from the previous sewage manuring is to be found in the difference in the chemical composition of the produce from the respective plots. This point is illustrated by the results of analyses given in the following Table. For further details, see Appendix, Table XLII *p.* 196.

TABLE XXII.—Composition of the Grass obtained in 1864, without Sewage.

	First Crop.								3d Cro.
	Five-acre Field.				Ten-acre Field.				Five-acre Field.
	Un-sewaged.	Sewaged in 1861-2-3.			Un-sewaged.	Sewaged in 1861-2-3.			Plot #
	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	
Per-cent. in the Fresh Grass.									
Dry Substance - -	34.70	30.19	28.12	20.33	33.36	30.84	23.63	21.50	55. —
Per-cent. in the Dry Substance of the Grass.									
Nitrogenous substance (N x 6.3) - - -	10.09	13.73	14.91	16.07	11.00	12.96	15.63	13.47	15. —
Fatty matter (ether extract) - - -	3.87	3.66	3.75	4.43	3.88	4.14	5.03	4.85	5. —
Woody fibre - - -	26.64	26.64	27.61	28.21	27.36	27.12	23.21	26.94	29. —
Other non-nitrogenous substances - -	51.65	48.39	45.33	42.20	50.96	48.17	43.52	46.12	41. —
Mineral matter (ash) -	7.75	7.58	8.40	9.09	6.81	7.63	8.61	8.62	7. —
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100. —

The first point to remark is, that as the proportion of dry or solid substance was much lower in the earlier cut and less matured sewaged grass than in the later cut unsewaged grass, the difference in the relative amount of the produce per acre on the respective plots was in reality not so great at the actual dates of cutting as the amounts of fresh or green grass recorded in the Table would indicate. The lower per-centages of dry substance in the sewaged grass indicate indeed a less degree of maturity or ripeness at the time of cutting; and with these characters a higher per-centage of nitrogenous substance in the solid matter of the grass would be expected. But the differences in the per-centage of nitrogenous substance which the Table shows are greater than can be accounted for by the earlier or later cutting, and consequent less or greater degree of ripeness.

The solid matter of the unsewaged produce of the one field contained 10, and of the other 11 per cent. of nitrogenous substance; whilst in that of the produce from the previously sewaged plots it ranged from about 13 to about 16 per cent., and was, with one exception, the higher the greater the quantity of sewage previously applied.

It has been shown in Section VIII. that the proportion of nitrogenous substance in the solid matter of the grass was much increased under the influence of sewage, and the results here recorded, taken together with those relating to the amounts of produce per acre, clearly show a considerable effect from the unexhausted residue from the previous sewage manuring, even in so extremely unfavourable a season; and it would doubtless have

been much greater under more favourable circumstances, and will probably be manifest for some time to come.

Although the amount and the composition of the grass obtained in 1864 have clearly shown the effect of the previous sewage manuring, the only partial investigation to which the soils have been submitted does not further illustrate the point. Calculation would, indeed, seem to indicate that there had been considerable accumulation, at a greater or less depth within the soil, of some of the most important manurial constituents of sewage where the larger quantities had been applied, but it was not thought desirable to incur either the necessary delay or the increased expenditure which a sufficiently detailed investigation of the subject would involve.

The results obtained are, however, of considerable interest as showing a very great difference in the character and composition of the soils of the two fields, which to a great extent explains the marked difference in the amount of produce which they respectively yielded without sewage.

Appendix Table XLIII. p. 196, shows the great difference in the general character, and Table XLIV. p. 197, in the chemical composition in some important respects, between the two soils.

It will be recollected that the unsewaged portion of the ten-acre field each year yielded much more produce than that of the five-acre field. The former was known to have a heavier soil, and a more sunny aspect; but these differences were not recognised as fully accounting for the great difference in the natural productiveness. The results of the partial mechanical and chemical analysis show, however, that, within a layer of 9 inches, taken immediately below as thin a sod as could be first removed, the lighter, gravelly, and naturally less productive soil of the five-acre field contained an average of nearly 20 per cent. of stones, whilst similar layers from the ten-acre field did not on the average contain half as much. Again, the separated fine mould from the soil of the five-acre field was sandy, contained about 15 per cent. of moisture, between 5 and 6 per cent. of organic matter, and scarcely one fifth per cent. of nitrogen, whilst that of the more productive ten-acre field was loamy, containing a good deal of clay, about one fourth more moisture, more than one half more organic matter, and about one third more ammonia, or nitrogen in some other form. Or, comparing the composition of the total fresh soils, including stones, instead of that of the separated fine moulds, the differences are in a still greater degree in favour of the soil of the ten-acre field, so far as the indications of natural fertility are concerned.

It is, however, very satisfactory to know, that the soil of the much less naturally fertile five-acre field gave fully as much produce per acre under the influence of liberal dressings of sewage, as that of the naturally much more productive ten-acre field. This result is quite confirmatory of the general opinion, founded on the results of practical experience, that light, and even naturally poor and unproductive soils, are capable of yielding very large crops of grass under the influence of sewage, and that they are, in fact, the best suited for its application.

XIII. *Concluding Observations; general Considerations on the Agricultural Utilisation of Town Sewage.*

There cannot be a doubt that to obtain a maximum amount and gross value of produce from a given amount of sewage it should be applied in small quantities per acre, and in dry weather. But it is clear that the maximum value per ton of sewage which would be obtainable under such conditions would be available only for short periods of the year; and, it is equally certain, that the constant daily supply, the year round, would, at all other times have to be disposed of at a very different rate, thus reducing the average value very considerably. It is obvious, indeed, that even supposing sewage were distributed over a sufficiently large area to command its full value, both as manure and as water, at the best periods of the year, the much larger remainder must either be sacrificed, or at the best used for grass at periods when its value even for that crop must be very much reduced.

Even assuming that during any considerable portions of each year the Metropolitan sewage would be worth to the farmer 2s per ton distributed over his arable land, there can still be no doubt that the average value the year round would be reduced to considerably below 1d. by the use of the remainder in large quantities to grass at the less favourable periods of the season.

Adopting the favourable supposition that as high an average as 500 tons of sewage per acre could be utilised on arable land, and that as low an average as 5,000 tons per acre were found sufficient for Italian rye and meadow grass, the important practical questions arise—would the increased productiveness, and increased gross money return per ton, in the former case, justify the extra cost of distribution over a ten-fold area, and to a great extent by pipes and hose and jet instead of by open runs?—or, having regard, not to the greatest amount of produce and of gross money return, but to the greatest profit, per ton of sewage, would it not be far more remunerative to limit the area, and cost of distribution at a certain sacrifice of the productiveness of the sewage?

The probability is, however, that the difference of area required under the two systems would be greater than that here assumed for the purpose of illustration; and then, of course, the difference in the cost of distribution of a given amount of sewage would be still further increased. In fact, to utilise the constituents of the Metropolitan sewage over an area at all corresponding to the rate per acre of the smaller and more productive application to oats at Rugby, it would require more nearly a twenty-fold than a ten-fold area as compared with that of such an application to grass land as above supposed.

The great dilution of the Metropolitan sewage, indeed of town sewage generally, its large daily supply at all seasons, and its greater amount in wet weather when the land can least bear, or at least requires, more water, render it extremely inappropriate for application on a comprehensive scale to arable land for the growth of corn and other ordinary rotation crops. But, apart from these difficulties, if it can only be distributed in small quantities over large areas at such a cost to the farmer as has as yet been pro-

posed, it is indeed vain to hope that any large proportion of the manurial constituents derived from the consumption of human food in our towns, can be distributed over the area from which they came.

A modified proposition is, to make arrangements for delivering the sewage over a large area, and to all crops, so as to obtain a high price per ton for so much as can be applied under the most favourable conditions of the land, the crop, and the season, having in reserve a sufficient tract of grass land to purify and utilise the surplus not so available. But this surplus would be very large, and the largest at those periods of the year when of the least value even for grass land, so that the gross value per ton of sewage the year round would be very much reduced.

Having regard to the cost of distribution, it is probable that a much more profitable mode of utilisation would be, to limit the area by specially adapting the arrangements for the application of at any rate the greater part, if not the whole, to permanent or other grasses, laid down to take it the year round, trusting to the occasional use to other crops within easy reach of the area so commanded, but relying mainly on the periodically broken up rye-grass land, and on the application to arable land of the solid manure resulting from the consumption of the sewage grass, for obtaining other produce than milk and meat by means of sewage.

The question arises—how much land would be requisite for the purification and utilisation of the sewage of a given population on such a plan?

Putting out of view, for the moment, the sanitary consideration of the sufficient purification of the sewage, and the economical one of the manurial value of its constituents, and looking merely to obtaining the largest possible amount of green produce from a given area of land, there is scarcely any limit to the amount of sewage that might be employed, even up to 40,000 or 50,000 tons per acre per annum. But, so far as existing experience furnishes data for a judgment on the point, it may be concluded that the use of about 5,000 tons per acre, judiciously applied to grass land properly laid down to receive it, would, in a great majority of cases, secure the most profitable utilisation. Where, however, the drainage from the sewage land must be turned into a river, other considerations than those relating only to the most profitable utilisation at once arise. Such an application as is here supposed would doubtless ensure a sufficient purification of the water to admit of its being turned into rivers without fear of detriment to fish; whilst, any streams receiving such drainage instead of that direct from the towns would, at any rate, be vastly improved from their previous condition as a water supply for other towns; but, whether or not, when this most important point has to be taken into consideration, the purification would be sufficient with an application of as much as 5,000 tons per acre per annum, is a question which requires the aid of further experience, and

further investigation, to answer satisfactorily, and which may, indeed, receive a different answer in different cases.

Assuming that the excretal matters of each individual of the Metropolitan area are, or will be, on the average, diluted with 100 tons of water per annum, including water supply, rainfall, and subsoil water, 5,000 tons of sewage would be contributed by fifty individuals in a year, and at this rate, a population of 3,000,000 would require, for the purification and utilisation of its sewage, an area of about 60,000 acres annually under irrigation. So far as Italian rye grass were grown, it might be estimated that the land devoted to it would one year in three be broken up, and some other crop be grown upon it, and to a corresponding extent the area laid down for irrigation would require to be extended beyond the 60,000 acres supposed. Then again, it is obvious that the manure produced by the consumption of the sewage grass must either be re-distributed by means of water, in which case the area under actual irrigation would be again increased, or, if collected and used in the solid form, it would be appropriate for application to arable land, and so to the growth of corn and other products; and it is obvious that for the most profitable utilisation in this way of the manure derived from the consumption of the sewage grass, such arable land would require to be either within the area laid down for irrigation, or so near its limits as to reduce the cost of carriage as far as possible.

The experiment with oats above referred to, and that with wheat made by the Chairman of the Commission, the Earl of Essex, the results of which his lordship gave in his evidence before the Sewage Committee of 1862, are the only cases in which exact quantitative results have been recorded of the effects of sewage applied to corn crops.

In the case of the experiment of the Earl of Essex, nothing is known of the strength of the sewage, and nothing is recorded of the characters of the season.

In the case of those with oats at Rugby, as already observed, the sewage was stronger than the average of that of Rugby, and much stronger than the average of that of the Metropolis, the weather was unusually dry at the time of the application, and the season was upon the whole one of very extraordinary productiveness, and, under these conditions, a very high gross return was obtained for a given amount of sewage. Judging from the results of the Earl of Essex, it is probable that the circumstances, both as to the strength of the sewage and character of the season, were in his case also unusually favourable.

At any rate, these isolated results, the one obtained under conditions known to be far above an average character, and the other under entirely unknown conditions, are obviously quite inappropriate as the basis for general conclusions as to the probable average results obtainable on the application of sewage to arable land for corn and other rotation crops.

It is, indeed, desirable that systematic trials should be made with different corn and other rotation crops, through several consecutive seasons, and that the results should be accurately recorded for the guidance of the public.

Although there is still wanting evidence of an exact and quantitative kind upon which to found estimates of the probable average results obtainable over various seasons, on the application of given quantities of sewage, of known strength, to corn and other rotation crops, yet evidence of common experience as to the applicability, in a practical or economical point of view, of sewage to such crops, is by no means wanting. The most extensive and systematic trials have been made at Rugby, Watford, and Alnwick.

At Rugby, the sewage from a population of between 6,000 and 7,000 individuals is collected in a receiving tank, from which it is pumped by a 12-horse power engine through iron pipes which are laid down for the distribution over 470 acres of mixed grass and arable land. These arrangements have been in existence for about 11 years. About 190 acres of the land so piped have, from the commencement, been held by J. A. Campbell, Esquire. But he has gradually limited the area of application, until, during the last few years, he has abandoned the use of the hose and jet, excepting occasionally on a small scale, and confined the application almost exclusively to from 12 to 20 acres of meadow and Italian rye grass. The greater part of the remainder of the 470 acres was, for some time previous to 1861, held by Mr. Berry Congreve, who, after trying sewage on arable as well as grass land, was glad to give up his holding, after having sustained considerable loss. The present tenant of the sewage works, and of the land formerly held by Mr. Congreve, is Mr. Bicknell Mullins, who entered into the occupation in 1861, and although he had between 250 and 300 acres laid down for the application of sewage to crops generally, and by hose and jet, he in practice confines it to about 100 acres of grass land, and applies it almost entirely by open runs.*

The result at Rugby is, then, that after about eleven years of practical experience, with arrangements adapted for the application of sewage to arable land, and to all crops, its use to any other crops than meadow and Italian rye grass forms no part of the general system adopted, and is, in fact, entirely exceptional.

In the neighbourhood of Watford, the Earl of Essex laid down pipes for the application of the sewage of the town by hose and jet to about 210 acres of mixed arable and grass land. The result which his lordship obtained on the application of only 134 tons of sewage to an acre of wheat has frequently been held to be conclusive proof of its applicability in small quantities, over large areas, and to all crops. But, in the evidence given by his lordship before the Sewage Committee of 1862, he stated, very

* Since the above was in type Mr. Mullins has informed us that during the dry season of 1864 he applied sewage to about four acres of roots, and apparently with good results.

emphatically, that his great error had been the piping of too much land; that the sewage of Watford, derived from a population of about 4,000, was not sufficient for more than about 60 or 70 acres; that he required 5,000 tons per acre for 10 acres of rye-grass; and that applying the remainder to 35 acres of meadow, he really had none to spare for wheat. It should be added, that, since the date of this evidence, the area of application has been still further contracted.

In other words, the result at Watford is, that although the abandonment of one acre of rye-grass would set free sewage enough for nearly 40 acres of wheat, if only applied at the rate which yielded the large profit which has been so frequently quoted, yet his lordship's practical experience has led him to prefer the application to the one acre of rye-grass, rather than to the nearly 40 acres of wheat.

In the neighbourhood of Alnwick, the Duke of Northumberland, some years ago, put down machinery and piping for the distribution of the sewage of from 6,000 to 7,000 individuals, over about 270 acres of mixed arable and grass land. It was applied in small quantities to various rotation crops, and in larger quantities to grass; but after a very short time the tenants who had the free use of the sewage for the cost of its application, entirely abandoned it; and the Bailiff of the district, who reports the failure, expresses his opinion strongly against the applicability of sewage to arable land. The failure at Alnwick has been attributed by those connected with the undertaking, to the great dilution of the sewage; and the analyses recorded of it would indicate a composition even much below that of the probable average of the Metropolitan sewage. But Mr. Rawlinson, who directed the sewerage arrangements at Alnwick, states, that not only are water-closets universal, but that the supply of water from all sources is certainly very much lower per head of the population contributing to the sewage there than in the case of the Metropolis; and it is obvious that, if this be the case, the average sewage must be in a corresponding degree the stronger.

At Edinburgh sewage has been applied to some portions of grass land for about 200 years, to a considerable portion for more than 60, and to most of the land now under irrigation, amounting to about 395 acres, for more than 30 years. It is there that larger amounts of sewage are applied per acre than anywhere else, and it is there that larger amounts of produce are obtained per acre than anywhere else. There is, however, no doubt, that at Edinburgh there is not only great waste of manurial constituents, but very imperfect purification of the sewage. The distribution is entirely by means of open runs. In two instances arrangements have been made for raising the sewage, by pumping, an inconsiderable number of feet; but it has been found that the cost has been too great to allow a sufficient quantity to be *applied per acre*, and hence the application in this way has been *much limited*, if not on some portions of the land entirely aban-

done. The application to ordinary rotation crops, on arable land, forms no part of the system adopted at Edinburgh.*

Next to Edinburgh, the attempt to utilize sewage on a large scale which has been the most successful so far as the amount of produce obtained per acre is concerned, is that of Mr. Marriage in the neighbourhood of Croydon, where about 250 acres of meadow and Italian rye grass annually receive an amount of sewage which averages rather more than 6,000 tons, and represents the excretal matters of between 60 and 70 persons per acre per annum. As, however, the fluid is always passed over several portions of land in succession, by which means a considerable portion is used on an average about $2\frac{1}{2}$ times over, it results that each acre receives annually $6,000 \times 2\frac{1}{2} = 15,000$ tons of fluid—less the amount which evaporates or passes off below the drains which collect and carry it off from one portion to be utilised on another. An enlargement of the area is, indeed, contemplated, which, notwithstanding the rapid increase of the population of the neighbourhood, will, if carried out as proposed, somewhat reduce the amount of fluid and of excretal matters available per acre below the quantities above stated. Mr. Marriage has not yet applied sewage in any systematic manner to arable land; but he was intending to try its effects upon root-crops during the past season (1864).†

In attempting to estimate by the aid of the evidence afforded by these various trials on a large scale, carried out by practical men with a view to profit, the value to the farmer of a ton of town sewage, we may, on account of the conditions above stated, exclude the Edinburgh results from our consideration.

At Croydon, again, the undertaking is of too recent establishment, the results have been obtained over too few seasons, and the present contract was made, on either side, under such disadvantages or uncertainty, that the experience there does not provide the adequate data for such an estimate. It may be observed, however, that after deducting £4 rental from the estimated gross value of the produce per acre at present prices, the gross return is, so far as can be calculated, with Italian rye grass from $\frac{3}{4}d.$ to $1d.$, and with meadow grass from $\frac{1}{2}d.$ to $\frac{3}{4}d.$ for each ton of sewage employed. But there can be little doubt, that if the supply of such produce were very greatly increased, the present market price would not be maintained.

At Rugby, where for about eleven years arrangements have been made for the distribution of small quantities of sewage over a large area, and to all crops, and where the sewage is much stronger than that of the Metropolis, the cost to the tenants averages about $\frac{3}{4}d.$ per ton delivered at the hydrants in the fields. Yet, both the present tenants have been glad, rather than incur the loss of using the sewage themselves at that cost, to get rid of it for the purposes of these experiments, at rates which,

* For further particulars relating to the Edinburgh sewage meadows, see Appendix No. 2. p. 198, et seq.

† For further details relating to Croydon, see Appendix No. 3. p. 202, et seq.

though three times as high during the six summer as during the six winter months, have averaged the year round scarcely, but very nearly, 1*d.* per ton at the hydrants.

Lastly on this point, in his evidence before the Sewage Committee of 1862, the Earl of Essex stated, as the result of his experience, which it will be remembered included the very favourable result with wheat, that in his opinion sewage would not be profitable to the farmer unless he could have it at from $\frac{1}{2}$ *d.* to $\frac{3}{4}$ *d.* per ton.

The experiments at Rugby to which this Report refers, having been conducted on feeding meadow land of more than average quality, the produce without sewage was doubtless considerably more than would be obtained from the average of such land as is likely to be devoted to the growth of grass by means of sewage on the large scale. For this reason, and also on account of the less perfect purification and utilisation of the sewage than would be attained where a large tract were so laid down as to allow of the passage of the fluid from one plot over a second, and so on, until it were properly exhausted, the amounts, and value, of the increase estimated according to the actual results of the experiments as due to the application of given quantities of sewage, are probably below those which would be attainable under good management in actual practice on the large scale.

Reviewing the whole of the results, both of the experiments and of the experience of common practice on the subject hitherto, with due regard to the circumstances under which they were obtained, and having regard also to both urban and rural interests, it is considered that an application of about 5,000 tons of sewage per acre per annum, to meadow or Italian rye grass, will, in a great majority of cases, prove to be the most profitable mode of utilisation. It is at the same time considered pretty certain that the farmer would not pay $\frac{3}{4}$ *d.*, and even very doubtful whether he could afford to pay $\frac{1}{2}$ *d.* per ton, the year round, for sewage of the average strength of that of the Metropolis (excluding storm water) delivered on his land.

SUMMARY.

The results of the whole inquiry may be briefly enumerated as follows:—

1. As there is a daily supply of sewage the year round, which, on sanitary and engineering grounds, it is essential to dispose of as soon as it is produced, and as passing it over land is the best mode both of purifying and utilising it, it should be employed for purposes of irrigation, and be applied in winter, when of comparatively little value, as well as in summer, when of more.

Results obtained on the Application of Sewage to Meadow and Italian Rye Grass.

2. By the application of sewage to grass land during the winter months a very early cut or bite of green food may be obtained, but the amount of increased produce due to the winter application is comparatively small for the amount of sewage employed.

3. By means of sewage irrigation the period during which an abundance of green food was available was extended considerably at the end as well as at the beginning of the season, and the more so the larger the quantity of sewage applied, almost up to the highest amount employed—namely, 9,000 tons per acre.

4. One of the experimental fields gave much less produce per acre without sewage than the other, and analysis showed its soil to be much less naturally fertile; but it gave fully as much produce per acre under the influence of liberal dressings of sewage as the naturally much more fertile soil.

5. Taking the average over three years, and in the two fields, the amount of produce obtained without sewage was about $9\frac{1}{2}$ tons, of green grass per acre per annum, equal about 3 tons of hay; and with 3,000, 6,000, and 9,000 tons of sewage per acre per annum the amounts were, respectively, about $22\frac{1}{2}$, $30\frac{1}{2}$, and $32\frac{1}{2}$ tons of green grass—equal respectively (reckoned according to the percentage of dry substance in each) about 5, $5\frac{1}{2}$, and $6\frac{1}{2}$ tons of hay.

6. The largest quantities of produce per acre were obtained in the third year of the experiments, and with 9,000 tons of sewage per acre per annum; namely, in one field 35 tons, and in the other 37 tons of green grass, equal respectively about 6 tons $12\frac{1}{2}$ cwt., and 7 tons 1 cwt., of hay.

7. The average increase obtained for each 1,000 tons of sewage was—when 3,000 tons per acre per annum were applied, about 5 tons of green grass; when 6,000 tons were applied, 4 tons $2\frac{1}{2}$ cwt.; and when 9,000 tons were applied, 3 tons $3\frac{1}{2}$ cwt. of green grass.

8. The amount of produce per acre was the greater the greater the quantity of sewage applied, up to 9,000 tons per acre; but the amount of increase of produce obtained for a given amount of sewage was the less where the greater amounts were applied.

9. Experiments with rye grass were made in one season only, sewage was not applied until the end of April, and comparatively small quantities were put on. The results so obtained indicated much about the same amount of increase of produce for a given amount of sewage as with meadow grass.

Results obtained with fattening Oxen.

10. When cut and given to fattening oxen tied up under cover, more sewage than unsewaged grass, reckoned in the fresh or green state, was both consumed by a given weight of animal within a given time, and required to produce a given weight of increase; but, of real dry or solid substance, less of that of the sewage than of the unsewaged grass was required to produce a given effect.

11. When cut grass was given alone the result was very unsatisfactory; but when oilcake was given in addition the amount of increase upon a given weight of animal within a given time, and for a given amount of dry substance of food consumed, was far short of the average result obtained when oxen are fed under cover on a good mixed diet.

12. The money return, whether reckoned per acre or for a given amount of sewage, was much less with fattening oxen than with milking cows.

Results obtained with milking Cows.

13. When cows were fed on unsewaged, or sewaged grass, as much as they chose to eat, a given weight of the animal was more productive, both of milk and increase, but especially of milk, on the unsewaged than on the sewaged grass.

14. From a given weight of unsewaged grass, reckoned in the fresh or green state, more milk was produced than from an equal weight of fresh sewaged grass; but a given weight of the dry or solid substance supplied in sewaged grass was on the average more productive than an equal weight supplied in unsewaged grass.

15. The milk producing quality of the grass was very different in different seasons, and at different periods of the same season. It was very inferior in the wet and cold season of 1862, and toward the close of the seasons as compared with the earlier periods. It appears probable that Italian rye grass deteriorates less toward the end of a season than meadow grass. On the average, about six parts by weight of fresh grass yielded one part by weight of milk.

16. By the aid of sewage, the time that an acre would keep a cow, and the amount of milk yielded from the produce of an acre were increased between three and four-fold.

17. So far as the results of the experiments afford the means of judging, it is estimated that with an application of about 5,000 tons of sewage per acre per annum to meadow land, an average gross produce of not less than 1,000 gallons of milk per acre per annum may be expected.

18. In experiments conducted with Italian rye grass (but in one season only), more milk was obtained by the use of a given amount of sewage applied to it than to meadow grass.

19. With an application of about 5,000 tons of sewage per acre per annum, an average gross return of from 30*l.* to 35*l.* per acre in milk at 8*d.* per gallon, may be anticipated.

Composition of the Rugby Sewage.

20. The mean of 93 analyses, of as many samples, of the Rugby sewage, collected over a period of 31 months, shows 6 grains of ammonia, and 87½ grains of total solid matter, per gallon; equal to 207½ lbs. of ammonia, and 2,803 lbs. of total solid matter per 1,000 tons. Or, taking the mean of the average composition fixed by the analyses for each of the 31 months instead of the direct mean of the total 93 analyses, the average contents would be almost exactly 7 grains of ammonia, and 92½ grains of total solid matter per gallon; equal to 224 lbs. or 2 cwts. of ammonia, and 2,960 lbs., or about 26½ cwts. of total solid matter, per 1,000 tons.

21. Although each sample analysed was the mixture of portions taken every two or three hours for several days together, the variation in composition at different times was very great; the amount of ammonia varying in the different mixed samples from $2\frac{1}{2}$ to about $15\frac{1}{2}$ grains per gallon, or from $81\frac{1}{2}$ to $500\frac{1}{2}$ lbs. per 1,000 tons, whilst the total solid matter varied from about $37\frac{1}{2}$ to about 270 grains per gallon, or from 1,203 to 8,637 lbs. per 1,000 tons.

22. 1,000 tons of the average sewage of Rugby represent the excretal and other matters of from 17 to 18 average individuals of a mixed population of both sexes and all ages for a year, and contain ammonia equal to that in from 11 to 12 cwt. of Peruvian guano; or, about 1,700 tons of such sewage would contain nitrogen reckoned as ammonia equal to that in 1 ton of Peruvian guano.

23. It is estimated that there are at Rugby, including rainfall, &c., on the average from 55 to 60 tons of sewage per head of the population per annum.

24. Judging from the average composition of the Rugby sewage, and of various crops, it is concluded that potass would be more likely than phosphoric acid to become deficient where town sewage was applied constantly to grass-land, whilst phosphoric acid would be more likely to become deficient than potass if it were applied to the ordinary crops of rotation.

Estimated average Composition of the Metropolitan Sewage.

25. There is as yet no record of the analysis of any samples or sample of sewage collected under circumstances fairly to represent the average Metropolitan sewage either with or without rainfall and subsoil water.

26. It is estimated that the Metropolitan sewage amounts on the average to about 60 tons without, and probably to about 100 tons with, rainfall and subsoil water, per head of the population per annum.

27. It is estimated that, including human excretal and other matters, there are annually contributed to the Metropolitan sewage about $12\frac{1}{2}$ lbs. of ammonia per head of the mixed population of both sexes and all ages.

28. Reckoned according to the currently adopted trade prices of the several constituents, taking dry and portable manures as standard, the total annual value of the manurial constituents contributed to the sewage, supposing them to be extracted and dried, would amount to 8s. 4d. per head of the population.

29. Accordingly, in the dry weather sewage of the Metropolis, reckoned at 60 tons per head per annum, there will be about $6\frac{1}{2}$ grains of ammonia per gallon, and the manurial constituents in 1 ton, if extracted and dried, would be worth about $1\frac{1}{2}$ d.; and in the average sewage with rainfall, &c., reckoned at 100 tons per head per annum, there will be scarcely 4 grains of ammonia per gallon, and the total manurial constituents in 1 ton will have an estimated value of 1d.

30. 1,000 tons of the average Metropolitan sewage without rainfall, reckoned at 60 tons per head per annum, represent the excretal and collateral manurial matters from nearly 17 average individuals, and contain ammonia equal to that in about 11 cwt. of Peruvian guano; and 1,000 tons with rainfall, reckoned at 100 tons per head per annum, represent the manurial matter from 10 average individuals, and contain ammonia equal to that in about $6\frac{1}{2}$ cwt. of Peruvian guano. In other words, about 1,800 tons of the average Metropolitan sewage without, and about 3,000 tons of the average sewage with rainfall, &c., would contain nitrogen reckoned as ammonia equal to that in 1 ton of Peruvian guano.

31. The value of the total manurial constituents in sewage reckoned according to the currently adopted trade prices of the several constituents, taking dry and portable manures as the standard, is pretty exactly indicated by putting a value of 8d. on every lb. of ammonia, or by giving a value of one farthing per ton for every grain of ammonia per gallon of the sewage. But this theoretical value, according to composition and the trade prices of the constituents, cannot, of course, be taken as directly indicating the value realized, or realizable, by the agricultural utilisation in various ways, of sewage of different strengths.

32. It is very desirable that as soon as the Main Drainage system is sufficiently advanced and in practical working, competent persons should be appointed to undertake the gauging, sampling, and analysis, of the Metropolitan sewage, in such manner as satisfactorily to determine its average composition in the condition in which it will have to be dealt with in any plan of utilisation.

Composition of the Drainage Water (Rugby).

33. Analyses of the drainage water passing from the experimentally sewage-irrigated land at Rugby showed that those constituents which are of the most value, because the most liable to become relatively exhausted, had been the most efficiently retained by the soil, but that the water still contained a considerable amount of valuable manurial matters, besides a large quantity of other substances less important as manure, but affecting the purity of the water.

34. When large quantities of sewage are applied to grass land the arrangements should be such as to allow of the water being used more than once, so that both the utilisation and the purification may be as complete as possible.

Chemical Composition of the Grass.

35. The sewage-drenched meadow grass, as cut and given to the animals, contained a less proportion of dry or solid substance than the unsewaged; and the grass cut during the later portion of the season (both unsewaged and sewage-drenched) contained less solid matter than that cut during the more genial periods of growth.

36. Italian rye grass, in the condition as cut, was also found to be more succulent and to contain less solid matter when grown with sewage than without it; but the proportion of dry substance diminished less as the season advanced in its case than in that of the meadow grass.

37. The proportion of nitrogenous substance (and also of impure waxy or fatty matter) was much greater in the solid matter of the sewaged, than in that of the unsewaged grass. The proportion of nitrogenous substance was also much higher in the solid matter of the grass grown towards the end than earlier in the season. The proportion of indigestible woody fibre was much about the same in the dry substance of the unsewaged and of the sewaged grass. It progressively diminished as the season advanced, and was generally lower in the dry substance of the Italian rye than in that of the meadow grass.

38. A given amount of the dry substance of grass grown in a cold and wet season, or during the cold and wet periods of the year, generally contains more nitrogenous substance, but is less productive than that of grass grown in more genial weather.

39. The greater productiveness in milk and increase of a given amount of the solid matter of the sewaged grass appears to depend more on a favourable condition of maturation, digestibility, and assimilability, of the constituents, than on the actual percentage amount of any of those determined, and above enumerated.

Effects of Sewage on the mixed Herbage of Grass Land.

40. The effect of sewage irrigation on the mixed herbage of grass land is to develop the Gramineaceous plants chiefly, nearly to exclude the Leguminous, and to reduce the prevalence of miscellaneous or weedy plants, but much to encourage individual species.

41. Among the grasses which have been observed to be the most encouraged by sewage are (according to locality or other circumstances) rough meadow grass, couch grass, rough cock's foot, woolly soft grass, and perennial rye grass; two or three only remaining in any considerable proportion after sewage has been liberally applied for some years.

42. The produce of sewage irrigated meadows being generally cut or grazed very young, the tendency which the great luxuriance of a few very free growing grasses has to give a coarse and stemmy later growth is not an objection as in the case of meadows left for hay; a given weight of the dry or solid substance of the more simple sewaged grass being, when consumed green, more productive than an equal weight of that of the more complex unsewaged herbage.

Composition of the Milk from the unsewaged and the sewaged Grass.

43. Although more milk was obtained from a given weight of the dry or solid substance of sewaged than of unsewaged grass, there was comparatively little difference in the composition or

richness of the milk from the two kinds of grass. That from the sewage grass was, however, slightly the less rich, containing somewhat less of casein, butter, sugar, and total solid matter (though more mineral matter) than that from the unsewaged.

44. When oilcake was given with the grass (whether sewage or unsewaged) the richness of the milk was notably increased.

Results obtained on the Application of Sewage to Oats.

45. In an experiment with oats in which $135\frac{1}{2}$ tons of sewage were applied per acre, the gross value of the increased produce amounted to more than 5d. per ton of the sewage employed, or to about three times the market value of the constituents of the sewage, supposing them to have been extracted and dried; and in another experiment in which 510 tons were applied per acre, the gross value of the increased produce amounted to about $1\frac{1}{2}$ d. per ton of the sewage employed.

46. In the experiment with the smaller quantity of sewage the supply of water was equivalent to something under an additional $1\frac{1}{2}$ inch of rain at the critical period of growth, and in that with the larger amount to about 5 inches, which proved to be a great excess at the period of the season at which it was applied, there being an over production of straw, and the crop being much laid. Both experiments were made in the unusually productive season of 1863, and with sewage of about double the average strength of that of the Metropolis, which was applied during a period of very dry weather. It is obvious, therefore, that the results were quite exceptional, and cannot be taken as indicating what might be expected from the application of small quantities of sewage to corn crops on different soils, and on the average of seasons.

47. It is probable that 500 tons of sewage per acre is more than would be appropriate to arable land otherwise treated in the ordinary way, taking the average of soils and seasons; and it is certainly more than would be appropriate for heavy lands, and for wet seasons.

General Conclusions.

48. To obtain a maximum amount and gross value of produce from a given amount of sewage, it should be applied in small quantities per acre, and in dry weather; but the great dilution of town sewage, its large daily supply at all seasons, and its greater amount in wet weather, when the land can least bear, or least requires, more water, render it quite inappropriate for application on a comprehensive scale to arable land for corn and other ordinary rotation crops.

49. Supposing arrangements were made for distributing sewage over a sufficiently large area to command a full value, both as manure and as water, at the most favourable periods of the year, the cost of main distribution would be very great, the application to the arable land would require to be chiefly by the expensive means of piping and hose and jet, instead of open runs, and but a

small proportion of the total sewage could be so used, leaving the remainder to be applied in large quantities to grass-land, at the less favourable periods of the year, and, of course, to realize a much lower value.

50. Having regard to the cost of distribution, it is probable that the most profitable mode of utilisation would be to limit the area by specially adapting the arrangements for the application of the greater part, if not the whole, to permanent or other grasses laid down to take it the year round, trusting to the occasional use to other crops within easy reach of the line or area so commanded, but relying mainly on the periodically broken up rye-grass land, and on the application to arable land of the solid manure resulting from the consumption of the sewaged grass, for obtaining other produce than milk and meat, by means of sewage.

51. It is probable that about 5,000 tons of sewage per acre, judiciously applied to grass-land properly laid down to receive it, would, in a great majority of cases, secure the most profitable utilisation.

52. Supposing an application of 5,000 tons of sewage per acre per annum to grass land, the purification of the water would doubtless be sufficient to admit of the drainage being turned into rivers without fear of detriment to fish; whilst any streams receiving such drainage instead of that direct from the towns would at any rate be vastly improved from their previous condition as a water supply; but whether the purification would be sufficient with such an application is a question which requires further experience and investigation to answer satisfactorily, and which will probably receive a different answer in different cases.

53. Assuming that the average dilution of the Metropolitan sewage, including rainfall and subsoil water, will amount to 100 tons per head per annum, 5,000 tons would represent the excretal and other matters of 50 average individuals; and a population of 3,000,000 would require about 60,000 acres constantly under irrigation.

54. The only records of exact quantitative results obtained on the application of town sewage to corn crops are those of the experiments of the Earl of Essex on wheat, and those of the experiments with oats at Rugby given in this Report, and in both cases the increase of produce represented a very high gross money return per ton of sewage employed. The circumstances of the experiments at Rugby were, however, quite exceptional; and, where the most extensive trials of the application of sewage to corn crops have been made with a view to profit, namely, at Watford, Rugby, and Alnwick, the practice has been abandoned; whilst neither at Edinburgh nor Croydon, where the best results have been obtained with grass, does the application to corn and other rotation crops constitute a part of the general system adopted.

55. Judging both from the results of the experiments, and from the experience of common practice, it is considered that the most

APPENDIX.

APPENDIX No. 1.

DETAILED RECORDS relating to the EXPERIMENTS made at RUGBY.

TABLE I.

DETAILED RECORD of the SEWAGE applied to PERMANENT GRASS LAND.
SECOND SEASON 1861-2.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average time taken to fill Gauge-tank (3'216 tons).	Time of application.	Sewage applied (calculated per acre).			Average time taken to fill Gauge-tank (3'216 tons).	Time of application.	Sewage applied (calculated per acre).		
			Plot 2. (Area '93375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area 1'00512 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99388 acre.)	Plot 4. (Area 1'00019 acre.)
Nov. 4	Mins.	H. M.	Tons.	Tons.	Tons.	Mins.	H. M.	Tons.	Tons.	Tons.
5	15'50	2 45	34'06	9'50	2 0	41'14
6	13'06	5 0	71'07	15'00	4 0	..	51'67	..
7	14'33	6 15	..	90'41	..	11'50	2 0	..	33'70	..
8	13'00	7 0	103'88
9	14'16	11 15	152'52	10'50	2 0	36'75
10	13'83	10 0	138'81	9'50	2 0	41'14
11	14'67	10 0	..	141'31	..	13'50	3 0	..	43'06	..
12	11'50	2 0	..	33'70	..
13	12'07	7 15	110'39
14	14'00	6 15	86'68	10'50	2 0	36'75
15	13'00	4 0	59'07
16	9'50	2 0	..	40'79	..
17	11'50	2 0	33'98
18	11'50	2 0	33'55
19	14'50	9 15	122'47
20	15'16	9 45	..	133'32	..	13'50	5 0	..	71'76	..
21	14'07	5 0	65'75
22	14'67	6 0	79'42
23	14'34	4 0	53'55
24	14'16	9 45	132'19
25	15'00	9 45	..	134'75	..	14'50	5 0	..	66'81	..
26	15'00	5 0	64'31
27	15'34	6 0	75'95
28	14'67	4 0	52'34
Tot.	(14'42)	114 0	313'12	409'79	745'01	(12'61)	59 15	116'26	341'49	451'38
Dec. 2	14'00	9 0	123'41
3	14'16	9 45	..	142'74	..	13'67	5 0	71'47
4	14'50	5 0	66'52
5	12'34	6 0	75'95
6	15'34	4 0	50'06
7	14'34	5 0	66'94	13'34	5 0	72'31
8	14'00	5 0	..	74'04
9	14'00	5 0	68'56
10	15'00	9 15	118'39
11	15'34	4 0	50'63	13'67	5 0	..	70'87	..
12	14'00	5 45	..	85'14	..	13'00	5 0	74'20
13	15'30	8 15	100'24
14	15'00	9 15	..	127'84	..	13'34	5 0	..	72'62	..
15	13'50	4 30	64'31
Tot.	(14'66)	80 15	126'58	429'76	527'60	(13'56)	34 30	71'47	143'49	277'34

Table I.—continued.
Detailed Record of the Sewage applied to Permanent Grass La
Second Season 1861-2.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average Time taken to fill Gauge-tank (3'216 tons).	Time of applica-tion.	Sewage applied (calculated per acre).			Average Time taken to fill Gauge-tank (3'216 tons).	Time of applica-tion.	Sewage appl (calculated per		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area 1'00512 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99588 acre.)	
Jan. 1	Mins.	H. M.	Tons.	Tons.	Tons.	Mins.	H. M.	Tons.	Tons.	
6	15'00	3 30	45'31
7	15'16	9 45	123'47
8	15'16	9 45	..	133'32	..	12'00	4 45	77'35
9	12'00	5 0
10	15'00	5 45	74'43
11	15'34	4 0	50'06
12	14'50	9 45	129'09
14	14'07	9 45	..	137'78	..	12'00	5 0	..	90'73	..
15	12'34	5 0
16	14'34	6 0	81'24
17	15'34	4 0	50'06
21	21'25	5 45	..	56'09
22	16'34	5 0	59'42
23	16'00	5 0	59'99
27	15'53	9 45	118'24
28	15'50	9 45	..	130'40	..	12'34	5 0	..	78'51	..
29	15'50	5 0	62'64	12'50	5 0
30	15'50	5 0
31	14'07	4 0	52'34
Total -	(15'41)	103 30	323'04	457'59	583'25	(12'20)	29 45	77'35	159'24	
Feb. 3	22'16	11 15	97'46
4	15'07	11 30	..	152'14	..	11'07	7 45	129'77
5	12'50	3 45
6	16'34	11 15	133'17
10	16'00	10 45	124'32
11	16'16	11 30	..	147'52	..	11'50	5 45	..	96'83	..
12	12'34	5 45
13	18'00	11 30	122'65
17	16'34	4 45	55'81
18	16'23	11 30	..	141'65	..	12'34	5 45	..	90'28	..
19	12'00	5 45
20	16'07	11 45	126'86
24	16'34	11 15	133'17
25	17'53	11 30	..	123'71	..	11'50	5 45	97'70
26	12'34	5 45
28	12'00	3 0	32'36
30	16'00	7 15	86'99
Total -	(17'03)	123 45	169'22	575'02	751'57	(11'98)	46 0	227'47	187'16	
Mar. 3	16'00	11 15	134'98
4	18'00	11 30	..	132'44	..	12'00	5 45	..	92'84	..
5	11'34	5 45
6	16'34	6 45	90'51
7	17'34	3 45	41'52
17 (8 tanks*)	11 0	25'60
21	16'00	11 30	..	149'00	..	12'00	5 45	..	92'84	..
22	11'33	5 45
23	14'50	4 15	56'91
24	12'07	5 45	80'75
25	12'00	4 45
26	11'33	11 30	..	126'35	..
28	12'00	6 45	109'91
29	14'07	11 0	143'96	..	4 15
30	12'07	11 30	..	174'40
31	12'33	6 45	93'23
31	14'00	3 45	51'42
31	12'16	11'30
Total -	(16'73)	96 45	235'44	455'84	475'22	(11'55)	61 45	109'91	374'08	

* When the number of tanks is given, the flow was too slow and irregular to estimate average time taken to fill the gauge-tank, and therefore the actual number of tanks counted.

Table I.—continued:
Detailed Record of the Sewage applied to Permanent Grass Land.
Second Season 1861-2.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average Time taken to fill Gauge-tank (3' 216 tons).	Time of applica-tion.	Sewage applied (calculated per acre).			Average Time taken to fill Gauge-tank (3' 216 tons).	Time of applica-tion.	Sewage applied (calculated per acre).		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '99381 acre.)	Plot 4. (Area 1' 00512 acre.)			Plot 2. (Area '99375 acre.)	Plot 3. (Area '99383 acre.)	Plot 3. (Area 1' 00019 acre.)
	Mins.	H. M.	Tons.	Tons.	Tons.	Mins.	H. M.	Tons.	Tons.	Tons.
April 1	11' 50	11 30	..	193' 76	..
2	12' 33	6 45	106' 97
3	15' 16	11 30	145' 63	11' 33	4 15	72' 37
4	15' 50	11 30	..	153' 89
5	15' 67	6 45	83' 64
6	15' 00	2 45	35' 20
7	11' 50	10 15	167' 58
8	12' 16	11 30	..	183' 24	..
9	12' 33	5 45	91' 12
10	15' 20	10 15	129' 46	11' 50	3 45	62' 01
11	14' 75	7 45	..	108' 92
12	15' 67	10 15	125' 58
13	12' 33	10 30	164' 29
14	12' 33	11 30	..	180' 71	..
15	12' 33	6 45	106' 97
16	11' 67	4 30	74' 39
17	17' 60	10 0	109' 08
18	15' 67	4 0	49' 56
19	15' 33	6 45	..	91' 28	..	12' 16	10 30	166' 59
20	12' 33	11 30	..	180' 71	..
21	12' 33	6 45	106' 97
22	11' 50	3 30	58' 71
23	16' 00	11 0	131' 98
24	16' 00	11 30	..	140' 00
25	16' 67	6 45	78' 62
26	16' 67	3 45	43' 19
27	12' 67	4 45	72' 83
28	13' 50	11 30	..	165' 05	..
29	13' 67	6 45	96' 49
30	13' 00	2 30	87' 10
Total ..	(15' 75)	114 30	211' 82	503' 00	720' 12	(12' 26)	144 45	508' 52	903' 47	876' 27
May 1	17' 67	4 45	51' 61
2	20' 67	4 45	44' 62
3	21' 00	6 45	..	66' 63
4	20' 67	10 30	97' 52
5	12' 16	10 30	163' 93
6	13' 50	4 30	..	64' 59	..
7	13' 67	5 45	81' 15
8	13' 33	5 45	84' 20
9	25' 67	13' 33	5 0	72' 36
10	25' 67	11 30	80' 00
11	25' 00	11 30	..	92' 87
12	23' 33	6 15	48' 54
13	..	4 15	34' 97
14	15' 40	8 45	109' 62
15	14' 67	5 45	..	75' 94	..
16	17' 00	6 0	68' 09
17	24' 50	16' 00	4 45	58' 01
18	16' 00	11 30	90' 11	12' 67	6 45	102' 78
19	15' 50	11 30	..	140' 00
20	16' 00	4 0	50' 11
21	..	6 45	80' 99
22	11' 83	10 30	171' 23
23	12' 67	5 45	..	87' 93	..
24	13' 33	5 45	83' 22
25	15' 00	11' 50	3 0	50' 97
26	16' 83	11 30	116' 20	12' 33	5 45	89' 97
27	17' 00	11 30	..	141' 65
28	17' 00	5 45	65' 68
29	..	4 45	53' 64
30	11' 83	10 30	171' 23
31	11' 33	11 30	196' 67
..	11' 67	5 45	96' 28
..	11' 33	5 45	97' 91
..	..	11 30	..	131' 20	110' 39
..	..	6 45	72' 82
..	..	3 45	41' 54
Total ..	(19' 56)	161 0	281' 77	551' 35	762' 97	(12' 90)	127 45	280' 55	425' 13	1261' 40

Table I.—continued.
Detailed Record of the Sewage applied to Permanent Grass Land.
Second Season 1861–2.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average time taken to fill Gauge- tank (3'216 tons).	Time of applica- tion.	Sewage applied (calculated per acre).			Average time taken to fill Gauge- tank (3'216 tons).	Time of applica- tion.	Sewage applied (calculated per acre).		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area '1'00512 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99588 acre.)	Plot 4. (Area '1'00019 acre.)
	Mins.	H. M.	Tons.	Tons.	Tons.	Mins.	H. M.	Tons.	Tons.	Tons.
June 2	13'00	4 45	70'49
3	12'00	7 45	..	125'14	..
4	12'00	3 45	60'29
5	13'33	5 45	84'29
6	18'50	11 30	117'43	13'00	5 45	85'33
7	16'53	11 30	..	141'65
8	17'00	6 45	77'10
9	16'33	3 45	44'09
10	10'50	4 0	..	73'81	..
11	12'00	6 45	108'52
12	14'00	9 15	..	128'02	..
13	16'53	11 30	181'18	12'67	5 45	88'68
14	18'00	2 0	..	23'03	..	13'00	5 45	85'33
Total -	(17'30)	47 0	77'10	164'68	292'70	(12'61)	59 15	172'97	326'97	409'96
July 5	36'50	4 0	21'04	17'75	7 0	76'08
6	14'00	11 0	151'58
7	11'33	11 0	..	188'11	..
8	11'17	12 0	209'92
9	16'00	12 0	143'98	12'00	12 0	192'92
10	16'50	12 0	..	150'77	..	12'00	12 0	..	193'76	..
11	15'67	11 0	136'30	11'83	11 0	179'39
12	15'67	11 0	134'76	11'67	11 0	181'85
13	15'50	12 0	..	160'40	..	11'67	12 0	..	199'24	..
14	15'50	12 0	150'33	12'33	12 0	190'17
15	16'16	12 0	142'56	13'50	12 0	171'49
16	15'60	11 0	..	146'17	..	11'60	11 0	..	183'74	..
17	15'33	11 0	137'75	11'67	11 0	181'85
18	14'53	11 0	142'40	11'67	11 0	181'85
19	15'50	4 45	..	63'53	..	11'00	4 45	..	83'67	..
20	16'33	12 0	142'69	11'33	12 0	..	205'29	..
21	15'50	12 0	148'63	12'00	12 0	195'40
22	15'50	12 0	..	160'40	..	11'50	12 0	201'31
23	14'17	11 0	150'73	12'50	5 0	77'15
24	15'33	11 0	137'75	12'00	2 0	82'15
25	15'00	11 30	..	158'93	..	11'60	12 0	199'59
26	14'67	12 0	157'04	11'50	12 0	..	202'18	..
27	14'60	12 0	157'79	12'20	12 0	189'79
Total -	(15'69)	217 15	580'05	840'38	1323'70	(12'00)	239 45	595'49	1255'92	2016'95
Aug. 1	14'67	12 0	157'04	11'40	12 0	..	203'96	..
2	14'67	11 0	143'95	11'25	11 0	188'6
3	14'67	11 0	143'95	11'60	11 0	182'9
4	14'67	12 0	157'04	11'60	12 0	..	200'44	..
5	15'33	12 0	151'99	11'60	12 0	199'5
6	15'00	12 0	153'58	11'60	12 0	202'14
7	15'17	12 0	161'86	11'60	12 0	..	200'44	..
8	15'33	11 0	..	148'75	..	11'80	11 0	179'8
9	11'00	5 0	87'6
10	12'00	12 0	..	193'76	..
11	11'00	11 30	191'2
12	15'33	12 0	150'28
13	15'33	12 0	..	162'27
14	15'00	7 30	95'99
15	12'40	11 0	171'1
16	11'50	12 0	..	202'18	..
17	15'00	12 0	153'58
18	15'00	12 0	..	165'84
19	15'67	11 0	134'76	11'50	11 0	184'5
20	14'83	12 0	157'12	12'00	12 0	195'40
Total -	(15'04)	171 30	309'11	475'86	1442'03	(11'65)	167 30	397'54	1000'78	1385'63

Table I.—continued.

Detailed Record of the Sewage applied to Permanent Grass Land.
Second Season 1861-2.

DATE.	Five-acre Field.					Ten-acre Field.				
	Average time taken to fill Gauge-tank (3' 216 tons).	Time of application.	Sewage applied (calculated per acre).			Average time taken to fill Gauge-tank (3' 216 tons).	Time of application.	Sewage applied (calculated per acre).		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area 1'00812 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99688 acre.)	Plot 4. (Area 1'00019 acre.)
Sept. 1	Mins. 14'53	H. M. 11 0	Tons. ..	Tons. ..	Tons. 142'40	Mins. 11'50	H. M. 11 0	Tons. ..	Tons. ..	Tons. 184'53
2	14'53	12 0	..	167'74	..	11'33	12 0	..	205'22	..
3	15'50	11 0	136'24
9	15'33	12 0	..	162'27
15	15'07	11 0	134'76	11'67	11 0	181'85
16	14'67	6 0	79'42	12'17	12 0	..	191'05	..
21	15'00	6 0	..	82'92
22	17'00	3 0	33'88	12'50	3 0	46'30
23	16'33	6 0	..	76'17	..	13'20	12 0	177'64
24	18'33	6 0	62'84
25	17'33	11 0	118'44	12'20	11 0	173'95
26	18'50	12 0	..	134'47	..	12'17	12 0	190'23
Total .	(15'09)	107 0	79'42	623'57	628'56	(12'03)	84 0	177'64	396'27	776'86
Oct. 6	24'00	11 0	85'84	23'40	11 0	90'60
7	23'07	12 0	..	86'77	..	26'20	12 0	..	88'74	..
13	16'00	11 0	131'98	11'50	11 0	184'54
14	16'00	6 0	72'82	11'16	12 0	..	208'34	..
15	16'33	6 0	..	76'17
20	12'33	11 0	172'11
21	12'67	11 0	..	168'22	..
22	12'50	11 0	171'95
23	15'33	11 0	137'75
24	15'25	8 0	..	108'75
25	16'00	11 0	131'98
27	12'33	5 15	83'20
28	12'00	1 15	..	20'18	..
29	12'33	4 30	70'41
30	16'00	11 0	131'98
31	16'23	9 0	..	114'81
32	16'50	3 0	34'90
33	16'00	12 0	145'03
34	16'00	5 30	60'75
35	16'33	6 30	76'42
Total .	(17'23)	123 0	285'20	386'50	730'85	(13'06)	90 0	255'15	485'48	517'75

TABLE II.
DETAILED RECORD of the SEWAGE applied to PERMANENT GRASS LAND.
THIRD SEASON 1862-3.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average time taken to fill Gauge-tank (3'216 tons).	Time of application.	Sewage applied (calculated per acre).			Average time taken to fill Gauge-tank (3'216 tons).	Time of application.	Sewage applied (calculated per acre).		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area 1'00512 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99588 acre.)	Plot 4. (Area 1'00019 acre.)
Nov. 1	Mins. 17'00	H. M. 11 0	Tons.	Tons.	Tons.	Mins. 12'22	H. M.	Tons.	Tons.	Tons.
3	12'67	4 0	60'91
4	12'50	10 0	..	155'01	..
5	12'83	10 0	150'37
6	16'67	10 0	115'16
7	16'33	10 0	..	126'95
8	15'83	10 0	121'27
10	13'50	10 0	142'91
11	12'67	10 0	..	152'93	..
12	13'00	10 0	150'31
13	16'83	10 0	114'07
14	16'00	10 0	..	129'56
15	16'25	7 30	89'62
17	12'60	9 30	145'46
18	13'00	10 0	..	149'05	..
19	12'50	10 0	154'34
20	17'00	10 0	112'93
21	16'33	10 0	..	126'95
22	15'40	9 0	112'19
24	12'33	7 30	117'35
25	12'40	9 0	..	140'63	..
26	12'40	8 45	137'88
27	16'60	9 15	106'98
28	16'00	10 0	..	129'56
29	16'00	9 15	112'26
Total -	(16'33)	120 0	201'88	513'02	806'82	(12'71)	108 45	288'19	597'62	771'34
Dec. 1	12'50	10 0	154'34
2	12'50	10 0	..	155'01	..
3	12'40	9 0	140'03
4	16'67	10 0	115'16
5	16'50	10 0	..	125'64
6	16'67	10 0	115'16
9	12'40	8 0	124'47
10	12'40	9 0	..	140'63	..
11	18'00	6 0	63'99
12	17'00	6 45	..	82'31
13	17'00	10 0	114'22
15	12'00	10 0	160'77
16	11'67	10 0	..	166'03	..
17	11'60	9 0	161'61
18	17'00	10 0	112'93
19	19'33	10 0	..	107'24
20	18'60	9 0	92'89
22	11'67	10 0	165'32
23	11'83	10 0	..	163'79	..
24	11'60	9 0	149'68
Total -	(17'34)	81 45	114'22	315'19	500'13	(12'04)	104 0	161'61	625'46	894'61
Jan. 5	11'67	10 0	165'32
6	12'00	10 0	162'84
7	11'40	9 0	152'31
8	16'33	10 0	112'04
9	16'33	10 0	117'56
10	16'67	10 0	116'48
12	11'83	10 0	163'08
13	11'67	10 0	..	166'03	..
14	11'67	10 0	165'32
15	17'83	10 0	107'67
16	16'80	9 0	..	111'06
17	16'17	10 0	118'72
19	12'00	8 30	40'19
20	12'00	10 0	..	161'47	..
21	11'83	10 0	165'18
22	16'33	10 0	117'56
23	16'00	10 0	..	129'56
24	15'83	10 0	122'66
26	11'83	10 0	163'08
27	12'50	7 0	..	108'50	..
28	11'80	10 0	165'60
29	16'67	10 0	..	124'86
30	16'17	10 0	118'72
31	16'67	10 0	..	124'86
Total	(16'55)	119 0	351'18	459'34	580'25	(11'82)	108 30	403'62	430'00	840'38

36. Italian rye grass, in the condition as cut, was also found to be more succulent and to contain less solid matter when grown with sewage than without it; but the proportion of dry substance diminished less as the season advanced in its case than in that of the meadow grass.

37. The proportion of nitrogenous substance (and also of impure waxy or fatty matter) was much greater in the solid matter of the sewaged, than in that of the unsewaged grass. The proportion of nitrogenous substance was also much higher in the solid matter of the grass grown towards the end than earlier in the season. The proportion of indigestible woody fibre was much about the same in the dry substance of the unsewaged and of the sewaged grass. It progressively diminished as the season advanced, and was generally lower in the dry substance of the Italian rye than in that of the meadow grass.

38. A given amount of the dry substance of grass grown in a cold and wet season, or during the cold and wet periods of the year, generally contains more nitrogenous substance, but is less productive than that of grass grown in more genial weather.

39. The greater productiveness in milk and increase of a given amount of the solid matter of the sewaged grass appears to depend more on a favourable condition of maturation, digestibility, and assimilability, of the constituents, than on the actual percentage amount of any of those determined, and above enumerated.

Effects of Sewage on the mixed Herbage of Grass Land.

40. The effect of sewage irrigation on the mixed herbage of grass land is to develop the Graminaceous plants chiefly, nearly to exclude the Leguminous, and to reduce the prevalence of miscellaneous or weedy plants, but much to encourage individual species.

41. Among the grasses which have been observed to be the most encouraged by sewage are (according to locality or other circumstances) rough meadow grass, couch grass, rough cock's foot, woolly soft grass, and perennial rye grass; two or three only remaining in any considerable proportion after sewage has been liberally applied for some years.

42. The produce of sewage irrigated meadows being generally cut or grazed very young, the tendency which the great luxuriance of a few very free growing grasses has to give a coarse and stemmy later growth is not an objection as in the case of meadows left for hay; a given weight of the dry or solid substance of the more simple sewaged grass being, when consumed green, more productive than an equal weight of that of the more complex unsewaged herbage.

Composition of the Milk from the unsewaged and the sewaged Grass.

43. Although more milk was obtained from a given weight of the dry or solid substance of sewaged than of unsewaged grass, there was comparatively little difference in the composition or

Table II.—continued.
Detailed Record of the Sewage applied to Permanent Grass Land.
Third Season 1862-3.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average time taken to fill Gauge-tank (3'216 tons).	Time of applica-tion.	Sewage applied (calculated per acre).			Average time taken to fill Gauge-tank (3'216 tons).	Time of applica-tion.	Sewage applied (calculated per acre).		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area 1'00512 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99588 acre.)	Plot 4. (Area 1'00019 acre.)
May 1	Mins.	H. M.	Tons.	Tons.	Tons.	Mins.	H. M.	Tons.	Tons.	Tons.
4	25'80	10 30	69'09	11'83	12 0	195'70
5	28'83	11 15	..	80'89
6	27'67	12 0	83'26
7	11'67	12 0	198'38
8	11'67	12 0	..	199'24	..
12	35'33	11 15	61'13
13	24'17	11 15	90'38
14	11'80	9 30	155'32
15	13'50	2 45	39'81
18	16'67	5 0	..	62'18
19	18'50	12 0	139'62
20	18'33	12 0	..	152'31
27	16'33	5 0	58'78
28	16'67	12 0	..	149'23	..	11'33	12 0	..	205'22	..
29	16'17	12 0	142'47	12'00	12 0	192'92
30	16'17	10 30	..	134'61	..	11'40	10 30	..	178'46	..
Total -	(20'28)	124 45	90'38	579'25	555'25	(11'72)	82 45	39'81	582'92	742'32
June 1	16 00	11 0	131'98	12'00	11 0	..	177'61	..
2	16'50	12 0	..	150'77	..	11'83	12 0	195'70
3	20'00	12 0	115'19	11'67	12 0	..	199'24	..
4	21'17	12 0	110'07	11'67	12 0	198'38
5	23'00	12 0	100'16	11'50	12 0	..	202'18	..
6	17'20	10 30	118'54	12'17	10 30	166'45
8	16'25	9 45	115'19
9	17'80	12 0	..	139'76
10	15'50	12 0	150'33
11	11'83	12 0	..	196'54	..
12	12'00	12 0	192'92
15	17'00	9 45	111'36
16	16'83	12 0	..	147'81
17	17'00	12 0	137'06
18	11'67	12 0	..	199'24	..
19	11'83	11 30	189'95
22	16'00	11 0	133'49
23	15'67	7 45	..	102'53
24	17'17	12 0	135'71
25	12'00	12 0	195'40
26	11'83	12 0	195'70
Total -	(17'41)	167 45	896'56	540'87	462'52	(11'83)	141 0	385'35	974'81	949'15
July 1	19'17	12 0	120'17
2	12'00	12 0	192'92
3	12'20	11 0	176'18
6	17'67	10 30	114'08
7	12'33	11 30	182'25
8	17'17	11 30	128'58
9	11'75	8 45	143'67
10	16'40	9 45	114'13
13	17'67	10 30	114'08
14	17'33	11 30	..	137'56
15	18'00	9 0	95'99
16	11'25	8 0	137'13
17	12'25	8 0	125'92
20	16'17	11 0	..	141'02
21	17'80	9 45	100'90
22	16'83	12 0	..	147'81
23	12'33	12 0	..	188'57	..
24	11'80	11 0	179'84
27	17'50	10 15	112'44
28	17'80	10 30	113'25
29	17'67	6 15	67'90
30	12'40	10 0	..	156'26	..
31	11'67	8 0	133'95
Total -	(17'44)	134 0	100'90	426'39	980'62	(12'02)	100 15	492'38	944'83	779'63

Table II.—continued.
Detailed Record of the Sewage applied to Permanent Grass Land.
Third Season 1862-3.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average time taken to fill Gauge-tank (3'216 tons).	Time of applica-tion.	Sewage applied (calculated per acre).			Average time taken to fill Gauge-tank (3'216 tons).	Time of applica-tion.	Sewage applied (calculated per acre).		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area 1'00512 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99588 acre.)	Plot 4. (Area 1'00019 acre.)
Aug. 3	Mins.	H. M.	Tons.	Tons.	Tons.	Mins.	H. M.	Tons.	Tons.	Tons.
4	16'00	2 30	30'00
5	16'00	12 0	143'98
6	17'17	12 0	..	144'88
7	13'00	12 0	178'08
10	16'00	10 0	119'99	14'00	10 45	150'04
11	17'33	12 0	..	143'55
12	19'67	7 0	68'32
17	16'33	11 0	129'32
18	18'40	10 45	..	121'12
19	23'00	1 30	12'52
20	12'33	11 0	172'11
21	12'40	11 0	171'14
24	15'40	10 0	121'66
25	15'50	8 0	99'09
26	15'50	12 0	124'53
27	11'83	12 0	195'70
31	17'00	10 15	111'81
Total -	(17'00)	119 0	..	409'55	964'22	(12'65)	56 45	150'04	..	717'03
Sept. 1	15'83	12 0	..	157'15
2	18'33	12 0	125'68
3	13'40	10 15	147'57
4	12'50	12 0	..	186'01	..
7	17'50	11 0	120'67
8	16'50	11 30	..	144'48
9	17'00	12 0	135'51
10	12'00	9 0	..	145'32	..
11	13'50	7 0	100'03
15	16'00	12 0	145'63
16	16'75	8 15	..	102'11
17	13'00	7 30	111'30
18	12'80	9 15	..	140'02	..
21	19'00	11 0	111'15
22	17'40	10 30	..	125'10
23	18'75	8 0	..	88'45
24	12'75	9 0	..	169'54	136'18
25	12'00	10 30
28	17'83	11 0	118'44
29	18'00	12 0	129'45
30	19'67	5 0	48'80
Total -	(17'39)	136 15	275'08	617'29	660'25	(12'68)	74 30	..	640'89	495'08
Oct. 1	12'33	12 0	187'76
2	13'50	8 0	115'79
5	16'50	6 30	..	81'67
6	16'50	4 30	52'36
7	18'00	9 30	101'32
7	19'25	8 45	87'26
8	11'80	10 30	171'67
9	12'33	12 0	..	188'57	..
12	17'67	11 0	119'51
13	16'83	12 0	136'88
14	17'67	12 0	..	140'78
15	13'00	12 0	178'08
16	12'00	12 0	..	193'76	..
17	12'50	11 0	171'95
19	17'00	5 0	57'11
20	18'00	6 0	63'99
21	17'17	12 0	134'17
22	17'33	6 0	66'47	12'00	2 45	44'21
23	14'20	8 45	118'83
24	13'67	6 0	85'77
25	15'00	6 0	..	77'50	..
26	20'00	1 0	..	10'37	..	11'00	1 0	17'76
27	16'00	4 45	56'99	14'00	7 0	96'46
Total -	(17'42)	99 0	57'11	252'82	818'95	(12'61)	109 0	391'27	459'83	797'03

TABLE III.
DETAILED RECORD of the Amounts of GREEN PRODUCE obtained in t
ments on PERMANENT GRASS LAND.
SECOND SEASON 1862.

Dates of Cuttings.	Green Grass obtained (calculated per acre).											
	Five-acre Field.								Ten-acre Field.			
	Without Sewage.		With Sewage.						Without Sewage.		With Sew.	
			Plot 1.*		Plot 2.		Plot 3.					
	tons.	cwt.	qrs.	lbs.	tons.	cwt.	qrs.	lbs.	tons.	cwt.	qrs.	lbs.
		</										

* Area of Plot 1, five-acre field = 0.80756 acre, and of Plot 1, ten-acre field = 0.79243 acre; for sewage plots, see preceding Tables I. and II. of sewage applied.

Table III.—continued.

Detailed Record of the Amounts of Green Produce obtained in the Experiments on Permanent Grass Land.

Second Season 1862.

Dates of sowing.	Green Grass obtained (calculated per acre).											
	Five-acre Field.								Ten-acre Field.			
	Without Sewage.		With Sewage.						Without Sewage.		With Sewage.	
			Plot 1.	Plot 2.	Plot 3.	Plot 4.						
	tons.	cwt.	grs.	lbs.	tons.	cwt.	grs.	lbs.	tons.	cwt.	grs.	lbs.
1	0 2 1 10	0 9 0 23	0 3 5 6	0 12 3 3
2	0 2 2 9	0 8 0 3	0 3 1 24	0 18 0 0
3	0 2 0 26	0 8 1 19	0 3 0 17	0 15 2 0
4	..	0 3 2 8	0 3 1 24	0 7 1 20
5	0 5 1 16	0 15 2 8	0 8 3 11	1 10 3 25
6	0 2 2 17	0 9 0 5	0 4 0 25	0 16 3 11
7	0 2 2 9	0 4 2 7	0 0 3 8	0 6 1 16	0 16 3 22
8	0 2 3 11	0 7 1 13	0 5 2 6	0 18 2 10
9	0 2 1 0	0 7 2 11	0 16 0 0
10	0 2 3 19	0 7 1 11	0 3 2 17	0 18 1 12
11	0 2 2 19	0 7 1 11	0 6 3 27	..	1 7 1 19	0 7 0 11
12	0 5 3 18	0 17 2 23
13
14	0 2 3 5	0 8 1 18	0 3 1 19	..	0 12 2 11	10 11 2 9
15	0 2 1 12	0 8 1 14	10 19 1 27	..	0 14 2 26	Total
16	0 2 1 10	0 7 1 25	0 18 3 1	2d Crop.
17	0 2 1 8	0 7 0 13	Total	..	0 14 1 26	..
18	0 2 2 8	0 8 3 0	1st Crop.	..	0 16 2 7	..
19	0 4 2 15	0 12 0 23	0 3 3 21	1 5 1 0	..
20
21	0 2 1 0	0 6 0 13	6 9 3 6	..
22	0 2 1 13	0 5 3 22
23
24	0 2 0 26	14 12 1 4	0 8 1 5	Total	..
25	0 2 2 13	0 8 2 4	2d Crop.	..
26	0 2 1 3	0 10 0 6	0 4 3 3
27	0 4 2 16	0 15 1 14
28	0 2 1 17	Total	0 9 0 17
29	0 1 2 22	1st Crop.	0 8 3 24
30	0 1 2 17	1 1 0 17
31	0 5 0 6	0 17 1 8
32	8 16 0 23	8 2 3 3	4 19 1 19	2 11 2 0	0 3 2 24	6 9 3 6	8 16 1 2
1	1st Crop. (cont.)	2d Crop.	2d Crop.	2d Crop. (cont.)	2d Crop.	2d Crop. (cont.)	2d Crop.
2	0 4 3 20	..	0 9 3 0	0 18 2 1
3	0 10 0 13	1 2 0 23
4	0 1 2 20	..	0 9 0 5	7 0 0 15
5	0 1 2 19	..	0 9 1 8
6	0 2 1 0	..	0 11 0 14	Total
7	0 2 0 1	..	0 12 3 8	2d Crop.
8	0 2 0 21	..	0 9 3 20
9	0 4 1 13	..	0 19 3 9
10	0 1 3 27	..	0 9 2 9
11	0 1 2 9	..	0 9 3 3
12	0 1 2 20	..	0 9 3 15
13	0 2 3 11	..	0 12 1 24
14	0 1 3 16	..	0 9 1 1
15	0 2 3 26	..	0 18 1 4
16	0 2 1 12	..	0 12 1 11
17	0 1 3 16	..	0 10 1 1
18	0 1 3 19	..	0 10 0 19
19
20
21	6 13 1 3	..	0 8 2 14
22
23	Total	0 10 1 4	9 13 0 25
24	1st Crop.	0 19 0 27
25	..	0 8 1 25	Total
26	..	0 9 2 20	2d Crop.
27	..	0 10 1 9	0 10 3 13
28	..	0 9 2 14	0 13 1 21
29	..	0 9 0 13	0 15 0 7
30	..	0 19 3 16	0 11 1 22
31
32	8 8 2 10	4 16 2 16	9 12 0 25	2 0 2 24	0 11 3 0	6 12 1 3	2 15 3 8

Table I.—continued.
Detailed Record of the Sewage applied to Permanent Grass Land
Second Season 1861-2.

DATES.	Five-acre Field.					Ten-acre Field.				
	Average Time taken to fill Gauge-tank (3'216 tons).	Time of application.	Sewage applied (calculated per acre).			Average Time taken to fill Gauge-tank (3'216 tons).	Time of application.	Sewage applied (calculated per acre).		
			Plot 2. (Area '99375 acre.)	Plot 3. (Area '93081 acre.)	Plot 4. (Area 1'00512 acre.)			Plot 2. (Area '9875 acre.)	Plot 3. (Area '99588 acre.)	Plot 4. (Area 1'00512 acre.)
Jan. 1	Mins.	H. M.	Tons.	Tons.	Tons.	Mins.	H. M.	Tons.	Tons.	Tons.
6	15'00	3 30	45'31
7	15'16	9 45	123'47
8	15'16	9 45	..	133'32	..	12'00	4 45	77'35
9	12'00	5 0	8
10	15'00	5 45	74'43
11	15'34	4 0	50'06
12	14'50	9 45	129'09
13	14'07	9 45	..	137'78	..	12'00	5 0	..	80'73	..
14	12'34	5 0	7
15	14'34	6 0	81'24
16	15'34	4 0	50'06
17	21'25	5 45	..	56'09
18	16'34	5 0	59'42
19	16'00	5 0	59'99
20	15'83	9 45	118'24
21	15'50	9 45	..	130'40	..	12'34	5 0	..	78'51	..
22	12'50	5 0	7
23	15'50	5 0	62'64
24	14'07	4 0	52'34
Total -	(15'41)	103 30	323'04	457'59	583'25	(12'20)	29 45	77'35	159'24	230
Feb. 3	22'16	11 15	97'46
4	15'07	11 30	..	152'14	..	11'07	7 45	129'77
5	12'50	3 45	5
6	16'34	11 15	132'17
7	16'00	10 45	124'32
8	16'16	11 30	..	147'52	..	11'50	5 45	..	96'88	..
9	12'34	5 45	8
10	18'00	11 30	122'65
11	16'34	4 45	55'81
12	16'83	11 30	..	141'65	..	12'34	5 45	..	90'28	..
13	12'00	5 45	9
14	16'07	11 45	136'86
15	16'34	11 15	132'17
16	17'83	11 30	..	133'71	..	11'50	5 45	97'70
17	12'34	5 45	8
18	18'00	3 0	32'36
19	16'00	7 15	80'99
Total -	(17'03)	128 45	169'22	575'02	751'57	(11'98)	46 0	227'47	187'16	330
Mar. 3	16'00	11 15	134'98
4	18'00	11 30	..	132'44	..	12'00	5 45	..	92'84	..
5	11'34	5 45	9
6	16'34	6 45	80'21
7	17'34	3 45	41'52
8	(8 tanks*)	11 0	25'60
9	16'00	11 30	..	149'00	..	12'00	5 45	..	92'84	..
10	11'33	5 45	9
11	14'50	4 15	56'91
12	13'67	5 45	80'75
13	12'00	4 45	7
14	11'83	11 30	..	188'35	..
15	12'00	6 45	109'91
16	11'67	4 15	7
17	14'07	11 0	143'95
18	13'67	11 30	..	174'40
19	13'35	6 45	98'32
20	14'00	3 45	51'42
21	12'16	11'30	18
Total -	(16'73)	98 45	235'44	455'84	478'22	(11'85)	61 45	109'91	374'03	524

* When the number of tanks is given, the flow was too slow and irregular to estimate it by average time taken to fill the gauge-tank, and therefore the actual number of tankfuls counted.

**ED RECORD of the AMOUNTS of GREEN PRODUCE obtained in the EXPERI-
MENTS ON PERMANENT GRASS LAND.**

Green Grass obtained (calculated per acre).

[illegible]

**Detailed Record of the Amounts of Green Produce obtained in the Experiment
on Permanent Grass Land.**

Green Grass obtained (calculated per acre).

Dates of Outtings.		Green Grass obtained (calculated per acre).																											
		Five-acre Field.												Ten-acre Field.															
		Without Sewage.				With Sewage.				Without Sewage.				With Sewage.															
		Plot 1.			Plot 2.	Plot 3.			Plot 4.	Plot 1.			Plot 2.	Plot 3.			Plot 4.												
		tons.	cwt.	grs.	lbs.	tons.	cwt.	grs.	lbs.	tons.	cwt.	grs.	lbs.	tons.	cwt.	grs.	lbs.	tons.	cwt.	grs.	lbs.								
		1st Crop.				2d Crop.				2d Crop.				1st Crop.				1st Crop.				1st Crop.				2d Crop.			
		(cont.)																(cont.)											
June	1	0 11 3 23								
	2	0 10 0 14								
	3	1 10 1 21								
	4								
	5								
	6								
	8								
	9								
	10								
	11								
	12								
	13								
	15	0 13 2 18								
	16	1 8 3 5								
	17	1 9 3 7								
	18	[10 14 1 0]								
	19	Total								
	20	1st Crop.	2 5 3 18								
	22	1 11 1 11								
	23								
	24	0 16 2 7								
	25	0 14 3 10								
	26	4 6 3 0								
	27								
	28	2 9 1 27								
	29	[10 13 2 0]								
	30	0 18 3 6								
		0 19 1 23								
		Total								
		2d Crop.								
Monthly Total		6 4 2 3	3 9 2 20								
		10 18 2 0								
		3 7 3 25								
		13 1 2 23								
		5 16 0 23								
		11 13 2								
July	1	1 0 3 0								
	2								
	3								
	4								
	6	0 16 1 0								
	7	0 17 3 20								
	8	1 0 0 9								
	9	2 6 0 6								
	10								
	11	0 4 0 23	1 7 1 12								
	13	0 3 0 23	0 15 0 17								
	14	0 4 0 18	0 10 0 19	[9 7 0 22]								
	15	0 3 0 1	0 13 1 11								
	16	..	0 16 1 11								
	17	..	1 0 0 0	Total								
	18	0 9 1 4	1 8 0 27	2d Crop.								
	20	0 3 2 17	1 4 2 23								
	22								
	23								
	24								
	25								
	27	0 5 1 11	0 15 0 2								
	28	[3 18 3 14]	[6 7 3 9]								
	29	Total	Total								
	30	1st Crop.	2d Crop.								
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
									
																					

Table IV.—*continued.*Record of the Amounts of Green Produce obtained in the Experiments
on Permanent Grass Land.

Third Season 1863.

Green Grass obtained (calculated per acre).											
Five-acre Field.						Ten-acre Field.					
Without Sewage.		With Sewage.				Without Sewage.		With Sewage.			
Plot 1.	Plot 2.	Plot 3.	Plot 4.		Plot 1.	Plot 2.	Plot 3.	Plot 4.			
tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.	tons. cwt. grs. lbs.		
..	..	3d Crop.	3d Crop.		3d Crop.	3d Crop.			
..	1 11 1 3				
..	0 18 0 18				
..	0 14 3 26				
..	0 19 2 3				
..	0 19 0 3				
..	..	1 12 3 10	0 15 2 0			
..	..	1 3 3 15			
..	0 15 0 0			
..	0 14 0 0			
..	0 16 3 5				
..	0 19 0 3				
..	..	1 6 1 8	[6 18 2 5]		0 15 1 0			
..	Total		0 15 2 0			
..	3d Crop.		0 16 1 0			
..	0 18 1 14			
..	0 18 3 0			
..	1 12 0 0			
..	0 14 3 7	[8 6 1 26]			
..	0 13 3 6	..			
..	0 15 3 7	Total			
..	1 0 1 9	3d Crop.			
..	0 15 3 7	..			
..	1 15 0 20	..			
..	..	4 2 3 5	7 15 3 16		5 14 3 0	7 18 2 14			
2d Crop.	3d Crop.	3d Crop.	4th Crop.	2d Crop.	3d Crop.	3d Crop.	4th Crop.	2d Crop.	3d Crop.		
..	0 18 2 3		
..	..	0 15 1 7	0 12 3 0		
..	..	1 8 1 17		
..	1 5 1 12		
..	..	0 16 1 0	[7 0 0 12]		
..		
..	..	[7 2 3 1]	1 8 0 11	Total		
..	..	Total	1 1 3 8	3d Crop.		
..	..	3d Crop.	0 18 0 11		
..	1 0 3 15	1 1 0 1		
..	0 14 0 10		
..	0 12 2 2	..	0 14 1 27		
..		
..		
..	0 14 2 13	..	0 11 3 9		
..	0 17 0 11		
..	1 1 1 23		
..	0 17 3 18		
0 5 2 8		
0 3 3 24		
0 10 3 23		
0 19 3 27	[8 11 1 13]	0 17 3 13		
Total	1 1 0 14	1 1 3 16		
2d Crop.	..	4th Crop.	Total	..	1 12 3 23	[6 1 0 2]		
..	..	1 2 0 25	4th Crop.	..	Total	Total		
..	..	1 11 0 17	2d Crop.	3d Crop.		
..	..	0 18 3 13		
10 3 27	2 7 1 27	6 12 0 23	7 14 0 2	1 12 3 23	6 1 0 2	1 5 1 12	0 12 3 0				

Table IV.—*continued.*

Detailed Record of the Amounts of Green Produce obtained in the Experiment on Permanent Grass Land.

Third Season 1863.

Green Grass obtained (calculated per acre).																	
Dates of Cuttings.	Five-acre Field.								Ten-acre Field.								
	Without Sewage.			With Sewage.					Without Sewage.			With Sewage.					
Plot 1.			Plot 2.		Plot 3.		Plot 4.		Plot 1.			Plot 2.		Plot 3.		P4	
tons. cwtls. qrs. lbs.			tons. cwtls. qrs. lbs.		tons. cwtls. qrs. lbs.		tons. cwtls. qrs. lbs.		tons. cwtls. qrs. lbs.			tons. cwtls. qrs. lbs.		tons. cwtls. qrs. lbs.		tons. cwtls. qrs. lbs.	
Oct. 1	..			3d Crop. (cont.)		4th Crop. (cont.)		5th Crop.			4th Crop.		4th (cont.)
2		1 10
3		0 19
5		1 12 1 5			1 10
6		0 17
7		0 9 0 20		0 14
8		1 4 0 25		5 5
9		1 16 2 3		T
10	..			0 15 3 11			3 9 3 20		4th
13	..			0 19 2 14			Total 4th Crop.		
14	..			4 2 3 24		0 13 2 0			
19	..			Total 3d Crop.		5 18 0 4		0 14 1 20			
20	..			4th Crop. 0 9 0 13		Total 4th Crop. ..		Total 5th Crop.		
21	..			0 7 3 5			
				0 16 3 18		Total 4th Crop.											
Monthly Total }	..			2 12 1 15		2 5 3 5		0 14 1 20			3 9 3 20		5 12
Nov. 27	..			5th Crop. 0 3 0 16		5th Crop. 0 5 2 10		6th Crop. 0 5 2 24		..			4th Crop. ..		5th Crop. ..		5th
28	..			Total 5th Crop.		Total 5th Crop.		Total 6th Crop.		..			0 9 1 6		0 10 0 5		0 11
													Total 4th Crop.		Total 5th Crop.		T 5th
Monthly Total }	..			0 3 0 16		0 5 2 10		0 5 2 24		..			0 9 1 6		0 10 0 5		0 11

TABLE V.
DETAILED RECORD of the SEWAGE applied, and of the AMOUNTS of
GREEN PRODUCE obtained in EXPERIMENTS on ITALIAN RYE-GRASS.
SEASON 1863.

DATE.	Average time taken to fill Gauge-tank (3'308 tons).	Time of applica- tion.	Sewage applied (calculated per acre).		Green Grass obtained (calculated per acre).					
			Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)	Without Sewage.	With Sewage.			Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)
					Plot 1. (Area 1'00484 acre.)	Plot 1.	Plot 2.	Plot 3.		
						tons. cwt. lbs.	tons. cwt. lbs.	tons. cwt. lbs.		
April 24	1st Crop.	0 17 0 2	1 10 2 6	1 10 2 6		
25	1 12 0 17	0 16 0 3	0 16 0 3		
27	0 14 1 1	1 9 1 6	1 9 1 6		
28	1 3 2 16	1 0 0 2	[3 15 3 15]			
29	91'67	12 0	..	20'03	1 5 2 31	[4 3 1 22]	Total	1st Crop.		
30	101'67	11 15	..	22'00	0 15 0 12	Total	1st Crop.			
Monthly Total	(96'24)	23 15	..	48'03	3 4 1 21	4 3 1 22	3 15 3 15			
May 1	120'00	12 0	19'87	..	0 11 0 22		
2	107'50	11 30	..	21'27	0 12 0 21		
3	137'50	5 0	..	7'23	0 14 1 8		
4		
5	123'33	12 0	..	17'89	[5 2 0 16]		
6	120'00	11 30	19'04		
7	117'50	11 30	..	19'46	Total		
8	122'50	11 15	..	18'20	1st Crop.		
9	120'00	11 0	18'21		
11	123'33	12 0	..	19'35		
12	127'50	10 0	..	14'46		
13	88'75	11 30	25'75		
14	112'50	12 0	..	21'21		
15	75'00	4 30	..	11'93		
16	72'50	12 0	32'89		
17	69'00	12 0	..	34'58		
18	67'50	4 30	..	13'25		
19	65'00	12 0	36'34		
20	71'25	10 45	..	30'00		
21	75'60	10 30	..	28'76		
Monthly Total	(95'80)	197 30	152'10	257'05	1 17 2 23	..	1 11 1 7			
June 1	88'75	10 15	22'93		
2	78'75	12 0	..	30'30		
3	101'67	8 45	..	17'11		
4	85'00	10 30	21'96		
5	90'00	12 0	..	26'51		
6	90'00	11 0	..	24'30		
7	97'50	9 15	..	18'86		
8	130'00	11 0	..	16'82		
9	85'00	10 15	..	23'97		
10	76'00	12 0	..	31'39		
11	87'50	12 0	..	27'27		
12	73'75	11 0	..	29'65		
13	110'67	8 30	14'48		
14	102'50	11 15	..	21'83		
15	96'25	8 45	..	18'07		
16	93'75	12 0	25'43		
17	86'25	12 0	..	27'68		
18	85'00	9 0	..	21'05		
19	86'25	8 45	20'16		
20	83'33	7 0	16'69		
21	80'00	12 0	..	19'88		
22	80'00	..	19'87		
23	5'75	..	19'04		
24	5'25	..	17'38		
25		
26		
27		
28		
29		
30		
Monthly Total	177'96	354'06	8 18 0 2	7 1 1 27	6 9 3 13			

When the number of tanks is given, the flow was too slow and irregular to estimate it by the time taken to fill the gauge-tank, and therefore the actual number of tankfuls was used.

10000

TABLE V.—continued.
Detailed Record of the Sewage applied, and of the Amounts of Green
Produce obtained in Experiments on Italian Rye-grass.

Season 1863.

DATES.	Average Time* taken to fill Gauge-tank (3'308 tons).	Time of applica-tion.	Sewage applied (calculated per acre).		Green Grass obtained (calculated per acre).					
			Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)	Without Sewage. Plot 1. (Area 1'00484 acre.)	With Sewage.				
						Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)			
	Mins.	H. M.	Tons.	Tons.	tons, cwt., qrs., lbs.	tons, cwt., qrs., lbs.	tons, cwt., qrs., lbs.	tons, cwt., qrs., lbs.		
July 1	(3'25tanks)	..	10'76	3d Crop.	0 6 1 1	
2	(5'75 ")	..	19'04	0 11 0 18		
3	(5'00 ")	16'57	0 7 0 1		
4	(3'00 ")	..	9'93	0 14 0 3		
6	(1'50 ")	4'97	0 9 1 2		
"	30'50	2 0	..	13'04		
7	(4'75tanks)	..	15'73	0 14 1 3		
"	31'50	2 0	12'62		
8	(2'75tanks)	9'11	0 5 3 1		
"	26'50	2 0	..	15'01		
9	(6'00tanks)	..	19'87	0 7 3 1		
10	(4'73 ")	15'67	0 9 0 12		
11	(6'30 ")	20'88	0 15 2 10		
13	(2'52 ")	8'35	3d Crop.	7 7 1 13		
"	24'50	2 0	..	16'23	..	0 10 0 11	..	Total 3d Crop.		
14	(3'00tanks)	9'94		
"	23'50	2 0	..	16'92	..	0 8 2 1		
15	(5'18tanks)	17'16	..	0 7 0 13		
"	24'50	2 0	..	16'23		
16	(6'75tanks)	22'37	..	0 7 0 6		
"	24'00	2 0	..	16'57		
17	(4'93tanks)	16'34	..	0 10 0 27		
"	24'50	2 0	..	16'23		
18	(6'00tanks)	19'88	..	0 13 0 1		
20	(3'00 ")	9'94	..	0 6 3 9		
"	23'50	2 0	..	16'92		
21	(4'00tanks)	..	13'25	0 7 0 15		
"	25'50	2 0	15'58		
22	(3'75tanks)	..	12'42	0 10 2 1		
"	25'50	2 0	15'58		
23	(7'00tanks)	23'20	..	0 5 2 25		
"	29'50	2 0	..	13'48		
24	(6'83tanks)	..	22'62	0 7 0 17		
"	32'50	2 0	12'23		
25	(2'30tanks)	7'62	..	0 13 2 22		
"	28'00	2 0	..	14'20		
27	(3'55tanks)	11'76	..	0 7 0 7		
"	29'50	2 0	..	13'48		
28	(1'00tank)	..	3'31	0 2 3 6	4th Crop.	0 6 0 15		
"	32'50	2 0	12'23		
29	(6'93tanks)	3'08	..	5 16 3 21	..	0 7 2 1		
30	(6'93 ")	..	22'95	Total 3d Crop.	0 9 1 8		
31	(5'43 ")	17'99	0 7 2 1		
Monthly Total	218'12	403'14	5 10 3 21	6 10 2 21		

* When the number of tanks is given, the flow was too slow and irregular to estimate it by the average time taken to fill the tanks. The actual number of tankfuls was

TABLE V.—continued.
Detailed Record of the Sewage applied, and of the Amounts of Green Produce obtained in Experiments on Italian Rye-grass.
Season 1863.

DATES.	Average time * taken to fill Gauge-tank (3'308 tons).	Time of application.	Sewage applied (calculated per acre).		Green Grass obtained (calculated per acre).					
					Without Sewage.		With Sewage.			
			Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)	Plot 1. (Area 1'00484 acre.)	Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)			
		H. M.	Tons.	Tons.	tons. cwt. qrs. lbs.	tons. cwt. qrs. lbs.	tons. cwt. qrs. lbs.	tons. cwt. qrs. lbs.	tons. cwt. qrs. lbs.	tons. cwt. qrs. lbs.
August 1	(4'00tnks.)	..	13'25	..	3rd Crop.	0 8 2 2
2	(0'83 "	2'75	0 5 3 2	0 5 0 3
4	(4'00 "	..	13'25	..	0 5 0 4	0 9 0 4
5	(3'00 "	9'94	0 1 1 21	0 5 2 23
6	(5'65 "	..	18'71	..	0 4 2 11	0 6 2 23
7	(3'48 "	11'53	0 3 2 26	0 5 3 1
8	(4'41 "	14'61	0 7 1 24	4th Crop.	..	0 9 3 26
10	(2'88 "	..	9'54	..	0 3 3 0	0 5 3 15	..	0 1 2 0
11	(3'20 "	..	10'60	..	0 2 2 4	0 4 1 16	4 2 223
12	(1'00 "	3'31	2 0 1 13	0 2 0 12	Total 4th Crop.
13	(2'62 "	8'68
14	(2'60 "	..	8'61	10'77	Total 3d Crop.
15	(3'25 "
17	(2'30 "	..	7'62
18	(1'00 "	3'31
20	(5'64 "	18'69
21	(3'40 "	..	11'26
22	(3'30 "	10'93
24	(3'15 "	..	10'43	0 2 2 20
25	(1'55 "	5'14	..	0 5 2 21
26	(3'56 "	11'80	..	0 4 3 0
27	(5'46 "	18'09	..	0 6 3 21
28	(5'40 "	17'89	..	0 1 8 20
29	(4'81 "	15'94	..	1 14 1 13	5th Crop.
31	(5'30 "	..	17'55	Total 4th Crop.	0 8 1 12
Monthly Total }	120'82	163'38	2 0 1 13	1 14 1 13	3 0 2 10
Sept. 1	(5'11tnks.)	16'93	0 4 0 17
2	(4'45 "	14'75	0 4 3 1
3	(3'45 "	..	11'42	0 5 0 21
4	(5'34 "	17'70	0 5 3 11
5	(3'81 "	12'63	..	5th Crop.	0 9 1 7
7	(4'40 "	..	14'57	0 1 3 4	0 2 3 10
8	(2'86 "	9'48	4th Crop.	0 1 2 24	2 0 1 23
9	(2'90 "	9'61	0 2 1 19	..	Total 5th Crop.
10	(5'61 "	18'59	0 4 0 8
11	(2'31 "	7'65	0 3 0 8
12	(2'46 "	..	8'15	..	0 9 2 7
15	(4'40 "	14'58
16	(1'86 "	6'16	Total 4th Crop.
17	(2'73 "	..	9'04
18	(3'60 "	11'93
19	(1'50 "	..	4'97
21	(3'98 "	13'19
22	(3'33 "	11'03	..	0 4 2 14
23	(1'60 "	..	5'30	0 4 2 10
24	(3'45 "	11'43	..	0 0 2 10
25	(4'00 "	13'25	..	0 13 1 6
26	(2'00 "	..	6'62	Total 5th Crop.
28	(4'44 "	14'71
29	(3'00 "	9'94
30	(1'90 "	6'30
Monthly Total }	60'07	219'86	0 9 2 7	0 13 1 6	1 12 0 11

* When the number of tanks is given, the flow was too slow and irregular to estimate it by the average time taken to fill the gauge-tank, and therefore the actual number of tankfuls was counted.

TABLE V.—*continued.*

Detailed Record of the Sewage applied, and of the Amounts of Green Produce obtained in Experiments on Italian Rye-grass.

Season 1863.

DATES.	Average time * taken to fill Gauge-tank (3'308 tons).	Time of application.	Sewage applied (calculated per acre).		Green Grass obtained (calculated per acre).											
			Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)	Without Sewage.	With Sewage.										
					Plot 1. (Area 1'00484 acre.)	Plot 2. (Area 0'999 acre.)	Plot 3. (Area 0'99831 acre.)									
					tons.	cwt.	qrs.	lbs.	tons.	cwt.	qrs.	lbs.	tons.	cwt.	qrs.	lbs.
Oct.	1 (7'28tnks.)	H. M.	Tons.	Tons.												
	2 (2'50 ")	..	24'11			
	3 (3'80 ")	..	12'58			
	5 (4'32 ")	14'32			
	6 (3'55 ")	11'76			
	7 (1'87 ")	..	6'19			
	8 (5'37 ")	17'79			
	9 (7'16 ")	23'73			
	10 (4'60 ")	..	15'23			
	15				6th Crop.	0 17 2		
	16				0 11 0			
	17			6th Crop.	0 12 2 1	0 14 2		
	21			0 13 1 15	2 3 0 1			
	23				5th Crop.			1 5 3 16	Total 6th Crop			
									0 6 0 9							
									Total 5th Crop.			Total 6th Crop.				
Monthly Total }	58'11	75'88	0 6 0 9				5 3 16			2 3 0 1				

* When the number of tanks is given, the flow was too slow and irregular to estimate it by the average time taken to fill the gauge-tank, and therefore the actual number of tankfuls was counted.

TABLE VI.

DETAILED RECORD of FOOD consumed, and INCREASE yielded, by OXEN fed on UNSEWAGED and SEWAGED MEADOW GRASS, with OILCAKE in addition.

SEASON 1862.

28 days, from May 8 to June 5.

Dates of weigh- ing.	Two Oxen on Unsewaged Grass.						Eight Oxen on Sewaged Grass.							
	Particulars of the Food consumed.													
	Grass.			Quantities (weighed green).	Oil- cake.†	Grass.			Quantities (weighed green).	Oil- cake.†				
	From		Crop.			From		Crop.						
Field.	Plot.*	Field.		Plot.	Crop.									
1862.				tons.	qrs.	lbs.	lbs.				tons.	qrs.	lbs.	lbs.
May 8	Five-acre	00	1	0 1 2 11		6	Five-acre	4	1	0 8 0 15			24	
9		00	1	0 2 0 11		6		4	1	0 8 2 14			24	
10		4	1	0 2 0 17		6		4	1	0 9 2 9			24	
11		4	1	0 2 0 17		6		4	1	0 9 2 9			24	
12		4	1	0 2 0 14		6		4	1	0 11 1 11			24	
13		4	1	0 2 0 24		6		4	1	0 9 3 0			24	
14		4	1	0 2 1 6		6		4	1	0 9 3 22			24	
15		4	1	0 2 1 17		6		4	1	0 10 1 6			24	
16		00	1	0 2 0 14		6		4	1	0 11 2 1			24	
17		4	1	0 2 0 12		6		4	1	0 10 2 21			24	
18		4	1	0 2 0 11		6		4	1	0 10 2 21			24	
19		4	1	0 2 0 24		6		4	1	0 12 0 27			24	
20		4	1	0 2 1 11		6		4	1	0 10 1 8			24	
21		4	1	0 2 1 22		6		4	1	0 10 3 15			24	
22		00	1	0 2 2 6		6		3	1	0 9 3 6			24	
23		00	1	0 2 0 14		6		3	1	0 9 0 7			24	
24		00	1	0 2 0 11		6		3	1	0 10 3 10			24	
25		3	1	0 2 0 14		6		3	1	0 10 3 9			24	
26		3	1	0 2 0 27		6		3	1	0 11 1 15			24	
27		00	1	0 2 3 16		6		3	1	0 10 2 1			24	
28		00	1	0 2 3 8		6		3	1	0 9 2 26			24	
29		0	1	0 2 0 4		6		3	1	0 8 2 13			24	
30		0	1	0 2 1 0		6		3	1	0 9 0 23			24	
31		0	1	0 2 0 18		6		3	1	0 8 3 22			24	
une 1		0	1	0 2 0 20		6		3	1	0 8 3 22			24	
2		0	1	0 2 1 13		6		3	1	0 9 3 1			24	
3		0	1	0 2 2 16		6		3	1	0 8 0 14			24	
4		0	1	0 2 1 1		6		3	1	0 9 0 7			24	
Total in 28 days				3 3 0 15	168	Total in 28 days				13 18 1 19	672			
Average per head per day				0 1 0 14	3	Average per head per day				0 1 0 27	3			

WEIGHTS AND INCREASE OF THE OXEN, &c.

Nos.	Weights.†		In-crease in 28 days.	Per 1,000 lbs. live-weight per week.			Weights.		In-crease in 28 days.	Per 1,000 lbs. live-weight per week.					
	May 8.	June 5.		Food consumed.		In-crease in weight.	May 8.	June 5.		Food consumed.		In-creas in weight			
				Grass.	Oilcake.					Grass.	Oilcake.				
1	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>				
2	1,129	1,234	105	} 745	17½	23·9	1,230	1,318	88	} 836	18	24·4			
3	1,130	1,252	122				1,141	1,232	91						
4				1,073	1,186	113
5				1,168	1,316	148
6	1,136	1,252	116						
7	1,066	1,170	104						
8	1,050	1,152	102						
	1,005	1,152	147						
Totals	2,259	2,486	227	8,869	9,778	909			
Means	1,130	1,243	113	1,109	1,222	114			

* Plot "0" is unmeasured land without sewage, and Plot "00" unmeasured land with sewage.
 † Equal parts linseed cake and rape cake.
 ‡ Until May 26th, the two oxen received sewaged grass, the unsewaged not being ready for cutting
 at that date, at which time the animals weighed 1,235 lbs. and 1,255 lbs. respectively.

TABLE VI.—*continued.*

Detailed Record of Food consumed, and Increase yielded, by Oxen fed on Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.

Season 1862.

28 days, from June 5 to July 3.

Dates of weigh- ing.	Two Oxen on Unsewaged Grass.						Eight Oxen on Sewaged Grass.					
	Particulars of the Food consumed.											
	Grass				Oil- cake.†	Grass				C ca		
	From			Quantities (weighed green).		From			Quantities (weighed green).			
	Field.	Plot.*	Crop.			Field.	Plot.*	Crop.				
1862.				tons. cwt. qrs. lbs.	lbs.				tons. cwt. qrs. lbs.			
June	Five-acre	0	1	0 2 0 2	6	Five-acre	3	1	0 7 3 25			
5	"	0	1	0 1 3 5	6	"	4	1	0 9 1 21			
6	"	0	1	0 1 3 26	6	"	4	1	0 7 3 0			
7	"	0	1	0 2 0 0	6	"	4	1	0 7 3 0			
8	"	0	1	0 2 0 1	6	"	4	1	0 7 3 22			
9	"	0	1	0 2 0 20	6	"	4	1	0 10 3 0			
10	"	0	1	0 2 1 9	6	"	4	1	0 8 1 13			
11	"	0	1	0 2 1 12	6	"	4	1	0 9 2 27			
12	"	0	1	0 2 1 9	6	"	4	1	0 9 3 3			
13	"	0	1	0 1 3 11	6	"	4	1	0 6 3 2			
14	"	0	1	0 1 3 13	6	"	4	1	0 6 3 4			
15	"	0	1	0 2 2 7	6	"	4	1	0 8 2 20			
16	"	0	1	0 2 1 11	6	"	4	1	0 8 1 25			
17	"	0	1	0 2 1 1	6	"	4	1	0 8 1 17			
18	"	0	1	0 2 0 9	6	"	3	1	0 7 3 23			
19	"	0	1	0 2 0 23	6	"	3	1	0 8 2 25			
20	"	0	1	0 2 0 2	6	"	3	1	0 7 1 27			
21	"	0	1	0 2 0 4	6	"	3	1	0 7 2 4			
22	"	0	1	0 2 0 1	6	"	3	1	0 5 3 12			
23	"	0	1	0 2 0 2	6	"	2	1	0 11 0 9			
24	"	0	1	0 1 2 14	6	"	2	1	0 8 0 5			
25	"	0	1	0 1 3 9	6	"	2	1	0 7 1 14			
26	"	0	1	0 2 0 8	6	"	00	1	0 6 3 13			
27	"	1	1	0 2 0 1	6	"	2	1	0 6 3 8			
28	"	1	1	0 2 0 0	6	"	2	1	0 6 3 14			
29	"	1	1	0 2 1 5	6	"	2	1	0 7 3 8			
30	"	1	1			"						
July												
1	"	1	1	0 1 3 14	6	"	2	1	0 8 2 4			
2	"	1	1	0 1 3 24	6	"	2	1	0 7 1 22			
Total in 28 Days - - 2 17 3 19 168 Total in 28 days - - 11 7 0 3												
Average per head per day - 0 1 0 4 3 Average per head per day 0 1 0 2												

WEIGHTS AND INCREASE OF THE OXEN, &c.

Nos.	Weights.		In-crease in 28 days.	Per 1,000 lbs. live-weight per week.			Weights.		In-crease in 28 days.	Per 1,000 lbs. live-w per week.		
	June 5.	July 3.		Food consumed.		In-crease in weight.	June 5.	July 3.		Food consumed.		
				Grass.	Oilcake.					Grass.	Oilcake.	
1	lbs. 1,234	lbs. 1,276	lbs. 42	lbs. 636	lbs. 16½	lbs. 12'6	lbs. 1,318	lbs. 1,402	lbs. 84	lbs. 629	lbs. 16½	lbs. 104
2	1,252	1,338	86	1,232	1,330	98
3	1,186	1,290	104
4	1,316	1,390	74
5	1,252	1,328	76
6	1,170	1,220	50
7	1,152	1,205	53
8	1,152	1,256	104
Totals	2,486	2,614	128	9,778	10,421	643
Means	1,243	1,307	64	1,222	1,302	80

unmeasured land without sewage, and Plot "00" unmeasured land with sewage. Inseed cake and rape cake.

TABLE VI.—continued.

Detailed Record of Food consumed, and Increase yielded, by Oxen fed on Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.

Season 1862.

28 days, from July 3 to July 31.

Dates of weigh- ing.		Two Oxen on Unsewaged Grass.						Eight Oxen on Sewaged Grass.							
		Particulars of the Food consumed.													
		Grass.			Oil- cake.†	Grass.			Oil- cake.†						
		From		Quantities (weighed green).		From		Quantities (weighed green).							
Field.	Plot.*	Crop.	Field.		Plot.*	Crop.									
1862.				tons.	cwt.	qrs.	lbs.			tons.	cwt.	qrs.	lbs.		
July	3	Five-acre	1	1	0	1	3 4	6	Five-acre	2	1	0	7 2 15	24	
	4	"	0	1	0	1	3 9	6	"	00 & 2	1	1	0 8 0 18	24	
	5	"	1	1	0	2	0 18	6	"	2	1	0	7 1 8	24	
	6	"	1	1	0	2	0 20	6	"	2	1	0	7 1 13	24	
	7	"	1	1	0	2	0 13	6	"	2	1	0	8 0 21	24	
	8	"	1	1	0	2	0 8	6	"	00, 2 & 4	1 & 2	1	0 8 1 18	24	
	9	"	1	1	0	2	0 19	6	"	2	1	0	6 2 10	24	
	10	"	1	1	0	1	3 6	6	"	2	1	0	6 2 27	24	
	11	"	1	1	0	2	0 15	6	"	2	1	0	6 3 10	24	
	12	"	1	1	0	2	1 9	6	"	2	1	0	8 1 16	24	
	13	"	1	1	0	2	1 8	6	"	2	1	0	8 1 16	24	
	14	"	1	1	0	2	1 0	6	"	2	1	0	7 3 15	24	
	15	"	1	1	0	1	3 15	6	"	2	1	0	7 2 21	24	
	16	"	1	1	0	1	3 14	6	"	2	1	0	5 3 17	24	
	17	"	1	1	0	1	3 11	6	"	2	1	0	6 2 5	24	
	18	"	1	1	0	1	3 21	6	"	2	1	0	8 0 10	24	
	19	"	1	1	0	1	3 10	6	"	2	1	0	5 3 9	24	
	20	"	1	1	0	1	3 14	6	"	2	1	0	5 3 9	24	
	21	"	1	1	0	1	3 10	6	"	2	1	0	5 2 17	24	
	22	"	1	1	0	1	3 4	6	"	2	1	0	5 1 16	24	
	23	"	1	1	0	1	3 4	6	"	4	2	0	8 1 5	24	
	24	"	1	1	0	2	0 9	6	"	4	2	0	8 0 14	24	
	25	"	1	1	0	1	3 8	6	"	4	2	0	9 2 25	24	
	26	"	1	1	0	1	3 11	6	"	4	2	0	7 2 0	24	
	27	"	1	1	0	1	3 14	6	"	4	2	0	7 2 0	24	
	28	"	1	1	0	1	3 19	6	"	4	2	0	8 3 11	24	
	29	"	1	1	0	1	2 5	6	"	4	2	0	8 1 24	24	
	30	"	1	1	0	1	2 2	6	"	4	2	0	8 0 12	24	
Total in 28 days		-	-	-	2	14	1 20	168	Total in 28 days		-	-	-	10 9 1 18	672
Average per head per day		-	-	-	0	0	3 25	3	Average per head per day		-	-	-	0 0 3 21	3

WEIGHTS AND INCREASE OF THE OXEN, &c.

WEIGHTS AND INCREASE OF THE OXEN, &c.												
Nos.	Weights.		In-crease in 28 days.	Per 1,000 lbs. live-weight per week.			Weights.		In-crease (or loss) in 28 days.	Per 1,000 lbs. live-weight per week.		
	July 3.	July 31		Food consumed.		In-crease in weight.	July 3.	July 31.		Food consumed.		In-crease in weight.
				Grass.	Oilcake.					Grass.	Oilcake.	
1	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>
2	1,276	1,332	56	} 570	15½	11·0	1,402	1,426	24	} 556	16	5·6
3	1,338	1,400	62				1,330	1,328	- 2			
4	1,290	1,288	- 2
5	1,390	1,448	58
6	1,328	1,372	44
7	1,220	1,266	46
8	1,205	1,242	37
	1,256	1,286	30
Totals	2,614	2,732	118	10,421	10,656	235
Means	1,307	1,366	59	1,303	1,332	29

* Plot "0" is unmeasured land without sewage, and Plot "00" unmeasured land with sewage.
 † Equal parts linseed cake and rape cake.

TABLE VI.—*continued.*

Detailed Record of Food consumed, and Increase yielded, by Oxen fed on Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Season 1862.

28 days, from July 31 to August 28.

Dates of weigh- ing.	Two Oxen on Unsewaged Grass.						Eight Oxen on Sewaged Grass.								
	Particulars of the Food consumed.														
	Grass				Oil- cake.†	Grass				Oil- cake.†					
	From			Quantities (weighed green).		From			Quantities (weighed green).						
Field.	Plot.*	Crop.			Field.	Plot.	Crop.								
July 31	Five-acre	1	1	tons. cwt. lbs.	0 1 1 20	lbs. 6	Five-acre	4	2	tons. cwt. lbs.	0 6 2 3	lbs. 24			
Aug. 1	"	1	1	0 1 2 13	6	"	4	2	0 8 0 27	24					
2	"	1	1	0 1 1 7	6	"	4	2	0 10 0 16	24					
3	"	1	1	0 1 1 6	6	"	3	2	0 9 0 8	24					
4	"	1	1	0 1 1 9	6	"	3	2	0 8 0 18	24					
5	"	1	1	0 1 2 4	6	"	3	2	0 8 2 15	24					
6	"	1	1	0 1 3 5	6	"	3	2	0 9 3 20	24					
7	"	1	1	0 1 2 12	6	"	3	2	0 11 2 8	24					
8	"	1	1	0 1 1 21	6	"	3	2	0 9 0 22	24					
9	"	1	1	0 1 3 1	6	"	3	2	0 9 0 6	24					
10	"	1	1	0 1 3 0	6	"	3	2	0 9 0 6	24					
11	"	1	1	0 1 1 27	6	"	3	2	0 8 2 16	24					
12	"	1	1	0 1 0 17	6	"	3	2	0 8 3 13	24					
13	"	1	1	0 1 1 9	6	"	3	2	0 8 3 2	24					
14	"	1	1	0 2 1 3	6	"	3	2	0 11 2 7	24					
15	"	1	1	0 1 2 0	6	"	3	2	0 8 1 13	24					
16	"	1	1	0 1 0 21	6	"	3	2	0 8 1 26	24					
17	"	1	1	0 1 0 23	6	"	3	2	0 8 2 0	24					
18	"	1	1	0 1 3 3	6	"	3	2	0 11 0 25	24					
19	"	1	1	0 1 2 1	6	"	3	2	0 9 0 21	24					
20	"	1	1	0 1 2 3	6	"	3	2	0 9 1 20	24					
21	"	0	1	0 1 0 19	6	"	3	2	0 7 2 21	24					
22	"	0	1	0 1 1 7	6	"	2	2	0 9 3 22	24					
23	"	0	1	0 1 0 9	6	"	2	2	0 9 1 18	24					
24	"	0	1	0 1 0 12	6	"	2	2	0 9 1 23	24					
25	"	0	1	0 1 0 5	6	"	2	2	0 8 1 14	24					
26	"	0	1	0 1 0 6	6	"	2	2	0 9 2 9	24					
27	"	0	1	0 1 0 2	6	"	2	2	0 10 0 13	24					
Total in 28 days				-	-	1 19 2 13	168	Total in 28 days				-	-	12 17 0 20	672
Average per head per day				-	-	0 0 2 23	3	Average per head per day				-	-	0 1 0 17	3

WEIGHTS AND INCREASE OF THE OXEN, &c.

Nos.	Weights.		In-crease in 28 days.	Per 1,000 lbs. live-weight per week.			Weights.		In-crease (or loss) in 28 days.	Per 1,000 lbs. live-weight per week.		
	July 31.	Aug. 28.		Food consumed.		In-crease in weight.	July 31.	Aug. 28.		Food consumed.		In-crease (or loss in weight
				Grass.	Oilcake.					Grass.	Oilcake.	
	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	
1	1,332	1,374	42	} 399	15	9'0	1,426	1,420	- 6	} 677	15½	-1'1
2	1,400	1,458	58				1,328	1,272	-56			
3				1,288	1,290	2			
4				1,448	1,458	10			
5				1,372	1,356	14			
6				1,266	1,273	7			
7				1,242	1,240	- 2			
8				1,286	1,272	-14			
Totals	2,732	2,832	100	10,656	10,611	-45
Means	1,366	1,416	50	1,332	1,326	- 6

* Plot "0" is unmeasured land without sewage.

† Equal parts linseed cake and rape cake.

Table VI.—*continued*.

Record of Food consumed, and Increase yielded, by Oxen fed on
unsewaged and Sewaged Meadow Grass, with Oilcake in addition.

Season 1862.

28 days, from August 28 to September 25.

Two Oxen on Unsewaged Grass.					Eight Oxen on Sewaged Grass.				
Particulars of the Food consumed.									
Grass.				Oil- cake.†	Grass.				Oil- cake.†
From			Quantities (weighed green).		From			Quantities (weighed green).	
Field.	Plot.*	Crop.			Field.	Plot.*	Crop.		
			tons. cwt. qrs. lbs.	lbs.				tons. cwt. qrs. lbs.	lbs.
Five-acre	0	1	0 0 3 25	8	Five-acre	2	2	0 9 0 20	32
28	0	1	0 1 0 8	8	28	2	2	0 8 2 15	32
28	0	2	0 1 3 3	8	28	2	2	0 9 2 25	32
28	0	2	0 1 3 5	8	28	2	2	0 9 3 2	32
28	0	2	0 1 3 3	8	28	00 & 2	2	0 11 3 13	32
28	0	2	0 1 3 8	8	28	2	2	0 10 2 20	32
28	0	2	0 1 3 13	8	28	2	2	0 10 3 0	32
28	0	2	0 1 3 16	8	28	2	2	0 9 3 25	32
28	0	2	0 1 3 15	8	28	2	2	0 9 3 26	32
28	0	2	0 1 3 8	8	28	2	2	0 9 3 0	32
28	0	2	0 1 3 10	8	28	2	2	0 9 3 3	32
28	0	2	0 1 3 1	8	28	2	2	0 7 0 20	32
28	0	2	0 1 3 12	8	28	00	2	0 9 0 9	32
28	0	2	0 1 3 15	8	28	00	2	0 9 0 26	32
28	0	2	0 1 3 17	8	28	00 & 4	2 & 3	0 9 1 3	32
28	0	2	0 1 3 18	8	28	4	3	0 10 0 15	32
28	0	2	0 1 3 2	8	28	4	3	0 8 3 22	32
28	0	2	0 1 3 1	8	28	4	3	0 8 3 10	32
28	0	2	0 1 2 23	8	28	4	3	0 11 0 3	32
28	0	2	0 1 2 20	8	28	4	3	0 10 1 11	32
28	0	2	0 1 2 15	8	28	4	3	0 11 1 4	32
28	0	2	0 1 3 2	10	28	4	3	0 11 2 15	40
28	0	2	0 1 2 13	10	28	4	3	0 10 3 15	40
28	0	2	0 1 3 3	10	28	4	3	0 8 3 19	40
28	0	2	0 1 3 6	10	28	4	3	0 8 3 19	40
28	0	2	0 1 3 7	10	28	4	3	0 9 3 17	40
28	0	2	0 1 3 16	10	28	4	3	0 8 3 10	40
28	0	2	0 1 2 21	10	28	4	3	0 10 0 1	40
28 days	-	-	2 8 2 26	238	Total in 28 days	-	-	13 14 2 4	962
per head per day	-	-	0 0 3 13	4½	Average per head per day	-	-	0 1 0 25	4½

WEIGHTS AND INCREASE OF THE OXEN, &c.

Weights.			Per 1,000 lbs. live-weight per week.			Weights.			Per 1,000 lbs. live-weight per week.		
		In- crease (or loss) in 28 days.	Food consumed.		In- crease in weight.			In- crease in 28 days.	Food consumed.		In- crease in weight.
Aug. 28.	Sep. 25.†		Grass.	Oilcake.		Aug. 28.	Sep. 25.†		Grass.	Oilcake.	
lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
374	1,414	40	479	20½	3'3	1,420	1,508	88	700	21½	16'7
458	1,496	— 2				1,272	1,425	153			
"	"	"				1,290	1,403	118			
"	"	"				1,458	1,524	66			
"	"	"				1,386	1,468	82			
"	"	"				1,273	1,334	61			
"	"	"				1,240	1,310	70			
"	"	"	"	"	"	1,272	1,369	97			
832	2,870	38	"	"	"	10,611	11,346	735	"	"	"
416	1,435	19	"	"	"	1,326	1,418	92	"	"	"

* is unmeasured land without sewage, and Plot "00" unmeasured land with sewage.
† cake only, excepting on August 28 and 29, on which days a mixture of linseed cake and rape
cake were in fact, for convenience, taken on September 26, but in the calculations are supposed
September 25.

Table VI.—continued.

Detailed Record of Food consumed, and Increase yielded, by Oxen fed Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.

Season 1862.

18 days, from September 25 to October 13.

Dates of weigh- ing.	Two Oxen on Unsewaged Grass.						Eight Oxen on Sewaged Grass.									
	Particulars of the Food consumed.															
	Grass.				Oil- cake.†	Grass.										
	From		Quantities. (weighed green).	From		Quantities (weighed green).										
Field.	Plot.*	Crop.		Field.	Plot.*		Crop.									
1862.				tons.	cwt.	qrs.	lbs.	lbs.				tons.	cwt.	qrs.	lbs.	
Sept. 25	Five-acre	0	2	0	1	3	24	12	Five-acre	4	3	0	11	1	2	
26	"	0	2	0	1	3	3	12	"	4	3	0	9	3	3	
27	"	0	2	0	1	2	19	12	"	4	3	0	8	3	21	
28	"	0	2	0	1	2	21	12	"	4	3	0	8	3	26	
29	"	0	2	0	1	3	23	12	"	4	3	0	10	1	6	
30	"	0	2	0	1	3	20	12	"	4	3	0	10	0	1	
Oct. 1	"	0	2	0	1	3	23	12	"	3	3	0	13	0	16	
2	"	0	2	0	2	0	6	12	"	3	3	0	9	2	23	
3	"	0	2	0	1	3	3	12	"	2 & 3	3	0	12	0	11	
4	"	1	2	0	1	3	17	12	"	2	3	0	8	3	5	
5	"	1	2	0	1	3	20	12	"	2	3	0	8	3	10	
6	"	1	2	0	1	3	11	12	"	00 & 2	2 & 3	0	10	0	8	
7	"	1	2	0	1	3	8	12	"	2	3	0	11	0	11	
8	"	1	2	0	1	3	9	12	"	2	3	0	9	0	22	
9	"	1	2	0	1	3	9	12	"	2	3	0	9	1	3	
10	"	1	2	0	1	3	24	12	"	2	3	0	9	1	11	
11	"	1	2	0	1	3	6	12	"	2	3	0	7	2	16	
12	"	1	2	0	1	3	8	12	"	2	3	0	7	2	16	
13	Five-acre	1	2	0	1	3	7	16	Five-acre†	00	3	0	11	0	13	
14	"	1	2	0	1	3	14	16	"	4	4	0	9	2	7	
15	"	1	2	0	1	3	8	16	"	4	4	0	8	2	2	
16	"	1	2	0	1	3	11	16	"	4	4	0	8	3	23	
Total in 18 days (to Oct. 13) -				1	13	2	2	216	Total in 18 days (to Oct. 13)				8	16	0	15
Average per head per day -				0	0	3	20	6	Average per head per day				0	1	0	25

WEIGHTS AND INCREASE OF THE OXEN, &c.

Nos.	Weights.		In-crease (or loss) in 18 days.	Per 1,000 lbs. live-weight per week.			Weights.		In-crease (or loss) in 18 days.	Per 1,000 lbs. live-weight per week.				
	Sept. 25. §	Oct. 13.		Food consumed.		In-crease (or loss) in weight.	Sept. 25. §	Oct. 13.		Food consumed.		Grass.	Oilcake.	or (or loss) in weight.
				Grass.	Oilcake.					Grass.	Oilcake.			
1	<i>lbs.</i> 1,414	<i>lbs.</i> 1,356	} 515 29½ -10·3	<i>lbs.</i> 1,508	<i>lbs.</i> 1,488	<i>lbs.</i> -20	} 681 29½ -	<i>lbs.</i> 1,425	<i>lbs.</i> 1,378	<i>lbs.</i> -47				
2	1,456	1,439		1,408	1,414	6								
3								
4								
5	1,524	1,531		7						
6	1,468	1,442		-26						
7	1,334	1,332		-2						
8	1,310	1,288		-22						
	1,369	1,311	-58								
Totals	2,870	2,795	-75	11,346	11,184	-162			
Means	1,435	1,397	-38	1,418	1,398	-20			

* Plot "0" is unmeasured land without sewage, and plot "00" unmeasured land with sewage.

† Linseed cake.

On October 11, 3 cwt. 0 qrs. 13 lbs. on October 13, the whole quantity, viz. 7 cwt. 8 qrs. 16 lbs. and 8 cwt. 2 qrs. 23 lbs. of the grass came from another (not experimental) field. weights were in fact, for convenience, taken on September 26, but in the calculations apply to September 25.

TABLE VII.
SUMMARY OF THE WEIGHTS AND INCREASE OF THE OXEN FED RESPECTIVELY ON UNSEWAGED AND SEWAGED MEADOW GRASS, WITH OILCAKE IN SOLUTION.
SEASON 1962.

	Weights.							Increase in Weight.			
								May 8 to Oct. 13 = 22½ weeks.†			
	May 8.*	May 29.*	June 5.	July 3.	July 31.	Aug. 28.	Sept. 25.†	Oct. 13.	Total.	Per head live-weight per week.	Per 1,000 lbs. live-weight per week.
TWO OXEN ON UNSEWAGED GRASS.											
1	1,129	1,225	1,254	1,276	1,332	1,374	1,414	1,536	227	10 1	8 2
2	1,130	1,255	1,252	1,338	1,400	1,458	1,456	1,459	309	10 1	7 8
Totals -	2,259	2,480	2,486	2,614	2,732	2,832	2,870	2,795	536	—	—
Averages -	1,130	1,240	1,243	1,307	1,366	1,416	1,435	1,398	268	11 14	9 6
									309	—	—
									155	8 5	6 5
EIGHT OXEN ON SEWAGED GRASS.											
1	1,230	—	1,318	1,402	1,426	1,420	1,508	1,458	258	11 7	8 7
2	1,141	—	1,252	1,330	1,328	1,372	1,425	1,378	237	10 8	8 5
3	1,073	—	1,186	1,290	1,288	1,290	1,408	1,414	341	15 2	12 2
4	1,168	—	1,316	1,390	1,448	1,438	1,524	1,531	363	16 1	11 14
5	1,136	—	1,202	1,328	1,372	1,386	1,468	1,442	306	13 9	10 8
6	1,063	—	1,170	1,220	1,266	1,273	1,334	1,332	266	11 13	9 13
7	1,050	—	1,162	1,205	1,242	1,240	1,310	1,288	238	10 9	8 12
8	1,005	—	1,152	1,256	1,286	1,272	1,369	1,311	305	13 9	11 11
Totals -	8,869	—	9,778	10,421	10,556	10,611	11,546	11,184	2,315	—	—
Averages -	1,109	—	1,222	1,303	1,322	1,326	1,418	1,398	289	12 13	10 4
									176	9 7	7 4

* Both lots of animals were weighed on May 8, but as the unsewaged grass was not ready to cut until May 29, the 2 animals were then re-weighed.
† Sept. 25 was the proper day of weighing, but as a matter of convenience, it did not, in fact, take place until Sept. 26.
‡ This period includes the first 5 weeks during which the 2 oxen, otherwise fed on unsewaged grass, received sewaged grass.
§ This period excludes the first month of the experiment during 3 weeks of which the 2 oxen, otherwise fed on unsewaged grass, received sewaged grass.

TABLE VIII.
SUMMARY OF FOOD CONSUMED AND INCREASE YIELDED BY THE OXEN FED ON UNSEWAGED AND SEWAGED MEADOW GRASS, WITH OILCAKE IN ADDITION.
SEASON 1862.

Periods.		TWO OXEN ON UNSEWAGED GRASS.						EIGHT OXEN ON SEWAGED GRASS.					
Dates.	Number of Weeks.	Food consumed.				Increase (or Loss) in Weight.		Food consumed.				Increase (or Loss) in Weight.	
		Per head per week.		Per 1,000 lbs. live-weight per week.		Per head per week.	Per 1,000 lbs. live-weight per week.	Per head per week.		Per 1,000 lbs. live-weight per week.		Per head per week.	Per 1,000 lbs. live-weight per week.
		Grass.	Oilcake.	Grass.	Oilcake.			Grass.	Oilcake.	Grass.	Oilcake.		
		<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs. ozs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs.</i>	<i>lbs. ozs.</i>
May 8 to June 5 -	4	884	21	745	17½	23	6	974	21	886	18	23	6
June 5 to July 3 -	4	811	21	686	16½	16	0	765	21	629	16½	20	3
July 3 to July 31 -	4	763	21	570	13½	14	13	733	21	556	16	7	6
July 31 to Aug. 28 -	4	555	21	389	15	13	8	900	21	677	15½	-1	6
Aug. 28 to Sept. 25 -	4	683	29½	479	20½	4	13	961	29½	700	21½	23	0
Sept. 25 to Oct. 13 -	2½	730	43	515	29½	-14	9	950	43	681	29½	-7	14
May 8 to Oct. 13 *	22½	738	25	594	19½	11	14	882	25	704	20	13	13
June 8 to Oct. 13 †	19½	706	23½	535	19½	8	5	863	23½	608	19½	9	7

* This period includes the first 5 weeks during which the 2 oxen, otherwise fed on unsewaged grass, received sewaged grass.
† This period excludes the first month of the experiment during 3 weeks of which the 2 oxen, otherwise fed on unsewaged grass, received sewaged grass.

DETAILED RECORD OF FOOD CONSUMED, AND MILK YIELDED, BY COWS FED ON THE OREGON PRATER
SECOND SEASON, 1862; 7 days.—May 2 to May 8.

DETAILED RECORD OF FOOD CONSUMED, AND THE WEIGHTS OF THE MILK PRODUCED, IN THE SECOND SEASON, 1862; 7 days.—May 2 to May 9.													
		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.			
THREE COWS.—UNSEWAGED GRASS.													
Food consumed.	Grass	From which Field, Plot, and Crop.							Tons.	cwt.	qrs.	lbs.	
		Quantities weighed		Oatcake (equal parts linseed and rape cake)									
Yield of Milk, &c.	Cows.	Breed.	Years old.	Dates of Calving.	Weights (May 1).								
					lbs.								
1 2 3	2 3 3	Cross Sht.-horn	6	Mar. 20	945								
		Cross Sht.-horn	7	Dec. 25	976								
		Cross Sht.-horn	Aged	Feb. 26	884								
Totals -					2,805								
Means -					935								

TWELVE COWS.—SEWAGED GRASS.

[illegible]

Table IX.—continued.

Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Second Season, 1862; 7 days.—May 9 to May 15.

THREE COWS.—UNSEWAGED GRASS.																												
Food consumed.	Grass	From which Field, Plot, and Crop.	Friday.			Saturday.			Sunday.			Monday.			Tuesday.			Wednesday.			Thursday.			Total in 7 days.	Per head per day.			
			Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.					
Field of Milk, &c.	Cows.	Breed.	Years old.	Dates of Calving.	Weights of (May 1st)			Quantities weighed			Oilcake (equal parts linseed and rape cake)			Quantities weighed			Oilcake (equal parts linseed and rape cake)			Quantities weighed			Oilcake (equal parts linseed and rape cake)					
					Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.			
					A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.
					A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.
1	Cross Shit-horn	6	Mar. 20	966	13	5	21	10	13	3	028	13	5	21	10	13	3	028	13	5	21	10	13	3	028			
2	Cross Shit-horn	7	Dec. 25	1,064	13	12	0	14	6	11	4	0	4	2	10	0	4	2	10	0	4	2	10	0	4			
3	Cross Shit-horn	Aged	Feb. 26	896	19	8	20	9	14	0	22	0	15	0	20	8	12	3	20	8	12	3	20	8	12			
Totals -				2,866	55	0	40	11	56	9	38	7	56	0	36	8	56	2	36	3	53	13	36	6	54			
Means -				955	18	5	13	10	18	14	12	13	18	11	12	3	18	11	12	1	17	13	12	2	18			

TWELVE COWS.—SEWAGED GRASS.																												
Food consumed.	Grass	From which Field, Plot, and Crop.	Friday.			Saturday.			Sunday.			Monday.			Tuesday.			Wednesday.			Thursday.			Total in 7 days.	Per head per day.			
			Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.					
Field of Milk, &c.	Cows.	Breed.	Years old.	Dates of Calving.	Weights of (May 1st)			Quantities weighed			Oilcake (equal parts linseed and rape cake)			Quantities weighed			Oilcake (equal parts linseed and rape cake)			Quantities weighed			Oilcake (equal parts linseed and rape cake)					
					Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.	Tons.	Cwts.	Lbs.			
					A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.
					A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.
1	Cross Shit-horn	8	Nov. 18	1,148	14	0	9	8	12	9	10	6	12	10	13	4	10	6	12	10	13	4	10	6	12			
2	Cross Shit-horn	8	Feb. 2	1,138	14	1	9	13	1	11	3	11	3	13	1	11	3	13	1	11	3	13	1	11	3			
3	Cross Shit-horn	6	Dec. 1	1,076	14	1	10	14	6	11	8	13	1	11	9	8	13	1	11	9	8	13	1	11	9			
4	Cross Shit-horn	Aged	Jan. 26	968	15	10	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10			
5	Cross Shit-horn	6	Nov. 17	1,076	13	10	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10			
6	Cross Shit-horn	Aged	Jan. 13	1,066	15	10	9	13	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10	13	10			
7	Cross Shit-horn	Aged	Apr. 28	1,126	26	9	17	4	26	0	18	1	26	4	18	6	20	4	18	6	20	4	18	6	20			
8	Cross Shit-horn	3	Apr. 13	723	14	4	8	15	14	2	9	14	1	9	13	10	9	13	10	9	13	10	9	13	10			
9	Cross Shit-horn	Aged	Apr. 16	965	14	0	10	5	14	6	9	14	6	9	14	6	9	14	6	9	14	6	9	14	6			
10	Cross Shit-horn	Aged	Apr. 3	866	21	2	14	5	20	8	14	5	20	8	14	5	20	8	14	5	20	8	14	5	20			
11	Cross Shit-horn	6	Mar. 30	781	16	3	11	7	17	0	13	0	17	4	12	10	15	8	11	0	15	8	11	0	15			
12	Cross Shit-horn	7	Nov. 18	984	16	3	11	7	17	0	13	0	17	4	12	10	15	8	11	0	15	8	11	0	15			
Totals -				11,488	148	0	108	18	148	0	108	18	148	0	108	18	148	0	108	18	148	0	108	18	148	0		
Means -				957	12	0	9	15	12	0	9	15	12	0	9	15	12	0	9	15	12	0	9	15	12	0		

Table IX.—continued.

Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Second Season, 1863; 7 days.—May 23 to May 29.

THREE COWS.—UNSEWAGED GRASS.																			
		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.		Per head per day.	
Food consumed.	Grass	From which Field, Plot, and Crop.		Ten-acre, Plot 0,* Crop 1.		Ten-acre, Plot 0,* Crop 1.		Ten-acre, Plot 0,* Crop 1.		Ten-acre, Plot 0,* Crop 1.		Ten-acre, Plot 0,* Crop 1.		Ten-acre, Plot 0,* Crop 1.		—		—	
		Quantities weighed		tons.		cwt.		qrs.		lbs.		tons.		cwt.		qrs.		tons.	
		Oilcake (equal parts linseed and rape-cake)		cwt.		qrs.		lbs.		tons.		cwt.		qrs.		lbs.		tons.	
		Weights (May 29).		lbs.		tons.		cwt.		qrs.		lbs.		tons.		cwt.		qrs.	
Yield of Milk, &c.	Cows.	Breed.		A.M.		P.M.		A.M.		P.M.		A.M.		P.M.		A.M.		P.M.	
		Cross Shit-horn		17 10 13		10 8		18 10 13		13 6		17 7 13		14 13		18 13 4		18 13 4	
		Cross Shit-horn		15 10 13		8 12		12 6		10 13		11 5		11 5		12 13		12 13	
		Cross Shit-horn		20 13 6		12 9		19 7		13 6		19 6		18 3		17 3		17 3	
Totals		3,044		32 12 37 11		51 11		50 5 38 9		49 5		36 10		46 9 43 0		16 6		49 3 41 3	
Means		1,015		17 9 12 9		17 4		16 12 14		16 7		12 3		13 8		14 5		13 12	
TWELVE COWS.—SEWAGED GRASS.																			
		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.		Per head per day.	
Food consumed.	Grass	From which Field, Plot, and Crop.		Ten-acre, Plot 3, Crop 1.		Five-acre, Plot 3, Crop 1.		Five-acre, Plot 3, Crop 1.		Five-acre, Plot 3, Crop 1.		Five-acre, Plot 3, Crop 1.		Five-acre, Plot 3, Crop 1.		—		—	
		Quantities weighed		tons.		cwt.		qrs.		lbs.		tons.		cwt.		qrs.		tons.	
		Oilcake (equal parts linseed and rape-cake)		cwt.		qrs.		lbs.		tons.		cwt.		qrs.		lbs.		tons.	
		Weights (May 29).		lbs.		tons.		cwt.		qrs.		lbs.		tons.		cwt.		qrs.	
Yield of Milk, &c.	Cows.	Breed.		A.M.		P.M.		A.M.		P.M.		A.M.		P.M.		A.M.		P.M.	
		Cross Shit-horn		12 10 13		10 4		12 14		16 7		12 3		13 8		14 5		13 12	
		Cross Shit-horn		15 10 13		8 12		12 6		10 13		11 5		11 5		12 13		12 13	
		Cross Shit-horn		20 13 6		12 9		19 7		13 6		19 6		18 3		17 3		17 3	
Totals		3,044		32 12 37 11		51 11		50 5 38 9		49 5		36 10		46 9 43 0		16 6		49 3 41 3	
Means		1,015		17 9 12 9		17 4		16 12 14		16 7		12 3		13 8		14 5		13 12	

[illegible]

Table IX.—continued.
 Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake addition.
 Second Season 1862; 7 days.—June 6 to June 12.

		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.
THREE COWS.—UNSEWAGED GRASS.										
Food con- sumed.	Grass	From which Field, Plot, and Crop.		Ten-acre, Plot 1, Crop 1.	Ten-acre, Plot 1, Crop 1.	Ten-acre, Plot 1, Crop 1.	Ten-acre, Plot 1, Crop 1.	Ten-acre, Plot 1, Crop 1.		
	Oilcake (equal parts linseed and rape cake)	lbs.	qrs.	lbs.	qrs.	lbs.	qrs.	lbs.	qrs.	lbs.
		tons.	cwt.	tons.	cwt.	tons.	cwt.	tons.	cwt.	tons.
		0 0 9	0 0 0	0 0 15	0 0 4	0 0 4	0 0 4	0 0 9	1 6 2	0 0 0
Yield of Milk, &c.	Breed.	Years old.	Dates of Calving.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
	1	6	May 29.	16 3	10 14	11 6	11 6	16 10	12 10	198 5
	2	7	Mar 20	15 12	10 14	11 10	10 8	10 14	12 0	187 3
	3	7	Dec. 25	10 4	11 15	15 8	11 6	16 8	12 11	198 15
Totals				42 3	35 8	43 12	33 13	43 12	38 11	554 7
Means				14 1	11 13	14 9	11 4	14 8	12 11	184 13
TWELVE COWS.—SEWAGED GRASS.										
Food con- sumed.	Grass	From which Field, Plot, and Crop.		Ten-acre, Plot 1, Crop 1.	Ten-acre, Plot 2, Crop 1.	Ten-acre, Plot 2, Crop 1.	Ten-acre, Plot 2, Crop 1.	Ten-acre, Plot 2, Crop 1.		
	Oilcake (equal parts linseed and rape cake)	lbs.	qrs.	lbs.	qrs.	lbs.	qrs.	lbs.	qrs.	lbs.
		tons.	cwt.	tons.	cwt.	tons.	cwt.	tons.	cwt.	tons.
		0 10 1	0 0 1	0 14 2	0 14 2	0 14 2	0 14 2	0 14 2	0 14 2	0 14 2
Yield of Milk, &c.	Breed.	Years old.	Dates of Calving.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
	1	6	Nov. 18	12 3	10 3	11 4	9 0	11 4	9 0	147 0
	2	8	Feb. 12	11 11	9 8	11 10	8 12	11 6	9 8	144 5
	3	8	Dec. 1	11 8	9 4	11 6	8 10	11 7	9 8	137 15
Yield of Milk, &c.	Breed.	Years old.	Dates of Calving.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
	4	6	Jan. 26	10 14	8 12	11 6	7 13	11 7	9 8	141 8
	5	6	Nov. 17	11 6	9 0	10 4	8 10	11 3	8 2	131 11
	6	6	Jan. 13	13 7	9 0	12 11	9 11	13 4	10 4	155 0
Yield of Milk, &c.	Breed.	Years old.	Dates of Calving.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
	7	6	May 12	20 11	14 6	19 3	14 11	20 12	15 11	245 6
	8	3	Apr. 15	10 6	8 14	10 6	7 9	10 1	7 14	119 12
	9	3	Feb. 16	11 3	8 1	12 2	10 0	13 11	10 0	152 11
Yield of Milk, &c.	Breed.	Years old.	Dates of Calving.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
	10	6	Apr. 3	12 10	9 7	13 2	10 0	13 4	10 6	210 3
	11	6	Apr. 3	12 4	10 3	13 5	10 13	13 13	11 5	163 7
	12	6	Apr. 3	12 4	10 3	13 5	10 13	13 13	11 5	163 7

THREE COWS.—UNWEALED GRASS.

[illegible]

TWELVE COWS.—SEWAGED GRASS.

[illegible]

Table IX.—continued.

Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Second Season, 1862; 7 days.—June 20 to June 26.

		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.												
THREE COWS.—UNSEWAGED GRASS.																						
Food consumed.	Grass	From which Field, Plot and Crop.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		—												
		Quantities weighed (equal parts linseed and rape cake)		lbs.		cwt.		cwt.														
		Tons.		cwt.		cwt.		cwt.														
Field of Milk, &c.	Breed.	Years old.	Dates of Calving.	Weights of (June 26).		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		—												
				1		2		3														
				Cross Sht.-horn		Cross Sht.-horn		Cross Sht.-horn														
1	2	3	Mar. 20	1,092	1,092	1,092	1,092	1,092	1,092	1,092												
2	3	7	Dec. 23	1,083	1,083	1,083	1,083	1,083	1,083	1,083												
3			Feb. 26	978	978	978	978	978	978	978												
Totals				3,161	42 12	32 6	41 12	32 6	43 13	33 9	40 13	33 3	40 14	32 15	41 5	33 12	40 0	34 12	524	4		
Means				1,054	14 4	10 13	13 14	10 13	14 10	11 3	13 10	11 1	13 10	11 0	13 12	11 4	13 5	11 9	174	12	24	15

TWELVE COWS.—SEWAGED GRASS.

Food consumed.	Grass	From which Field, Plot, and Crop.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre, Plot 2, Crop 1.			Ten-acre,		
----------------	-------	-----------------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	---------------------------	--	--	-----------	--	--

THREE COWS.—UNWEAVERED GRASS.

Food con- sumed.	Grass	From which Field, Plot, and Crop.				Weights (June 25).		Years old.	Dates of Calving.	Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																								Ten-acre, Plot 1, Crop 1.																							
------------------------	-------	-----------------------------------	--	--	--	-----------------------	--	---------------	-------------------------	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	------------------------------	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

TWELVE COWS.—SEWAGED GRASS.

Food con- sumed.	Grass	From which Field, Plot, and Crop.		Breed.	Years old.	Dates of Calving.	Ten-acre, Plot 2, Crop 1.		Ten-acre, Plot 2, Crop 1.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Tons. cwt.	qrs. lbs.	Tons. cwt.	qrs. lbs.																
		Tons. cwt.	qrs. lbs.				Tons. cwt.	qrs. lbs.	Tons. cwt.	qrs. lbs.	Tons. cwt.	qrs. lbs.	Tons. cwt.	qrs. lbs.	Tons. cwt.	qrs. lbs.	Tons. cwt.	qrs. lbs.																				
ed. D- ed.	Oilcake (rape-cake)	Quantities weighed	-	-	-	Weights of (June 20, 76).	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	Tons. cwt.	qrs. lbs.	Tons. cwt.	qrs. lbs.																
							Tons.	qtrs.	lbs.	Tons.	qtrs.	lbs.	Tons.	qtrs.	lbs.	Tons.	qtrs.	lbs.					Tons.	qtrs.	lbs.	Tons.	qtrs.	lbs.										
							11	11	11	11	11	11	11	11	11	11	11	11					11	11	11	11	11	11	11	11	11							
							12	12	12	12	12	12	12	12	12	12	12	12					12	12	12	12	12	12	12	12	12							
							13	13	13	13	13	13	13	13	13	13	13	13					13	13	13	13	13	13	13	13	13							
							14	14	14	14	14	14	14	14	14	14	14	14					14	14	14	14	14	14	14	14	14							
							15	15	15	15	15	15	15	15	15	15	15	15					15	15	15	15	15	15	15	15	15	15						
							16	16	16	16	16	16	16	16	16	16	16	16					16	16	16	16	16	16	16	16	16	16						
							17	17	17	17	17	17	17	17	17	17	17	17					17	17	17	17	17	17	17	17	17	17	17					
							18	18	18	18	18	18	18	18	18	18	18	18					18	18	18	18	18	18	18	18	18	18	18					
							19	19	19	19	19	19	19	19	19	19	19	19					19	19	19	19	19	19	19	19	19	19	19					
							20	20	20	20	20	20	20	20	20	20	20	20					20	20	20	20	20	20	20	20	20	20	20					
Totals -				12,894	141	4	102	5	139	4	103	9	147	13	114	8	153	11	96	0	139	13	113	11	152	4	103	14	154	13	106	10	1,753	7				
Means -				1,075	11	12	8	8	11	10	8	13	12	5	9	9	11	2	8	0	11	11	9	8	12	11	8	13	12	14	9	14	146	2				

Table IX.—continued.
Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Second Season, 1862; 7 days.—July 4 to July 10.

		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.			
THREE COWS.—UNSEWAGED GRASS.													
Cows.	Grass.	From which Field, Plot, and Crop.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		—			
		Breed.	Years old.	Dates of Calving.	Weights of (July 10).	Quantities weighed (rap-cake)	Tons.	cwt.	lbs.		Tons.	cwt.	lbs.
1	Cross Sht.-horn	6	Mar. 20	1,057	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
2	Cross Sht.-horn	7	Dec. 25	1,104	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
3	Cross Sht.-horn	Aged	Feb. 20	988	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
Totals		-		3,175	41 6 33 3	40 8 28 12	40 7 31 2	39 5 29 12	33 14 30 4	40 9 32 15	41 0 29 13	49 14	-
Means		-		1,058	13 13 11 1	13 8 9 9	13 8 10 6	13 2 9 15	11 5 10 1	13 8 11 0	13 10 9 15	16 1/2 5	23 8

TWELVE COWS.—SEWAGED GRASS.																	
Cows.	Grass.	From which Field, Plot, and Crop.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		—							
		Breed.	Years old.	Dates of Calving.	Weights of (July 10).	Quantities weighed (rap-cake)	Tons.	cwt.	lbs.		Tons.	cwt.	lbs.	Tons.	cwt.	lbs.	
1	Cross Sht.-horn	6	Nov. 18	1,190	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	
2	Cross Sht.-horn	6	Feb. 12	1,293	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
3	Cross Sht.-horn	6	Dec. 1	1,204	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
4	Cross Sht.-horn	Aged	Jan. 25	983	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
5	Cross Sht.-horn	Aged	Nov. 17	1,132	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
6	Cross Sht.-horn	Aged	Jan. 13	1,118	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
7	Cross Sht.-horn	Aged	May 2	1,029	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
8	Cross Sht.-horn	3	Apr. 13	822	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2
9	Cross Sht.-horn	Aged	Feb. 16	1,082	0 2 0 21 1/2	0 2 2 10 1/2	0 3 1 8 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2	0 3 0 10 1/2

		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.		Average per day.	
THREE COWS.—UNSEWAGED GRASS.																			
Grass		From which Field, Plot, and Crop.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.	
Quantities weighed		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.	
Oilcake (1 part linseed and 3 parts rape cake)		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.	
		Weights (July 10).		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.	
		Breed.		Years of Calving.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.	
		Cross Sht.-horn		6		1,087		9 3		9 12		10 9		10 5		10 5		10 5	
		Cross Sht.-horn		7		1,100		9 3		9 12		10 9		10 5		10 5		10 5	
		Cross Sht.-horn		Aged		988		13 8		12 6		13 6		10 8		13 6		10 8	
		Totals		-		3,175		37 4		31 9		35 15		30 3		37 4		31 9	
		Means		-		1,058		12 7		10 8		12 0		10 1		12 7		10 1	
TWELVE COWS.—SEWAGED GRASS.																			
Grass		From which Field, Plot, and Crop.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.		Ten-acre, Plot 4, Crop 2.	
Quantities weighed		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.	
Oilcake (1 part linseed and 3 parts rape cake)		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.	
		Weights (July 10).		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.	
		Breed.		Years of Calving.		lbs.		cwt.		lbs.		cwt.		lbs.		cwt.		lbs.	
		Cross Sht.-horn		6		1,100		10 7		10 7		10 7		10 7		10 7		10 7	
		Cross Sht.-horn		8		1,293		10 11		10 11		10 11		10 11		10 11		10 11	
		Cross Sht.-horn		Aged		953		11 6		11 6		11 6		11 6		11 6		11 6	
		Cross Sht.-horn		Aged		1,132		12 14		12 14		12 14		12 14		12 14		12 14	
		Cross Sht.-horn		Aged		1,118		12 14		12 14		12 14		12 14		12 14		12 14	
		Cross Sht.-horn		Aged		1,069		12 14		12 14		12 14		12 14		12 14		12 14	
		Cross Sht.-horn		Aged		822		12 5		12 5		12 5		12 5		12 5		12 5	
		Cross Sht.-horn		Aged		1,082		12 15		12 15		12 15		12 15		12 15		12 15	
		Cross Sht.-horn		Aged		918		17 11		17 11		17 11		17 11		17 11		17 11	
		Cross Sht.-horn		Aged		828		13 0		13 0		13 0		13 0		13 0		13 0	
		Cross Sht.-horn		Aged		1,110		13 1		13 1		13 1		13 1		13 1		13 1	
		Totals		-		12,755		154 14		145 13		169 9		150 10		166 6		151 8	
		Means		-		1,594		19 27		18 1		21 1		19 12		20 8		18 10	

Table IX.—*continued.*
 Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
 Second Season, 1862; 7 days.—July 18 to July 24.

THREE COWS.—UNSEWAGED GRASS.											
From which Field, Plot, and Crop.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.	
Grass	Oilcake	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	
Quantities weighed		0 0 3	0 0 18	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Oftake (1 part linseed and 3/4 parts rape cake)		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Weights of (July 24).		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Years old.		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Calving.		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Breed.		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
1 Cross Sht.-horn		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
2 Cross Sht.-horn		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
3 Cross Sht.-horn		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Totals		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Means		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
1,070		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	

THREE COWS.—UNSEWAGED GRASS.											
From which Field, Plot, and Crop.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.		Ten-acre, Plot 0, Crop 1.	
Grass	Oilcake	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	
Quantities weighed		0 0 3	0 0 18	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Oftake (1 part linseed and 3/4 parts rape cake)		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Weights of (July 24).		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Years old.		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Calving.		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Breed.		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
1 Cross Sht.-horn		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
2 Cross Sht.-horn		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
3 Cross Sht.-horn		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Totals		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
Means		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	
1,070		0 0 0	0 0 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	0 0 1 13	

TWELVE COWS.—SEWAGED GRASS.											
From which Field, Plot, and Crop.		Ten-acre, Plot 3, Crop 2.		Ten-acre, Plot 3, Crop 2.		Ten-acre, Plot 3, Crop 2.		Ten-acre, Plot 3, Crop 2.		Ten-acre, Plot 3, Crop 2.	
Grass	Oilcake	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	
Quantities weighed		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
Oftake (1 part linseed and 3/4 parts rape cake)		0 0 0	0 0 1 26	0 0 1 26	0 0 1 26	0 0 1 26	0 0 1 26	0 0 1 26	0 0 1 26	0 0 1 26	
Weights of (July 24).		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
Years old.		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
Calving.		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
Breed.		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
1 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
2 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
3 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
4 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
5 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
6 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
7 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
8 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
9 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
10 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
11 Cross Sht.-horn		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
Totals		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
Means		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	
1,070		0 0 13	0 0 1 27	0 0 1 26	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	0 0 1 27	

field of milk, &c.

Table IX.—continued.

THREE COWS.—UNSEWAGED GRASS.														Per head per day.			
		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.	
From which Field, Plot, and Crop	Grass	Five-acre, Plot 1, Crop 1.		Five-acre, Plot 1, Crop 1.		—		—		Ten-acre, Plot 0*, Crop 2		Ten-acre, Plot 0*, Crop 2		—		—	
		tons.	cwt.	gvs.	lbs.	tons.	cwt.	gvs.	lbs.	tons.	cwt.	gvs.	lbs.	tons.	cwt.	gvs.	lbs.
Quantity weighed	Oilcake	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Breed.	Dates of Calving.	Weights of (July 24).		—		—		—		—		—		—		—	
		Years old.	—		—		—		—		—		—		—		—
1	Cross Sht.-horn	6	Mar 30	1,080	—	—	—	—	—	—	—	—	—	—	—	—	—
2	Cross Sht.-horn	7	Dec. 26	1,127	—	—	—	—	—	—	—	—	—	—	—	—	—
3	Cross Sht.-horn	Agd	Feb. 28	1,011	—	—	—	—	—	—	—	—	—	—	—	—	—
Totals		39	13	29	1	30	12	25	9	33	6	27	13	35	2	26	13
Means		13	4	9	11	10	4	8	9	11	2	9	4	11	12	8	15

[illegible]

		From which Field, Plot, and Crop.		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.		Per head per day.	
Food con- sumed.	Grass	Ton-acre, Plot 0, Crop 2.		+		+		+		+		+		+		+		-		-	
		Quantities weighed		lbs.		tons.		cwt.		qrs.		lbs.		tons.		cwt.		qrs.		lbs.	
		A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 10	15 1	10 15	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5	13 0	10 12	14 6	10 5
	Oilcake (rape-cake)	0 7	1 13	0 6	0 0	0 8	0 13	0 0	0 12	0 0	0 12	0 4	0 16	0 0	0 12	0 0	0 12	0 0	0 12	0 0	0 4
	Grass	14 6	10 0	13 13	10 11	15 5	10 1														

Table IX.—continued.
Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Second Season, 1862; 7 days.—August 15 to August 21.

THREE COWS.—UNSEWAGED GRASS.																																																																																																																																																																																																																																																																																																																																																																																																																
Food on-med.	Grass	From which Field, Plot, and Crop.		Weights (Aug. 21).		Dates of Calving.		Years old.		Breed.																																																																																																																																																																																																																																																																																																																																																																																																						
		Quantities weighed		lbs.		lbs.		lbs.		lbs.																																																																																																																																																																																																																																																																																																																																																																																																						
		Oilcake (rape-cake)	Oilcake (rape-cake)	tons.	qtrs.	lbs.	tons.	qtrs.	lbs.	tons.	qtrs.																																																																																																																																																																																																																																																																																																																																																																																																					
field of ilk, &c.	Cows.	Nos.	Breed.	Years old.	Dates of Calving.	Weights (Aug. 21).	lbs.	tons.	qtrs. <td rowspan="3">lbs.</td> <td rowspan="3">tons.</td>	lbs.	tons.																																																																																																																																																																																																																																																																																																																																																																																																					
												Cross Sht.-horn	6	Mar. 20	1,076	10 10 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12	0 0 0 12

May 22 to August 28.

Food consumed.		From which Field, Plot, and Crop.		Grass		Oats (equal parts linseed and rape cake)		Quantities weighed		Weights (Aug. 21).		Dates of Calving.		Breed.		Years old.		Cows.		Milk, &c.	
										</											

[illegible]

Table IX.—continued.

Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.

Second Season, 1862: 7 days.—August 29 to September 4.

[illegible]

Second Season, 1862; 7 days.—September 5 to September 11.

		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
THREE COWS.—UNSEWAGED GRASS.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																					
Food consumed.	From which Field, Plot, and Crop.	Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
		tons.	cwt.s.	tons.	cwt.s.	tons.	cwt.s.	tons.	cwt.s.	tons.	cwt.s.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
		lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
Yield of Milk, &c.	Oilcake (equal parts linseed and rape cake)	0	3	2	11½	0	3	1	23½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½	0	3	3	24½

TWELVE COWS.—SEWAGED GRASS.

Food consumed.	Grass	From which Field, Plot, and Crop.		Ten-acre, Plot 4, Crop 3.		Ten-acre, Plot 4, Crop 3.		Ten-acre, Plot 3 and 4, Crop 3.		Ten-acre, Plot 3 and 4, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop	
----------------	-------	-----------------------------------	--	---------------------------	--	---------------------------	--	---------------------------------	--	---------------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	---------------------------	--	------------------------	--

Table IX.—continued.

Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.

Second Season, 1862; 7 days.—September 12 to September 18.

THREE COWS.—UNSEWAGED GRASS.																								
Food Consumed.	From which Field, Plot, and Crop.		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.													
	Grass	Oileake (equal parts linseed and rape cake)	Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Tons.	Cwt.	Lbs.											
			Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.				Lbs.										
															Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.				
Quantities weighed		Tons.		Cwt.		Lbs.		Tons.		Cwt.		Lbs.												
Yield of Milk, &c.	Breed.	Years old.	Dates of Calving.	Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).										
				Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.									
																Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1										
	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2									
	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3									
	Totals		3,276		3,276		3,276		3,276		3,276		3,276		3,276									
	Means		1,092		1,092		1,092		1,092		1,092		1,092		1,092									

TWELVE COWS.—SEWAGED GRASS.																								
Food Consumed.	From which Field, Plot, and Crop.		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.													
	Grass	Oileake (equal parts linseed and rape cake)	Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Ten-acre, Plot 3, Crop 3.		Tons.	Cwt.	Lbs.											
			Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.				Lbs.										
															Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.				
Quantities weighed		Tons.		Cwt.		Lbs.		Tons.		Cwt.		Lbs.												
Yield of Milk, &c.	Breed.	Years old.	Dates of Calving.	Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).		Weights (Sept. 18).										
				Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.									
																Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.	Tons.	Cwt.	Lbs.
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1										
	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2									
	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3									
	Totals		12,910		12,910		12,910		12,910		12,910		12,910		12,910									
	Means		1,630		1,630		1,630		1,630		1,630		1,630		1,630									

Table IX.—continued.
Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Second Season, 1862; 7 days.—September 26 to October 2.

THREE COWS.—UNSEWAGED GRASS.																												
Food consumed.		Grass		From which Field, Plot, and Crop.		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.		Per head per day.						
Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights of (Sept. 18), lbs.	Ten-acre, Plot 0*, Crop 2.			Ten-acre, Plot 0*, Crop 2.			Ten-acre, Plot 0*, Crop 3.			Ten-acre, Plot 0*, Crop 3.			Ten-acre, Plot 0*, Crop 3.			tons.	qrs.	lbs.						
					Quantities weighed			Quantities weighed			Quantities weighed			Quantities weighed			Quantities weighed											
					tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.				tons.	qrs.	lbs.			
1	Cross Sht.-horn	6	Mar. 20	1,068	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
2	Cross Sht.-horn	7	Dec. 25	1,190	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
3	Cross Sht.-horn	8	Feb. 28	1,020	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0					
Totals				3,276	34	2	25	14	32	12	25	7	32	4	23	1	26	8	26	10	32	2	20	10	380	0	18	9
Means				1,092	11	6	8	10	10	13	8	8	10	3	8	2	10	12	7	13	10	11	6	14	130	0	18	9

TWELVE COWS.—SEWAGED GRASS.																												
Food consumed.		Grass		From which Field, Plot, and Crop.		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.		Per head per day.						
Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights of (Sept. 18), lbs.	Five-acre, Plot 3, Crop 3.			Five-acre, Plot 3, Crop 3.			Five-acre, Plot 3, Crop 3.			Five-acre, Plot 3, Crop 3.			Five-acre, Plot 3, Crop 3.			Ten-acre, Plot 2, Crop 3.			Ten-acre, Plot 2, Crop 3.			tons.	qrs.	lbs.
					Quantities weighed			Quantities weighed			Quantities weighed			Quantities weighed			Quantities weighed			Quantities weighed			Quantities weighed					
					tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.	qrs.	lbs.			
1	Cross Sht.-horn	6	Nov. 18	1,270	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	Cross Sht.-horn	8	Feb. 12	1,267	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
3	Cross Sht.-horn	6	Dec. 1	1,288	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	Cross Sht.-horn	6	Jan. 26	1,073	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	Cross Sht.-horn	6	Nov. 17	1,199	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6	Cross Sht.-horn	6	Jan. 13	1,150	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
7	Cross Sht.-horn	6	May 2	1,106	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8	Cross Sht.-horn	3	Apr. 13	860	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
9	Cross Sht.-horn	3	Apr. 13	1,232	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10	Ayrshire	6	Feb. 16	1,000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
11	Guernsey	4	Mar. 30	824	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
12	Cross Sht.-horn	7	Nov. 16	1,146	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

THREE COWS.—UNSEWAGED GRASS.

[illegible]

TWELVE COWS.—SEWAGED GRASS.

[illegible]

at This cow calved again Oct. 7, 1862.

† Not experimental.

Unmeasured land, designated Plot 0 when unsewaged.

Table IX.—continued.
 Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
 Second Season, 1862; 7 days.—October 10 to October 16.

THREE COWS.—UNSEWAGED GRASS.										
Food consumed.	Grass	From which Field, Plot, and Crop.		Quantities weighed		Weights (Oct. 10).		Ten-acre, Plot 3, Crop 2.		—
		Oilcake (3 parts linseed and 2 parts rape cake)		Oats		lbs.		tons.		
		Breed.		Years of Calving.		lbs.		qrs.		
		Cows.		old.		lbs.		cwt.		
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2	Cross Sht.-horn	7	Dec. 25	1,219	1,219	0 3 1	0 27	—
		3	Cross Sht.-horn	7	Feb. 25	1,058	1,058	0 3 1	0 15	—
		Totals		—		3,335	3,335	0 10 2	0 57	—
		Means		—		1,112	1,112	0 3 4	0 19	—
THREE COWS.—UNSEWAGED GRASS.										
Field of milk, &c.	Grass	1	Cross Sht.-horn	6	Mar. 20	1,058	1,058	0 4 0	0 15	—
		2</								

Third Season, 1863; 7 days.—April 28 to May 4.

FIVE COWS.—UNSEWAGED MEADOW GRASS.

Grass consumed.	From which Field, Plot, and Crop	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights (April 27) lbs.	Per head per day.						
							Monday.	Tuesday.	Wednesday.	Thursday.	Friday.	Saturday.	Sunday.
Yield of Milk, &c.	Quantities weighed (tons, cwt, qrs, lbs.)	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.
		A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
1	Cross short-horn Aged	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
2	Cross short-horn 6	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
3	Cross short-horn 3	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
4	Cross short-horn 7	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
5	Cross short-horn 7	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
Totals		89 11	68 4	84 10	67 13	81 1	74 0	66 9	81 9	65 9	73 2	80 6	68 4
Means		17 15	13 10	17 5	13 9	16 5	14 10	13 5	16 5	13 2	14 6	16 1	14 1

TEN COWS.—SEWAGED MEADOW GRASS.

Grass consumed.	From which Field, Plot, and Crop	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights (April 27) lbs.	Per head per day.						
							Monday.	Tuesday.	Wednesday.	Thursday.	Friday.	Saturday.	Sunday.
Yield of Milk, &c.	Quantities weighed (tons, cwt, qrs, lbs.)	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.
		A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
1	Cross short-horn Aged	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
2	Cross short-horn 6	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
3	Cross short-horn 3	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
4	Cross short-horn 7	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
5	Cross short-horn 7	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
6	Cross short-horn 6	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
7	Cross short-horn 6	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
8	Cross short-horn 6	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
9	Cross short-horn 7	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
10	Half-short-horn Aged	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
Totals		177 4	131 4	171 0	123 8	138 0	118 8	127 5	121 7	123 11	123 12	123 12	123 12
Means		17 11	13 2	17 1	12 6	13 13	11 14	12 2	12 2	12 6	12 6	12 6	12 6

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

Grass consumed.	From which Field, Plot, and Crop	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights (April 27) lbs.	Per head per day.						
							Monday.	Tuesday.	Wednesday.	Thursday.	Friday.	Saturday.	Sunday.
Yield of Milk, &c.	Quantities weighed (tons, cwt, qrs, lbs.)	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.	Rye, Plot 1, Crop 1.
		A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
1	Cross short-horn Aged	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
2	Cross short-horn 6	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
3	Cross short-horn 3	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
4	Cross short-horn 7	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
5	Cross short-horn 7	17 6	12 8	18 4	15 3	14 10	17 6	12 8	18 4	15 3	14 10	17 6	12 8
Totals		89 11	68 4	84 10	67 13	81 1	74 0	66 9	81 9	65 9	73 2	80 6	68 4
Means		17 15	13 10	17 5	13 9	16 5	14 10	13 5	16 5	13 2	14 6	16 1	14 1

by Cows fed on Unsewaged and Sewaged Meadow Grass, and on Italian Eye Grass (without Oilcake). Third Season, 1863 : 7 days.—May 5 to May 11.

				Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.		Per head per day.	

TEN COWS—SEWAGED MEADOW GRASS.

[illegible]

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)										From which Field, Plot, and Crop																						
	R_{10} Plot 10, Crop 1.	R_{10} Plot 07, Crop 1.	R_{10} Plot 01, Crop 1.	R_{10} Plot 07, Crop 1.	R_{10} Plot 01, Crop 1.	R_{10} Plot 07, Crop 1.	R_{10} Plot 01, Crop 1.	R_{10} Plot 07, Crop 1.	R_{10} Plot 01, Crop 1.	R_{10} Plot 07, Crop 1.																							
1. Grass short-horn, aged	1,096	281	6	17	10	268	0	25	2	283	18	1	35	2	16	0	37	4	18	4	37	8	16	7	34	10	30	6	501	11	43	2	
2. Grass short-horn, aged	April 20	1,096	281	6	17	10	268	0	25	2	283	18	1	35	2	16	0	37	4	18	4	37	8	16	7	34	10	30	6	501	11	43	2
3. Grass short-horn, aged	May 1	1,096	281	6	17	10	268	0	25	2	283	18	1	35	2	16	0	37	4	18	4	37	8	16	7	34	10	30	6	501	11	43	2
4. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
5. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
6. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
7. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
8. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
9. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
10. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
11. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
12. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
13. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0
14. Grass short-horn, aged	April 15	1,155	227	2	16	5	225	8	19	6	231	10	3	22	1	22	15	23	10	26	0	31	2	25	0	34	11	34	0	396	5	40	0

Cows	Breed.	Years old.	Dates of Calving.	From which Field, Plot, and Crop.		Friday.		Saturday.		Sunday.		Monday.		Tuesday.		Wednesday.		Thursday.		Total in 7 days.		Per head per day.	
				Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.
THREE COWS.—UNSEWAGED GRASS.																							
				Quantities weighed (equal parts linseed and rape cake)		tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.
1	Cross Sht.-horn	6	Mar. 20	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	1,076	
2	Cross Sht.-horn	7	Dec. 25	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	1,170	
3	Cross Sht.-horn	Agd	Feb. 26	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	1,012	
Totals				3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	3,258	
Means				1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	1,086	
TWELVE COWS.—SEWAGED GRASS.																							
				Quantities weighed (equal parts linseed and rape cake)		tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.	tons.	cwt.	lbs.
1	Cross Sht.-horn	6	Nov. 12	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	
2	Cross Sht.-horn	8	Feb. 2	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	
3	Cross Sht.-horn	6	Dec. 1	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	
4	Cross Sht.-horn	Agd	Jan. 26	973	973	973	973	973	973	973	973	973	973	973	973	973	973	973	973	973	973	973	
5	Cross Sht.-horn	6	Nov. 17	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	1,158	
6	Cross Sht.-horn	Agd	Jan. 13	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	1,125	
7	Cross Sht.-horn	6	May 2	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	1,109	
8	Cross Sht.-horn	6	Apr. 13	834	834	834	834	834	834	834	834	834	834	834	834	834	834	834	834	834	834	834	
9	Cross Sht.-horn	Agd	Apr. 16	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	1,136	
10	Cross Sht.-horn	6	Apr. 3	950	950	950	950	950	950	950	950	950	950	950	950	950	950	950	950	950	950	950	
11	Cross Sht.-horn	6	Nov. 12	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	1,224	
12	Cross Sht.-horn	7	Nov. 12	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	1,237	

[illegible]

Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., lbs.)										Five-acre, Plot 2, Crop 1.					Five-acre, Plot 2, Crop 1.					Five-acre, Plot 2, Crop 1.					Five-acre, Plot 2, Crop 1.											
		Five-acre, Plot 2 & 3/4, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.										
Yield of milk, etc.	1	1,125	17	6	13	0	59	0	13	5	30	7	13	13	91	1	14	5	18	1	13	0	4	13	12	10	4	13	12	4	13	3	19	0	1	13		
	2	1,010	253	8	12	5	32	12	14	25	12	15	9	25	6	15	12	13	12	13	12	13	18	2	13	12	10	4	13	12	4	13	3	19	0	1	13	
	3	1,060	200	12	1	17	0	23	6	13	6	13	6	13	6	13	6	13	6	13	6	13	18	2	13	12	10	4	13	12	4	13	3	19	0	1	13	
	4	1,280	14	9	11	12	16	10	13	7	11	16	4	11	4	16	14	10	14	10	14	10	14	10	4	13	12	10	4	13	12	4	13	3	19	0	1	13
	5	984	14	6	11	4	13	15	9	11	16	4	11	4	16	14	10	14	10	14	10	14	10	4	13	12	10	4	13	12	4	13	3	19	0	1	13	
	6	1,040	19	8	12	1	21	2	12	12	21	4	14	9	23	8	14	2	22	1	12	5	20	6	13	12	10	4	13	12	4	13	3	19	0	1	13	
	7	1,038	17	0	12	5	18	8	11	11	29	5	13	0	20	8	12	15	18	10	11	17	2	10	4	13	12	10	4	13	12	4	13	3	19	0	1	13
	8	1,054	19	1	13	6	16	1	19	1	19	1	13	8	11	1	11	1	11	1	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	
	9	1,054	19	1	13	6	16	1	19	1	19	1	13	8	11	1	11	1	11	1	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	
	10	1,258	12	5	8	10	12	8	8	11	12	2	9	10	13	0	8	11	13	10	8	11	13	8	9	10	6	13	12	4	13	3	19	0	1	13		
	Total	10,312	105	1	124	15	177	12	121	12	180	8	131	3	180	13	182	13	173	12	128	0	174	6	121	9	100	6	105	13	2,065	10	—	—	—	—		
	Mean	1,031	10.5	12	8	17	12	12	12	18	1	13	1	18	1	18	1	18	1	17	1	12	0	17	6	12	1	10	6	10	13	2,065	0	—	—	—		

[illegible]

Table IX—continued.
Detailed Record of Food consumed, and Milk yielded, by Cows fed on the Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.
Second Season, 1862; 7 days.—September 12 to September 18.

		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	Total in 7 days.	Per head per day.
THREE COWS.—UNSEWAGED GRASS.										
Food consumed.	Grass	From which Field, Plot, and Crop.		Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.	Ten-acre, Plot 1, Crop 2.
		Quantities weighed		tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.
		Oilcake (equal parts linseed and rape cake)		tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.
		Weights of (Sept. 18).		A.M.	P.M.	P.M.	A.M.	P.M.	P.M.	P.M.
Milk, &c.	1	Years old.	Dates of Calving.	14 15	14 15	14 15	14 15	14 15	14 15	14 15
	2	Breed.	6	Mar. 20	1,096	1,096	1,096	1,096	1,096	1,096
	3	Cross Sht.-horn	7	Dec. 25	1,190	1,190	1,190	1,190	1,190	1,190
	4	Cross Sht.-horn	Agd	Feb. 23	1,020	1,020	1,020	1,020	1,020	1,020
Totals				37 11	30 4	37 4	31 7	34 2	26 3	36 11
Means				12 9	10 1	12 7	10 8	11 6	8 12	12 4
				12 9	10 1	12 7	10 8	11 6	8 12	12 4
TWELVE COWS.—SEWAGED GRASS.										
Food consumed.	Grass	From which Field, Plot, and Crop.		Ten-acre, Plot 3, Crop 3.	Ten-acre, Plot 3, Crop 3.	Ten-acre, Plot 3, Crop 3.	Ten-acre, Plot 3, Crop 3.	Ten-acre, Plot 3, Crop 3.	Ten-acre, Plot 3, Crop 3.	Ten-acre, Plot 3, Crop 3.
		Quantities weighed		tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.
		Oilcake (equal parts linseed and rape cake)		tons.	qrs.	lbs.	tons.	qrs.	lbs.	tons.
		Weights of (Sept. 18).		A.M.	P.M.	P.M.	A.M.	P.M.	P.M.	P.M.
Milk, &c.	1	Years old.	Dates of Calving.	10 14	7 14	13 4	10 14	7 14	13 4	10 14
	2	Breed.	6	Nov. 18	1,270	1,270	1,270	1,270	1,270	1,270
	3	Cross Sht.-horn	8	Dec. 2	1,297	1,297	1,297	1,297	1,297	1,297
	4	Cross Sht.-horn	Agd	Jan. 26	973	973	973	973	973	973
Totals	Means			37 11	30 4	37 4	31 7	34 2	26 3	36 11
				12 9	10 1	12 7	10 8	11 6	8 12	12 4
				12 9	10 1	12 7	10 8	11 6	8 12	12 4
				12 9	10 1	12 7	10 8	11 6	8 12	12 4

		Friday.	Saturday.	Sunday.	Monday.	Tuesday.	Wednesday.	Thursday.	7 days.	per day.
THREE COWS.—UNSEWAGED GRASS.										
Food con- sumed.	From which Field, Plot, and Crop.	Grass		Oil-cake (equal parts linseed and rape cake)		Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		—
		lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	
Field of Milk, &c.	Breed.	Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		—
		lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	
Cows.	Years old.	Weights (Stent. 18)		Weights (Stent. 18)		Weights (Stent. 18)		Weights (Stent. 18)		—
		Calving.	Dec.	Calving.	Dec.	Calving.	Dec.	Calving.	Dec.	
1	Cross Sht.-horn	6	Mar. 20	1,065	Mar. 20	1,065	Mar. 20	1,065	Mar. 20	1,065
2	Cross Sht.-horn	7	Dec. 25	1,190	Dec. 25	1,190	Dec. 25	1,190	Dec. 25	1,190
3	Cross Sht.-horn	8	Feb. 26	1,020	Feb. 26	1,020	Feb. 26	1,020	Feb. 26	1,020
Totals		3,275				3,275				
Means		1,092				1,092				
TWELVE COWS.—SEWAGED GRASS.										
Food con- sumed.	From which Field, Plot, and Crop.	Grass		Oil-cake (equal parts linseed and rape cake)		Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		—
		lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	
Field of Milk, &c.	Breed.	Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		Ten-acre, Plot 0.7 Crop 2.		—
		lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	lbs.	qtrs.	
Cows.	Years old.	Weights (Stent. 18)		Weights (Stent. 18)		Weights (Stent. 18)		Weights (Stent. 18)		—
		Calving.	Dec.	Calving.	Dec.	Calving.	Dec.	Calving.	Dec.	
1	Cross Sht.-horn	6	Nov. 18	1,270	Nov. 18	1,270	Nov. 18	1,270	Nov. 18	1,270
2	Cross Sht.-horn	8	Feb. 2	1,257	Feb. 2	1,257	Feb. 2	1,257	Feb. 2	1,257
3	Cross Sht.-horn	6	Dec. 1	1,284	Dec. 1	1,284	Dec. 1	1,284	Dec. 1	1,284
4	Cross Sht.-horn	6	Nov. 29	1,073	Nov. 29	1,073	Nov. 29	1,073	Nov. 29	1,073
5	Cross Sht.-horn	6	Nov. 17	1,199	Nov. 17	1,199	Nov. 17	1,199	Nov. 17	1,199
6	Cross Sht.-horn	6	Nov. 13	1,150	Nov. 13	1,150	Nov. 13	1,150	Nov. 13	1,150
7	Cross Sht.-horn	6	May 2	1,098	May 2	1,098	May 2	1,098	May 2	1,098
8	Cross Sht.-horn	6	Feb. 16	1,252	Feb. 16	1,252	Feb. 16	1,252	Feb. 16	1,252
9	Cross Sht.-horn	6	Apr. 3	1,090	Apr. 3	1,090	Apr. 3	1,090	Apr. 3	1,090
10	Cross Sht.-horn	6	Nov. 16	1,146	Nov. 16	1,146	Nov. 16	1,146	Nov. 16	1,146
Total		13,275				13,275				
Means		1,327				1,327				

Second Season, 1862; 7 days.—September 26 to October 2.

THREE COWS.—UNSEWAGED GRASS.									
From which Field, Plot, and Crop.		Breed.		Years old.	Dates of Calving.	Weights of (Sept. 18).		Quantities weighed - Oilcake (3 parts lined and 2 parts rape cake)	
Cow No.								Tons.	cwt.
1	Cross Shit-horn	6	Mar. 20	1,066					
2	Cross Shit-horn	7	Dec. 25	1,190					
3	Cross Shit-horn	Aged	Feb. 28	1,020					
Totals				3,276					
Means				1,092					
TWELVE COWS.—SEWAGED GRASS.									
From which Field, Plot, and Crop.		Breed.		Years old.	Dates of Calving.	Weights of (Sept. 18).		Quantities weighed - Oilcake (3 parts lined and 2 parts rape cake)	
Cow No.								Tons.	cwt.
1	Cross Shit-horn	6	Nov. 18	1,270					
2	Cross Shit-horn	8	Feb. 2	1,297					
3	Cross Shit-horn	6	Dec. 1	1,288					
4	Cross Shit-horn	Aged	Jan. 29	973					
5	Cross Shit-horn	6	Nov. 17	1,199					
6	Cross Shit-horn	Aged	Jan. 13	1,156					
7	Cross Shit-horn	6	May 2	1,106					
8	Cross Shit-horn	3	Apr. 13	850					
9	Cross Shit-horn	Aged	Feb. 16	1,252					
10	Ayrshire	6	Apr. 3	900					
11	Guernsey	4	Mar. 20	834					

FIVE COWS.—UNSEWAGED.													
Grass consumed.	From which Field, Plot, and Crop			Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Yield of Milk, &c.	
	Rye, Plot 2, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.				
	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.			
1	0 3 0 0		0 5 2 0		0 7 0 0		0 6 0 0		0 3 0 8		0 8 0 7	115 0 15	0 1 0 0
2													
3													
4													
5													
Totals	5,032	89 11	68 4	88 10	67 13	81 1	74 0	86 9	61 9	81 0	65 9	82 0	20 4
Means	1,010	17 15	13 10	17 5	13 9	16 4	14 13	17 5	12 5	16 35	13 2	16 6	11 14

TEN COWS.—SEWAGED MEADOW GRASS.

Grass consumed, lb.	From which Field, Plot, and Crop		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.		Five-acre, Plot 4, Crop 1.	
------------------------	----------------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--	----------------------------	--

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

Grass consumed.	From which Field, Plot, and Crop										Yield of Milk, &c.								
	Quantities weighed (tons, cwt., qrs., lbs.)		Rye, Plot 2, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.										
	1	2	3	4	5	6	7	8	9	10									
1	Cross short-horn	April 20	1,066	25 6	15 3	25 1	17 0	24 4	20 10	23 8	16 9	21 13	18 4	20 8	15 11	37 9	15 0	284	6
2	Ayrshire	May 13	874	22 8	17 3	27 10	16 14	22 3	35 1	21 12	14 1	21 8	14 5	21 0	13 6	23 2	15 8	523	0
3	Cross short-horn	May 13	1,250	18 10	9 3	27 6	18 8	27 4	50 14	23 10	10 14	23 1	14 2	27 0	13 8	23 8	15 6	1,166	5
4	Half short-horn	May 13	1,114	25 6	15 3	25 4	19 0	13 0	13 0	17 4	13 14	15 1	13 0	10 10	17 12	30 10	11 12	280	14
5	Half short-horn	Feb. 8	1,114	15 9	15 13	25 4	19 0	13 0	11 9	15 12	11 2	13 0	10 10	17 12	30 10	11 12	210	1	
Totals			5,229	103 7	75 3	116 7	81 15	115 5	88 10	108 14	77 8	84 6	57 15	66 4	63 11	94 13	55 4	1,089	5

Table X.—continued.
 Diet Record of Food consumed, and Milk yielded, by Cows fed on Unsewaged and Sewaged Meadow Grass, and on Italian Rye Grass (without Oilcake).
 Third Season, 1863; 7 days.—May 5 to May 11.

FIVE COWS.—UNSEWAGED MEADOW GRASS.																								
Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Cows. Nos.	Breed.	Years old.	Dates Calving.	Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.	Per head per day.		
							Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.				Rye, Plot 1, Crop 1.	
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.			A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	1	1013	1013	Mar. 1	0 6 2 3	0 9 0 16	0 4 0 0	0 4 0 0	0 5 2 20	0 6 1 16	0 3 2 7	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	1 18 3 11	0 1 0 12		
	2	Cross short-horn	2	1117	1117	Feb. 15	0 6 2 3	0 9 0 16	0 4 0 0	0 4 0 0	0 5 2 20	0 6 1 16	0 3 2 7	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	1 18 3 11	0 1 0 12		
	3	Cross short-horn	3	1117	1117	Feb. 15	0 6 2 3	0 9 0 16	0 4 0 0	0 4 0 0	0 5 2 20	0 6 1 16	0 3 2 7	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	1 18 3 11	0 1 0 12		
	4	Cross short-horn	4	1117	1117	Feb. 15	0 6 2 3	0 9 0 16	0 4 0 0	0 4 0 0	0 5 2 20	0 6 1 16	0 3 2 7	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	1 18 3 11	0 1 0 12		
	5	Cross short-horn	5	1117	1117	Feb. 15	0 6 2 3	0 9 0 16	0 4 0 0	0 4 0 0	0 5 2 20	0 6 1 16	0 3 2 7	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	0 3 2 6	1 18 3 11	0 1 0 12		
Totals							6038	68 0 58 6	84 7 56 9	84 10 55 12	85 0 31 1	87 8 54 12	85 14 50 9	87 0 57 7	900 15	—	—	—	—	—	—			
Means							1203	16 6 11 11	16 14 11 5	16 15 11 2	17 0 10 4	17 8 10 15	17 3 10 2	17 6 11 8	186 3	—	—	—	—	—	—			
TEN COWS.—SEWAGED MEADOW GRASS.																								
Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Cows. Nos.	Breed.	Years old.	Dates Calving.	Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.	Per head per day.		
							Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.				Rye, Plot 1, Crop 1.	
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.			A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	1	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22		
	2	Cross short-horn	2	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
	3	Cross short-horn	3	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
	4	Cross short-horn	4	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
	5	Cross short-horn	5	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
Totals							770 120 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6		
Means							154 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.																								
Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Cows. Nos.	Breed.	Years old.	Dates Calving.	Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.	Per head per day.		
							Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.		Rye, Plot 1, Crop 1.				Rye, Plot 1, Crop 1.	
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.			A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	1	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22		
	2	Cross short-horn	2	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
	3	Cross short-horn	3	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
	4	Cross short-horn	4	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
	5	Cross short-horn	5	1064	1064	Mar. 10	0 15 1 19	0 12 0 6	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22	0 13 2 22			
Totals							770 120 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6	77 9 125 6		
Means							154 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	15 4 24 1	

Third Season, 1863; 7 days.—May 19 to May 25.

FIVE COWS.—UNSEWAGED MEADOW GRASS.											
From which Field, Plot, and Crop	Grass consumed.	Quantities weighed (tons, cwt., lbs.)	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.
Cows.	Breed.	Years old.	Dates of Calving.	Weights (May 25) lbs.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.
10	Cross short-horn	Aged	May 1	1,064	10 6	2 0	10 6	2 0	10 6	2 0	10 6
2	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
3	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
4	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
5	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
6	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
7	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
8	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
9	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
10	Cross short-horn	Aged	May 15	1,084	10 6	2 0	10 6	2 0	10 6	2 0	10 6
Totals				5,254	85 8	31 14	84 2	49 2	85 8	31 14	84 2
Means				1,053	17 2	10 6	10 13	9 13	16 10	11 1	17 2
TEN COWS.—SEWAGED MEADOW GRASS.											
From which Field, Plot, and Crop	Grass consumed.	Quantities weighed (tons, cwt., lbs.)	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.
Cows.	Breed.	Years old.	Dates of Calving.	Weights (May 25) lbs.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.
1	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
2	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
3	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
4	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
5	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
6	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
7	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
8	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
9	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
10	Cross short-horn	Aged	Mar 10	1,125	17 2	10 6	10 13	9 13	16 10	11 1	17 2
Totals				10,212	172 3	125 13	184 14	126 12	175 12	137 11	175 12
Means				1,021	17 4	12 9	18 8	12 11	17 9	13 19	17 8
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.											
From which Field, Plot, and Crop	Grass consumed.	Quantities weighed (tons, cwt., lbs.)	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.	Plot of Crop 1.
Cows.	Breed.	Years old.	Dates of Calving.	Weights (May 25) lbs.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.
1	Cross short-horn	Aged	Apr 20	1,122	17 2	10 6	10 13	9 13	16 10	11 1	17 2
2	Cross short-horn	Aged	Apr 20	1,122	17 2	10 6	10 13	9 13	16 10	11 1	17 2
3	Cross short-horn	Aged	Apr 20	1,122	17 2	10 6	10 13	9 13	16 10	11 1	17 2
4	Cross short-horn	Aged	Apr 20	1,122	17 2	10 6	10 13	9 13	16 10	11 1	17 2
5	Cross short-horn	Aged	Apr 20	1,122	17 2	10 6	10 13	9 13	16 10	11 1	17 2
Totals				5,254	85 8	31 14	84 2	49 2	85 8	31 14	84 2
Means				1,053	17 2	10 6	10 13	9 13	16 10	11 1	17 2

FIVE COWS.—UNSEWAGED MEADOW GRASS.															
Grass consumed.	From which Field, Plot, and Crop			Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.	
	Quantities weighed (tons, cwt., qrs., lbs.)			+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.	
	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights of (May 23).	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	Aged	May 1	1,054	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	2	Cross short-horn	Aged	Feb. 15	1,184	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	3	Cross short-horn	Aged	April 13	819	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	4	Cross short-horn	Aged	May 20	1,054	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	5	Cross short-horn	Aged	April 17	1,112	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
Totals					5,274	108 9	65 2	107 9	67 9	104 15	67 14	102 13	71 14	104 2	65 15
Means					1,055	21 11	13 1	21 8	13 8	21 0	13 9	20 15	12 9	20 9	14 6
TEN COWS.—SEWAGED MEADOW GRASS.															
Grass consumed.	From which Field, Plot, and Crop			Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.	
	Quantities weighed (tons, cwt., qrs., lbs.)			+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.	
	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights of (May 23).	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	Aged	Mar. 10	1,156	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	2	Cross short-horn	Aged	April 7	1,010	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	3	Cross short-horn	Aged	Feb. 10	1,054	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	4	Cross short-horn	Aged	Feb. 15	1,054	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	5	Cross short-horn	Aged	Feb. 22	954	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
Yield of Milk, &c.	6	Cross short-horn	Aged	April 14	954	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	7	Cross short-horn	Aged	April 12	858	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	8	Cross short-horn	Aged	April 16	1,054	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	9	Cross short-horn	Aged	April 13	1,054	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	10	Cross short-horn	Aged	Oct. 30	1,054	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
Totals					10,212	165 1	124 15	177 12	121 12	160 8	131 3	182 13	113 12	174 6	102 12
Means					1,021	16 10	12 8	17 12	12 3	18 1	13 2	18 11	13 4	17 6	12 13
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.															
Grass consumed.	From which Field, Plot, and Crop			Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.	
	Quantities weighed (tons, cwt., qrs., lbs.)			+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.		+ Crop 1.	
	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights of (May 23).	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	Aged	April 20	1,122	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	2	Cross short-horn	Aged	Mar. 15	902	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	3	Cross short-horn	Aged	Mar. 1	1,106	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	4	Cross short-horn	Aged	Jan. 15	1,176	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
	5	Cross short-horn	Aged	Jan. 2	1,048	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1	'820 1
Totals					5,324	106 7	72 1	107 0	67 13	110 5	66 6	108 2	62 15	109 14	73 11

Table X.—continued.
Dead Record of Food consumed, and Milk yielded, by Cows fed on Unsewaged and Sewaged Meadow Grass, and on Italian Rye Grass (without Oilcake).
 Third Season, 1863; 7 days.—June 2 to June 8.

FIVE COWS.—UNSEWAGED MEADOW GRASS.																			
Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Cows Nos.	Breed.	Year old.	Dates of Calving.	Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.	Rye, Crop 1.	Rye, Crop 1.	Total in 7 days.	Per head per day.		
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.						A.M.	P.M.
1	Cross short-horn	Aged	May 1	1,092 7	14 13 10	0 5 8 14	0 4 2 2	0 9 3 14	0 4 2 0	0 4 0 0	0 5 2 4	0 5 2 4	1 19 3 10	0 1 0 15					
Yield of Milk, lbs.	2	Cross short-horn	Aged	Feb. 15	1,177 4	14 13 10	0 5 8 14	0 4 2 2	0 9 3 14	0 4 2 0	0 4 0 0	0 5 2 4	0 5 2 4	1 19 3 10	0 1 0 15				
	3	Cross short-horn	8	April 13	821 2	14 6 17	16 15 9 9	1 5 4 0	10 4 10 9	8 13 5 2	8 13 5 2	14 10 9 0	14 14 9 0	171 3 24 7	25 0 6				
	4	Cross short-horn	8	May 20	1,094 2	24 12 17	25 3 12	23 6 10	27 4 16	12 20 7	19 1 25	18 4 26	16 8 26	306 5 43 12	45 9 32				
	5	Cross short-horn	7	April 17	1,110 2	20 6 12	30 10 12	8 12 8	20 7 12	13 10 20	7 11 12	20 3 11	1 1 12	227 12 32 9	32 9 9				
	Totals		4,264		98 13 63 13		105 6 61 1		100 12 63 14		96 1 64 15		98 10 63 12		96 6 59 14		1,133 9		
Means		1,066		19 12 12 12		29 11 12 13		25 2 12 12		19 3 13 0		19 12 12 12		19 6 12 1		226 11			
TEN COWS.—SEWAGED MEADOW GRASS.																			
Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Cows Nos.	Breed.	Year old.	Dates of Calving.	Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.		Rye, Crop 1.	Rye, Crop 1.	Rye, Crop 1.	Total in 7 days.	Per head per day.		
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.						A.M.	P.M.
1	Cross short-horn	Aged	Mar. 10	1,136	17 3 13 6	17 3 13 6	0 11 2 2	0 17 0 9	0 12 1 23	0 5 2 7	0 14 3 8	0 14 3 13	4 5 2 21	0 1 0 25					
Yield of Milk, lbs.	2	Cross short-horn	Aged	April 7	1,010	21 3 13 6	22 4 13 14	13 11 14	12 6 12	13 11 14	12 6 12	13 11 14	12 6 12	13 11 14	12 6 12	35 7 30 6	5 7 30 6		
	3	Cross short-horn	6	Feb. 20	1,060	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12		
	4	Cross short-horn	6	Feb. 15	960	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12	16 6 11 12		
	5	Cross short-horn	8	Feb. 22	884	13 11 10 14	14 7 19 8	15 3 10 5	10 5 10	14 8 11 7	16 2 9 8	15 2 10 7	13 4 18 12	17 7 14 24	24 13 24 13	24 13 24 13	24 13 24 13		
	6	Cross short-horn	8	April 14	884	13 11 10 14	14 7 19 8	15 3 10 5	10 5 10	14 8 11 7	16 2 9 8	15 2 10 7	13 4 18 12	17 7 14 24	24 13 24 13	24 13 24 13	24 13 24 13		
Yield of Milk, lbs.	7	Cross short-horn	Aged	April 14	1,096	19 11 12 10	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12		
	8	Cross short-horn	8	April 16	1,096	12 11 12 10	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12		
	9	Cross short-horn	8	April 13	954	28 6 32 10	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12	12 6 17 12		
	10	Half short-horn	Aged	Oct. 30	1,558	9 3 5 10	10 6 6 14	10 11 6 8	11 2 6 13	10 12 7 0	9 2 1 10	8 6 4 1	1 10 8 2	15 7 15 7	15 7 15 7	15 7 15 7	15 7 15 7		
	Totals		10,312		161 3 117 10		165 3 120 8		164 3 121 11		166 4 124 14		170 6 122 0		165 5 101 3		164 5 100 3		
Means		1,031		16 2 11 12		16 8 12 1		16 7 12 3		16 10 12 8		17 1 12 3		16 8 10 2		16 5 10 0			
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.																			
Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Cows Nos.	Breed.	Year old.	Dates of Calving.	Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.	Rye, Plot 3, Crop 2.	Rye, Plot 3, Crop 2.	Total in 7 days.	Per head per day.		
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.						A.M.	P.M.
1	Cross short-horn	Aged	April 20	1,122	28 6 18 8	28 6 18 8	0 7 1 9	0 7 2 7	0 11 0 23	0 4 3 25	0 6 3 30	0 6 3 25	2 12 1 36	0 1 2 0					
Yield of Milk, lbs.	2	Cross short-horn	Aged	April 20	1,122	28 6 18 8	28 6 18 8	0 7 1 9	0 7 2 7	0 11 0 23	0 4 3 25	0 6 3 30	0 6 3 25	2 12 1 36	0 1 2 0				
	3	Cross short-horn	8	April 15	1,122	28 6 18 8	28 6 18 8	0 7 1 9	0 7 2 7	0 11 0 23	0 4 3 25	0 6 3 30	0 6 3 25	2 12 1 36	0 1 2 0				
	4	Cross short-horn	8	April 15	1,122	28 6 18 8	28 6 18 8	0 7 1 9	0 7 2 7	0 11 0 23	0 4 3 25	0 6 3 30	0 6 3 25	2 12 1 36	0 1 2 0				
	5	Cross short-horn	8	April 15	1,122	28 6 18 8	28 6 18 8	0 7 1 9	0 7 2 7	0 11 0 23	0 4 3 25	0 6 3 30	0 6 3 25	2 12 1 36	0 1 2 0				

FIVE COWS.—UNSEWAGED MEADOW GRASS.												
From which Field, Plot, and Crop		Ten-acre, Plot 1 st , Crop 1.		Ten-acre, Plot 2 nd , Crop 1.		Ten-acre, Plot 3 rd , Crop 1.		Ten-acre, Plot 4 th , Crop 1.				
Quantities weighed (cons. cwt. grs. lbs.)		0 0 3 4		0 6 0 5		0 9 3 17		0 8 3 2				
Cows. Nos.	Breed.	Year old.	Dates of Calving.	Weights of (May 25) lbs.	A.M.		P.M.		A.M.		P.M.	
					Yield of milk, &c.		Weights of (May 25)		Ten-acre, Plot 1 st , Crop 1.		Ten-acre, Plot 2 nd , Crop 1.	
1	Cross short-horn	Aged	May 1	1,081	19 13	13 0	19 0	13 12	19 0	13 12	19 0	13 12
2	Cross short-horn	6	Feb. 15	1,077	13 0	13 0	13 0	13 12	13 0	13 12	13 0	13 12
3	Cross short-horn	8	Feb. 15	1,077	13 0	13 0	13 0	13 12	13 0	13 12	13 0	13 12
4	Cross short-horn	8	Mar. 20	1,084	24 6	19 1	24 0	23 4	17 9	23 4	17 9	23 4
5	Cross short-horn	7	April 17	1,110	19 8	13 4	19 8	23 4	17 9	23 4	17 9	23 4
Totals				5,364	91 2	64 6	97 8	63 12	92 1	59 15	87 3	60 6
Means				1,083	18 4	12 14	19 8	13 9	18 6	13 0	17 7	12 1
											</	

TEN COWS.—SEWAGED MEADOW GRASS.

Grass con- sumed.	From which Field, Plot, and Crop										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten-acre, Plot 2, Crop 1.										Ten									
-------------------------	----------------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	---------------------------	--	--	--	--	--	--	--	--	--	-----	--	--	--	--	--	--	--	--	--

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

From which Field, Plot, and Crop		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.		Rye, Plot 3, Crop 2.	
Quantities weighed (tons, cwt., grs., lbs.)		0 9 3 23		0 6 0 15		0 6 3 3		0 11 0 10		0 7 2 10		0 8 1 23		0 8 1 26		2 18 2 9	
1	Cross short-born Aged April 30	98	10	18	1	15	5	26	0	18	5	25	4	16	14	23	42
2	Cross short-born Aged April 13	102	10	30	3	26	3	16	11	10	10	15	10	19	6	10	295
3	Cross short-born Aged Mar. 1	1,102	17	0	10	31	16	14	15	11	13	15	14	11	3	10	314
4	Cross short-born Aged April 15	1,176	23	2	16	3	22	8	14	12	22	0	12	22	5	8	318
5	Cross short-born Aged Jan. 2	1,048	13	1	9	8	14	10	6	12	23	9	14	21	12	23	437
Totals		97	13	67	10	101	5	61	10	97	3	65	11	59	13	59	1,106

Table X.—continued.

FIVE COWS.—UNSEWAGED MEADOW GRASS.																			
From which Field, Plot, and Crop		Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.		Per head per day.	
Quantities weighed (tons, cwt., qrs. lbs.)		Five-acre, Plot 0, Crop 1.		Five-acre, Plot 0, Crop 1.		Five-acre, Plot 0, Crop 1.		Five-acre, Plot 0, Crop 1.		Five-acre, Plot 0, Crop 1.		Five-acre, Plot 0, Crop 1.		Five-acre, Plot 0, Crop 1.		Five-acre, Plot 0, Crop 1.			
Cow No.	Breed.	Years old.	Dates of Calving.	Wt. (June 23, lbs.)	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
1	Cross short-born	Aged	May 1	1,078	19 7	12 4	20 8	12 13	20 4	12 11	10 0	12 11	20 4	12 11	10 0	12 11	20 4	12 11	10 0
2	Cross short-born	Aged	Feb. 15	1,100	19 7	12 4	20 8	12 13	20 4	12 11	10 0	12 11	20 4	12 11	10 0	12 11	20 4	12 11	10 0
3	Cross short-born	Aged	April 13	1,804	19 7	12 4	20 8	12 13	20 4	12 11	10 0	12 11	20 4	12 11	10 0	12 11	20 4	12 11	10 0
4	Cross short-born	Aged	May 17	1,407	19 7	12 4	20 8	12 13	20 4	12 11	10 0	12 11	20 4	12 11	10 0	12 11	20 4	12 11	10 0
5	Cross short-born	Aged	April 17	1,407	19 7	12 4	20 8	12 13	20 4	12 11	10 0	12 11	20 4	12 11	10 0	12 11	20 4	12 11	10 0
Totals				5,438	88	1	57 14	90 1	50 13	90 1	60 0	92 5	59 10	18 7	11 15	17 6	13 2	33 11	90 7
Means				1,048	17 10	11 9	18 0	11 15	18 0	12 0	18 7	11 15	17 6	13 2	33 11	10 12	18 1	11 4	20 9

TEN COWS.—SEWAGED MEADOW GRASS.																			
From which Field, Plot, and Crop		Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.		Per head per day.	
Quantities weighed (tons, cwt., qrs. lbs.)		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.		Five-acre, Plot 2, Crop 1.			
Cow No.	Breed.	Years old.	Dates of Calving.	Wt. (June 23, lbs.)	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
1	Cross short-born	Aged	Mar. 10	1,121	13 4	10 5	15 0	11 4	14 14	11 7	14 7	14 7	14 7	14 7	14 7	14 7	14 7	14 7	14 7
2	Cross short-born	Aged	June 15	1,550	17 0	10 13	18 13	12 0	18 0	12 11	19 5	12 7	18 10	12 8	19 0	12 8	19 0	12 8	19 0
3	Cross short-born	Aged	Feb. 20	1,661	17 0	10 13	18 13	12 0	18 0	12 11	19 5	12 7	18 10	12 8	19 0	12 8	19 0	12 8	19 0
4	Cross short-born	Aged	Feb. 15	1,496	13 6	9 10	14 0	10 0	14 0	10 0	14 0	10 0	14 0	10 0	14 0	10 0	14 0	10 0	14 0
5	Ayrshire	Aged	Feb. 22	1,400	13 6	9 10	14 0	10 0	14 0	10 0	14 0	10 0	14 0	10 0	14 0	10 0	14 0	10 0	14 0
6	Cross short-born	Aged	April 14	1,508	17 0	10 13	18 13	12 0	18 0	12 11	19 5	12 7	18 10	12 8	19 0	12 8	19 0	12 8	19 0
7	Montreal	Aged	April 13	1,604	17 0	10 13	18 13	12 0	18 0	12 11	19 5	12 7	18 10	12 8	19 0	12 8	19 0	12 8	19 0
8	Cross short-born	Aged	April 15	1,604	17 0	10 13	18 13	12 0	18 0	12 11	19 5	12 7	18 10	12 8	19 0	12 8	19 0	12 8	19 0
9	Cross short-born	Aged	April 15	1,604	17 0	10 13	18 13	12 0	18 0	12 11	19 5	12 7	18 10	12 8	19 0	12 8	19 0	12 8	19 0
10	Half short-born	Aged	April 23	1,196	18 4	6 0	6 13	8 12	8 12	6 4	8 10	7 0	8 2	6 8	8 4	6 12	11 0	10 4	14 13
Totals				10,523	130	2	100 11	144 3	107 7	142 3	110 6	144 7	103 12	141 15	107 12	141 9	87 10	134 7	86 8
Means				1,052	13 15	10 1	14 7	10 12	14 3	11 1	14 7	10 6	14 5	10 13	14 3	8 12	13 7	8 10	160 4

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.																			
From which Field, Plot, and Crop		Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.		Per head per day.	
Quantities weighed (tons, cwt., qrs. lbs.)		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.			
Cow No.	Breed.	Years old.	Dates of Calving.	Wt. (June 23, lbs.)	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
1	Cross short-born	Aged	April 20	1,122	22 8	14 6	25 6	17 14	27 0	16 2	23 3	16 10	23 14	17 0	22 10	14 2	21 6	14 11	270 12
2	Ayrshire	Aged	Mar. 15	878	17 11	11 6	18 10	11 10	13 13	11 10	18 10	11 2	17 11	9 4	17 11	8 4	9 14	20 5	80 13
3	Cross short-born	Aged	Mar. 1	1,047	14 5	10 0	15 4	10 0	14 15	11 13	14 15	10 5	14 12	9 5	14 15	8 4	15 8	6 14	24 8
4	Cross short-born	Aged	April 15	1,180	19 5	13 12	20 10	13 0	21 8	12 2	21 2	13 10	20 0	12 14	20 2	14 1	20 1	12 7	253 12
5	Cross short-born	Aged	Jan. 2	1,126	14 2	8 6	13 6	10 3	12 10	8 15	13 8	10 3	13 1	8 6	13 3	8 2	13 8	8 0	102 9
Totals				5,353	88	2	100 11	144 3	107 7	142 3	110 6	144 7	103 12	141 15	107 12	141 9	87 10	134 7	86 8
Means				1,071	18 15	10 1	14 7	10 12	14 3	11 1	14 7	10 6	14 5	10 13	14 3	8 12	13 7	8 10	160 4

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.																			
From which Field, Plot, and Crop		Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.		Sunday.		Monday.		Total in 7 days.		Per head per day.	
Quantities weighed (tons, cwt., qrs. lbs.)		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.		Five-acre, Plot 2, Crop 2.			
Cow No.	Breed.	Years old.	Dates of Calving.	Wt. (June 23, lbs.)	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
1	Cross short-born	Aged	April 20	1,122	22 8	14 6	25 6	17 14	27 0	16 2	23 3	16 10	23 14	17 0	22 10	14 2	21 6	14 11	270 12
2	Ayrshire	Aged	Mar. 15	878	17 11	11 6	18 10	11 10	13 13	11 10	18 10	11 2	17 11	9 4	17 11	8 4	9 14	20 5	80 13
3	Cross short-born	Aged	Mar. 1	1,047	14 5	10 0	15 4	10 0	14 15	11 13	14 15	10 5	14 12	9 5	14 15	8 4	15 8	6 14	24 8
4	Cross short-born	Aged	April 15	1,180	19 5	13 12	20 10	13 0	21 8	12 2	21 2	13 10	20 0	12 14	20 2	14 1	20 1	12 7	253 12
5	Cross short-born	Aged	Jan. 2	1,126	14 2	8 6	13 6	10 3	12 10	8 15	13 8	10 3	13 1	8 6	13 3	8 2	13 8	8 0	102 9
Totals				5,353	88	2	100 11	144 3	107 7	142 3	110 6	144 7	103 12	141 15	107 12	141 9	87 10	134 7	86 8
Means				1,071	18 15	10 1	14 7	10 12	14 3	11 1	14 7	10 6	14 5	10 13	14 3	8 12	13 7	8 10	160 4

FIVE COWS.—UNSEWAGED MEADOW GRASS.

Cows consumed.	From which Field, Plot, and Crop		Quantities weighed (tons, cwt., qrs., lbs.)												Yield of Milk, &c.				
			Ten-acre, Plot 0 th , Crop 1.		Five-acre, Plot 0 th , Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.		Ten-acre, Plot 1, Crop 1.						
Cows. Nos.	Breed.	Years old.	Dates of Calving.	Weights (June 22) lbs.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	Ten-acre, Plot 1, Crop 1.	Ten-acre, Plot 1, Crop 1.	Ten-acre, Plot 1, Crop 1.
1	Cross short-horn	Ag'd	May 1	1,078	10 12 2 22	0 4 0 4	0 2 8 23½	0 4 0 14	0 4 8 5½	0 3 1 23½	0 3 1 28	—	—	—	—	—	—	—	—
2	Cross short-horn	6	Feb. 15	1,100	10 12 2 22	0 4 0 4	0 2 8 23½	0 4 0 14	0 4 8 5½	0 3 1 23½	0 3 1 28	—	—	—	—	—	—	—	—
3	Cross short-horn	8	April 13	1,811	10 12 2 22	0 4 0 4	0 2 8 23½	0 4 0 14	0 4 8 5½	0 3 1 23½	0 3 1 28	—	—	—	—	—	—	—	—
4	Cross short-horn	5	April 17	1,467	10 12 2 22	0 4 0 4	0 2 8 23½	0 4 0 14	0 4 8 5½	0 3 1 23½	0 3 1 28	—	—	—	—	—	—	—	—
5	Cross short-horn	7	April 17	1,467	10 12 2 22	0 4 0 4	0 2 8 23½	0 4 0 14	0 4 8 5½	0 3 1 23½	0 3 1 28	—	—	—	—	—	—	—	—
Totals				3,225	50 60 10 110	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100	20 80 12 100
Means				1,048	16 12	11 9	17 7	11 0	17 6	11 7	16 9	11 4	16 6	11 1	17 0	9 13	17	10 5	27 15

TEN COWS.—SEWAGED MEADOW GRASS.

Grass consumed.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)										Yield of milk, &c.								
		Five-acre, Plot 4, Crop 2.		Five-acre, Plot 3, Crop 2.		Five-acre, Plot 3, Crop 2.		Ten-acre, Plot 3, Crop 2.		Ten-acre, Plot 3, Crop 2.										
		Mar. 10	June 15	Mar. 10	June 15	Mar. 10	June 15	Mar. 10	June 15	Mar. 10	June 15									
1	Cross short-horn	1,321	18 4	10 4	15 4	10 8	16 1	10 8	14 0	11 14	12 0	11 8	16 12	15 4	47 5	10	0	1	1	0
2	Cross short-horn	1,221	18 8	9 11	14 3	10 4	16 1	10 8	14 0	11 14	12 0	11 8	16 12	15 4	48 3	10	0	1	1	0
3	Cross short-horn	1,066	13 0	9 11	14 3	10 4	16 1	10 8	14 0	11 14	12 0	11 8	16 12	15 4	47 5	10	0	1	1	0
4	Cross short-horn	906	12 2	8 1	13 2	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0
5	Ayrshire	1,091	12 5	8 3	13 2	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0
6	Cross short-horn	958	12 8	11 10	17 6	10 7	18 4	10 0	17 12	10 6	18 4	11 4	18 7	9 9	16 14	11 0	13 7	11	22	8
7	Moncrel	894	12 3	8 0	13 1	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0	13 4	9 0
8	Cross short-horn	1,068	10 2	11 14	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12
9	Cross short-horn	1,168	10 2	11 14	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12	10 6	11 12
10	Half short-horn	1,196	23 3	16 0	22 0	14 2	22 0	14 2	22 0	14 2	22 0	14 2	22 0	14 2	22 0	14 2	22 0	14 2	22 0	14 2
Totals		10,332	140 1	104 14	158 10	108 14	158 10	108 14	158 10	108 14	158 10	108 14	158 10	108 14	158 10	108 14	158 10	108 14	158 10	108 14
Means		1,033	14 0	10 8	15 4	10 14	15 7	10 5	14 11	30 11	13 5	11 11	16 7	10 7	16 10	10 14	185 2	36	8	8

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

Cows con- sumed.	From which Field, Plot, and Crop	- {		Ives,		Ives,		Ives,		Ives,		Ives,		Ives,		Ives,		Yield of milk, &c.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
		Plot 2, Crop 2.		Plot 2, Crop 2.		Plot 2, Crop 2.		Plot 2, Crop 2.		Plot 2, Crop 2.		Plot 2, Crop 2.		Plot 2, Crop 2.		Plot 2, Crop 2.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
		Ives,	Plot 2, Crop 2.	Ives,	Plot 2, Crop 2.	Ives,	Plot 2, Crop 2.	Ives,	Plot 2, Crop 2.	Ives,	Plot 2, Crop 2.	Ives,	Plot 2, Crop 2.	Ives,	Plot 2, Crop 2.	Ives,	Plot 2, Crop 2.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
Quantities weighed (tons, cwt., qrs., lbs.)		-		-		-		-		-		-		-		-																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
1	Cross short-born	April 20	21	6	16	0	25	12	14	5	22	0	11	6	22	0	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	14	10	1

TABLE A.—Continued.
Record of Food consumed, and Milk yielded, by Cows fed on Unsewaged and Sewaged Meadow Grass, and on Italian Rye Grass (without Oilcake).
Third Season, 1863; 7 days.—June 30 to July 6.

FIVE COWS.—UNSEWAGED MEADOW GRASS.													
Grass consumed.	From which Field, Plot, and Crop	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights (June 22) (lbs.)	Ten-acre, Plot 1, Crop 1.						
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	Per head per day.
Yield of Milk, &c.	Quantities weighed (tons, cwt., qrs., lbs.)	1	Cross short-horn	6	May 15	1,078	0 2 2 21	0 4 1 22	0 6 0 10	0 5 3 10	0 4 1 13	0 4 0 1	1 12 1 4
		2	Cross short-horn	6	Feb. 15	1,058	0 2 2 21	0 4 1 22	0 6 0 10	0 5 3 10	0 4 1 13	0 4 0 1	1 12 1 4
		3	Cross short-horn	8	April 15	1,050	0 2 2 21	0 4 1 22	0 6 0 10	0 5 3 10	0 4 1 13	0 4 0 1	1 12 1 4
		4	Cross short-horn	8	May 20	1,069	0 2 2 21	0 4 1 22	0 6 0 10	0 5 3 10	0 4 1 13	0 4 0 1	1 12 1 4
		5	Cross short-horn	7	April 17	1,097	0 2 2 21	0 4 1 22	0 6 0 10	0 5 3 10	0 4 1 13	0 4 0 1	1 12 1 4
	Totals					5,293	8 4 1 51	8 4 9 51	8 3 0 53	8 2 6 53	8 0 1 50	8 5 3 50	9 46 15
	Means					1,048	16 13 10	16 14 10	16 9 10	16 7 10	17 3 10	17 1 10	189 6
TEN COWS.—SEWAGED MEADOW GRASS.													
Grass consumed.	From which Field, Plot, and Crop	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights (June 22) (lbs.)	Ten-acre, Plot 1, Crop 1.						
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	Per head per day.
Yield of Milk, &c.	Quantities weighed (tons, cwt., qrs., lbs.)	1	Cross short-horn	8	Mar. 10	1,221	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		2	Cross short-horn	8	June 15	1,280	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		3	Cross short-horn	6	Feb. 20	1,056	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		4	Cross short-horn	7	Feb. 15	1,061	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		5	Cross short-horn	8	Feb. 22	1,069	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
	Totals					5,696	8 4 1 51	8 4 9 51	8 3 0 53	8 2 6 53	8 0 1 50	8 5 3 50	9 46 15
	Means					1,139	16 13 10	16 14 10	16 9 10	16 7 10	17 3 10	17 1 10	189 6
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.													
Grass consumed.	From which Field, Plot, and Crop	Cows Nos.	Breed.	Years old.	Dates of Calving.	Weights (June 22) (lbs.)	Ten-acre, Plot 1, Crop 1.						
							A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	Per head per day.
Yield of Milk, &c.	Quantities weighed (tons, cwt., qrs., lbs.)	1	Cross short-horn	8	Mar. 10	1,221	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		2	Cross short-horn	8	June 15	1,280	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		3	Cross short-horn	6	Feb. 20	1,056	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		4	Cross short-horn	7	Feb. 15	1,061	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
		5	Cross short-horn	8	Feb. 22	1,069	0 17 3 18	0 18 5 27	0 12 0 0	0 15 1 21	0 12 2 12	0 12 3 18	5 6 2 7
	Totals					5,696	8 4 1 51	8 4 9 51	8 3 0 53	8 2 6 53	8 0 1 50	8 5 3 50	9 46 15
	Means					1,139	16 13 10	16 14 10	16 9 10	16 7 10	17 3 10	17 1 10	189 6

FIVE COWS.—UNSEWAGED MEADOW GRASS.

Grass consumed.	From which Field, Plot, and Crop			Weights (June 22)	Quantities weighed (tons, cwt., qrs., lbs.)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
	Cows Nos.	Breed.	Years old.		Dates of Calving.	Ten-acre, Plot 1, Crop 1.			Ten-acre, Plot 1, Crop 1.			Ten-acre, Plot 1, Crop 1.			Five-acre, Plot 1, Crop 1.			Five-acre, Plot 1, Crop 1.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																					
						A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.	A.M.	P.M.	F.M.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			
Field of Milk, &c.	1	Cross short-horn	Aged	May 1	1,678																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		

TEN COWS.—SEWAGED MEADOW GRASS.

[illegible]

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

[illegible]

Table X.—continued.
Detailed Record of Food consumed, and Milk yielded, by Cows fed on Unsewaged and Sewaged Meadow Grass, and on Italian Rye Grass, with Oilcake in addition.
Third season, 1903; 7 days.—September 28 to September 28.

		FIVE COWS.—UNSEWAGED MEADOW GRASS.										Total in 7 days.		Yer head per day.	
Food consumed.	Grass	From which Field, Plot, and Crop		Five-acre, Plot 0 th , Crop 2.		Five-acre, Plot 1, Crop 2.		Five-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.		Ten-acre, Plot 1, Crop 2.	
		Quantities weighed (tons, cwt., qrs., lbs.)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)	
		Cows Nos.	Years old.	Dates of Calving.	Weights (Sept. 14) lbs.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	Aged	May 1	1,294	20 8	12 12	21 8	13 13	20 12	11 13	21 6	13 10	20 8	12 14
	2	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
	3	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
	4	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
	5	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
Yield of Milk, &c.	1	Cross short-horn	Aged	May 1	1,294	20 8	12 12	21 8	13 13	20 12	11 13	21 6	13 10	20 8	12 14
	2	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
	3	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
	4	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
	5	Cross short-horn	2	Feb. 15	1,272	11 0	6 11	11 0	13 10	20 12	11 13	21 6	13 10	20 8	12 14
		Totals			5,270	84 5	53 2	87 2	53 8	85 1	49 7	87 3	53 11	84 9	54 7
		Means			1,141	16 14	10 10	17 7	10 11	17 0	9 14	17 7	10 12	16 15	10 14
TEN COWS.—SEWAGED MEADOW GRASS.															
Food consumed.	Grass	From which Field, Plot, and Crop		Five-acre, Plot 0 th , Crop 4.		Five-acre, Plot 1, Crop 4.		Five-acre, Plot 1, Crop 4.		Ten-acre, Plot 1, Crop 4.		Ten-acre, Plot 1, Crop 4.		Ten-acre, Plot 1, Crop 4.	
		Quantities weighed (tons, cwt., qrs., lbs.)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)	
		Cows Nos.	Years old.	Dates of Calving.	Weights (Sept. 14) lbs.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	Aged	Mar. 10	1,298	15 0	10 3	14 12	10 12	14 8	10 13	14 2	9 10	15 1	10 7
	2	Cross short-horn	8	June 15	1,294	16 10	11 8	17 12	11 11	18 6	11 11	17 11	11 6	18 5	11 8
	3	Cross short-horn	8	June 15	1,294	12 1	8 8	12 0	8 6	12 1	8 13	11 12	8 15	11 8	8 10
	4	Cross short-horn	8	June 15	1,294	12 1	8 8	12 0	8 6	12 1	8 13	11 12	8 15	11 8	8 10
	5	Cross short-horn	8	June 15	1,294	12 1	8 8	12 0	8 6	12 1	8 13	11 12	8 15	11 8	8 10
Yield of Milk, &c.	1	Cross short-horn	Aged	Mar. 10	1,298	15 0	10 3	14 12	10 12	14 8	10 13	14 2	9 10	15 1	10 7
	2	Cross short-horn	8	June 15	1,294	16 10	11 8	17 12	11 11	18 6	11 11	17 11	11 6	18 5	11 8
	3	Cross short-horn	8	June 15	1,294	12 1	8 8	12 0	8 6	12 1	8 13	11 12	8 15	11 8	8 10
	4	Cross short-horn	8	June 15	1,294	12 1	8 8	12 0	8 6	12 1	8 13	11 12	8 15	11 8	8 10
	5	Cross short-horn	8	June 15	1,294	12 1	8 8	12 0	8 6	12 1	8 13	11 12	8 15	11 8	8 10
		Totals			11,444	144 0	94 8	146 10	90 5	144 10	97 0	147 5	90 5	145 1	99 6
		Means			1,144	14 6	9 7	14 11	9 15	14 7	9 11	14 12	9 15	14 8	9 15
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.															
Food consumed.	Grass	From which Field, Plot, and Crop		Five-acre, Plot 0 th , Crop 4.		Five-acre, Plot 1, Crop 4.		Five-acre, Plot 1, Crop 4.		Ten-acre, Plot 1, Crop 4.		Ten-acre, Plot 1, Crop 4.		Ten-acre, Plot 1, Crop 4.	
		Quantities weighed (tons, cwt., qrs., lbs.)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)		Oilkake (3 parts cotton & 2 parts rape-cake) (ditto)	
		Cows Nos.	Years old.	Dates of Calving.	Weights (Sept. 14) lbs.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.
Yield of Milk, &c.	1	Cross short-horn	Aged	Apr. 30	1,121	20 5	12 0	20 0	12 4	19 14	13 1	19 10	12 0	18 6	11 4
	2	Cross short-horn	8	Mar. 15	1,094	17 0	11 15	17 10	10 13	17 10	11 15	16 2	10 11	16 2	10 11
	3	Cross short-horn	8	Mar. 15	1,094	17 0	11 15	17 10	10 13	17 10	11 15	16 2	10 11	16 2	10 11
	4	Cross short-horn	8	Mar. 15	1,094	17 0	11 15	17 10	10 13	17 10	11 15	16 2	10 11	16 2	10 11
	5	Cross short-horn	8	Mar. 15	1,094	17 0	11 15	17 10	10 13	17 10	11 15	16 2	10 11	16 2	10 11
		Totals			5,600	84 5	53 2	87 2	53 8	85 1	49 7	87 3	53 11	84 9	54 7
		Means			1,120	16 9	10 3	17 4	10 6	17 0	9 17	17 0	10 12	16 18	10 14

[illegible]

FIVE COWS.—UNSEWAGED MEADOW GRASS, *a*

[illegible]

TEN COWS.—SEWAGED MEADOW GRASS.

Food consumed.	Grass	From which Field, Plot, and Crop		—		Five-acre, Plot 4, Crop 3.		Five-acre, Plot 4, Crop 3.		Ten-acre, Plot 4, Crop 3.		Five-acre, Plot 3, Crop 3.		Five-acre, Plot 3, Crop 3.		—			
		Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)	Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)	Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)	Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)	Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)	Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)	Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)	Quantities weighed (tons, cwt., lbs.)	Oileate (cotton-cake)		
Yield of Milk, &c.	1	Cross short-horn	Aged	Mar. 10	1,154	17 10	12 12	17 14	13 9	18 3	12 10	17 12	12 9	17 0	13 14	17 6	11 4	17 8	212 12
	2	Cross short-horn	6	June 15	1,584	22 11	15 8	22 3	16 0	21 12	15 11	21 0	15 10	22 4	15 3	21 8	13 0	21 12	306 6
	3	Cross short-horn	8	Feb. 20	1,146	15 10	10 6	14 3	11 5	12 14	11 0	13 8	10 9	13 12	10 1	13 10	9 2	14 7	258 5
	4	Cross short-horn	8	Feb. 15	1,040	15 11	10 6	14 3	11 5	12 14	11 0	13 8	10 9	13 12	10 1	13 10	9 2	14 7	253 13
	5	Cross short-horn	8	Feb. 15	1,040	15 11	10 6	14 3	11 5	12 14	11 0	13 8	10 9	13 12	10 1	13 10	9 2	14 7	253 13
	6	Cross short-horn	8	Feb. 15	1,040	15 11	10 6	14 3	11 5	12 14	11 0	13 8	10 9	13 12	10 1	13 10	9 2	14 7	253 13
	7	Cross short-horn	8	Feb. 15	1,040	15 11	10 6	14 3	11 5	12 14	11 0	13 8	10 9	13 12	10 1	13 10	9 2	14 7	253 13
	8	Cross short-horn	8	Feb. 15	1,040	15 11	10 6	14 3	11 5	12 14	11 0	13 8	10 9	13 12	10 1	13 10	9 2	14 7	253 13
	9	Cross short-horn	8	Feb. 15	1,040	15 11	10 6	14 3	11 5	12 14	11 0	13 8	10 9	13 12	10 1	13 10	9 2	14 7	253 13
	10	Half short-horn	8	April 25	1,226	22 8	15 12	21 3	16 3	25 6	15 0	23 15	15 0	23 12	14 15	23 0	14 12	22 4	386 10
		Totals			11,035	109 13	110 15	168 2	129 4	171 5	117 4	169 1	117 7	168 7	118 0	170 6	168 2	1,995 7	
		Means			1,106	17 0	11 11	16 13	12 0	17 2	11 12	16 14	11 12	16 13	11 13	17 1	16 5	169 9	

FIVE COWS.—SEWAGED OR UNSEWAGED ITALIAN RYE GRASS.

Grass	From which Field, Plot, and Crop		Yrs.,		Yrs.,		Yrs.,		Yrs.,		Yrs.,		Yrs.,		Yrs.,		Yrs.,	
	Plot 3, Crop 4.		Plot 3, Crop 4.		Plot 3, Crop 4.		Plot 3, Crop 4.		Plot 3, Crop 4.		Plot 3, Crop 4.		Plot 3, Crop 4.		Plot 3, Crop 4.		Plot 3, Crop 4.	
Quantities weighed (tons, cwt., qrs., lbs.)	Oleate (centin-ease)		Oleate (centin-ease)		Oleate (centin-ease)		Oleate (centin-ease)		Oleate (centin-ease)		Oleate (centin-ease)		Oleate (centin-ease)		Oleate (centin-ease)		Oleate (centin-ease)	
1	1,140	April 20	21	8	15	6	22	10	14	8	23	2	12	10	24	0	12	8
2	910	Mar. 15	16	11	9	14	18	3	10	14	18	3	10	14	18	3	10	14
3	1,170	Mar. 1	15	12	10	17	16	10	10	16	1	14	1	14	1	14	1	14
4	1,218	April 15	21	2	11	12	15	13	13	12	14	13	12	14	13	12	14	13
5	1,184	Jan. 2	21	2	0	0	9	0	4	11	6	7	11	6	7	11	6	7
Total	5,682		87	8	58	2	87	0	56	7	89	1	53	2	88	11	50	12

Table X.—continued.
Detailed Record of Food consumed, and Milk yielded, by Cows fed on Unswaged and Sewaged Meadow Grass, and on Italian Rye Grass, with Oilcake in addition.
 Third Season, 1898; 7 days.—August 11 to August 17.

FIVE COWS.—UNSEWAGED MEADOW GRASS, a																	
Food con- sumed.	Grass	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Dates of Calving	Breed	Plot 1, Crop 3.		Rye, Crop 3.		Plot 1, Crop 3.		Rye, Crop 3.		Sunday.	Monday.	Total in 7 days.	Per head per day.
						A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.				
Yield of Milk, &c.	1	Cross short-horn	1,170	Aug. 17	Weighted	0 2 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10
	2	Cross short-horn	1,170	May 12	Weighted	0 2 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10
	3	Cross short-horn	1,170	May 12	Weighted	0 2 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10
	4	Cross short-horn	1,170	May 20	Weighted	0 2 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10
	5	Cross short-horn	1,170	April 17	Weighted	0 2 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10	0 3 10
	Totals						9 12 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59
TEX COWS.—SEWAGED MEADOW GRASS.																	
Food con- sumed.	1	Cross short-horn	1,184	Mar. 10	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	2	Cross short-horn	1,184	June 8	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	3	Cross short-horn	1,184	Feb. 15	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	4	Cross short-horn	1,184	Feb. 22	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	5	Cross short-horn	1,184	April 12	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	Totals						9 12 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59
Yield of Milk, &c.	1	Cross short-horn	1,184	Mar. 10	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	2	Cross short-horn	1,184	June 8	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	3	Cross short-horn	1,184	Feb. 15	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	4	Cross short-horn	1,184	Feb. 22	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	5	Cross short-horn	1,184	April 12	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	Totals						9 12 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59
Food con- sumed.	1	Cross short-horn	1,184	Mar. 10	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	2	Cross short-horn	1,184	June 8	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	3	Cross short-horn	1,184	Feb. 15	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	4	Cross short-horn	1,184	Feb. 22	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	5	Cross short-horn	1,184	April 12	Weighted	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12	0 1 3 12
	Totals						9 12 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59	9 11 59

FIVE COWS.—UNRWAQED MEADOW (RAB. 7

Food consumed.	Grass	From which Field, Plot, and Crop		Quantities weighed (tons, cwt, qrs. lbs.)												Oleate (cotton-seed)											
		Breed.	Years old.	Intakes of Calving.		Rye, Crop 2.		Rye, Crop 3.		Rye, Crop 4.		Rye, Crop 5.		Rye, Crop 6.		Rye, Crop 7.		Rye, Crop 8.		Rye, Crop 9.		Rye, Crop 10.					
				A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		
1	Cross short-horn	1	1876	May 1	12.0	13.4	12.0	13.4	12.0	13.4	12.0	13.4	12.0	13.4	12.0	13.4	12.0	13.4	12.0	13.4	12.0	13.4	12.0	13.4			
2	Cross short-horn	2	1258	Feb. 15	14.0	8.6	14.0	8.6	14.0	8.6	14.0	8.6	14.0	8.6	14.0	8.6	14.0	8.6	14.0	8.6	14.0	8.6	14.0	8.6			
3	Cross short-horn	3	878	April 13	13.12	9.7	16.0	10.3	13.6	9.6	16.0	10.3	13.6	9.6	16.0	10.3	13.6	9.6	16.0	10.3	13.6	9.6	16.0	10.3			
4	Cross short-horn	4	1116	May 30	24.6	16.6	23.5	15.7	28.6	16.5	28.6	16.5	28.6	16.5	28.6	16.5	28.6	16.5	28.6	16.5	28.6	16.5	28.6	16.5			
5	Cross short-horn	5	1194	April 17	18.0	11.1	18.6	11.0	18.3	11.0	18.6	11.0	18.3	11.0	18.6	11.0	18.3	11.0	18.6	11.0	18.3	11.0	18.6	11.0			
				Totals	93.8	58.7	90.3	68.15	96.2	60.0	94.8	78.3	98.0	57.15	96.5	63.9	90.12	64.4	1469	13							
				Means	18.11	11.11	19.4	11.13	19.4	12.0	18.14	11.10	19.3	11.9	19.1	10.11	19.6	10.14	213	15							

TEN COWS.—SEWAGED MEADOW GRASS.

Food consumed.	Yield milk, &c.	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)																
			Grass		Oaten (sown-cake)		Ten-acre, Plot 4, Crop 3.		Ten-acre, Plot 4, Crop 3.		Ten-acre, Plot 4, Crop 3.		Ten-acre, Plot 4, Crop 3.						
1	Cross short horn	Mar. 10	1,182	16 4	11 12	16 0	12 7	17 2	12 3	16 1	12 4	16 12	10 6	17 2	11 0	16 10	185 12	28 0	
2	Cross short horn	June 15	1,284	21 7	13 1	20 8	13 4	20 8	13 7	20 4	13 6	20 1	12 1	13 14	21 0	13 4	224 2	35 7	
3	Cross short horn	Feb. 11	1,283	15 12	10 2	12 0	13 0	13 0	13 0	13 0	13 0	13 0	13 0	13 0	13 0	13 0	137 2	22 7	
4	Cross short horn	Feb. 22	1,109	11 8	9 11	12 10	8 7	12 5	9 2	12 4	10 13	13 4	10 12	12 6	9 7	12 1	147 12	31 1	
5	Avyshire	Feb. 22	1,109	11 8	9 11	12 10	8 7	12 5	9 2	12 4	10 13	13 4	10 12	12 6	9 7	12 1	147 12	31 1	
6	Cross short horn	Mar. 14	1,048	19 0	13 5	18 11	11 5	17 12	13 1	19 0	11 0	13 0	12 0	18 10	19 1	10 7	276 8	33 13	
7	Monrel	Mar. 14	684	16 1	9 1	15 11	8 8	14 14	10 4	13 4	8 14	13 4	8 16	14 14	8 1	15 4	7 0	108 8	24 1
8	Cross short horn	Apr. 16	1,140	11 12	9 8	10 0	9 8	12 4	9 4	12 0	9 8	12 0	9 8	13 4	9 0	13 2	6 0	160 5	21 6
9	Cross short horn	Apr. 13	1,240	20 3	11 10	18 1	13 12	12 0	12 8	17 4	12 10	16 14	13 5	14 6	11 13	13 8	211 10	30 4	
10	Half short horn	Apr. 25	1,090	21 4	16 4	20 1	15 1	21 2	14 12	21 2	14 12	21 8	13 4	20 12	12 10	21 15	247 11	33 6	
	Totals		11,350	165 3	111 4	160 11	111 6	161 9	110 13	161 7	110 9	160 8	107 8	155 4	103 4	161 15	1,896 1	—	
	Means		1,421	16 8	11 2	16 1	11 2	16 3	11 1	16 2	11 1	16 1	10 12	13 8	10 9	16 2	186 10	36 15	

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

Food con- sumed.	From which Field, Plot, and Crop	Grams	Quantities weighed (tons, cwt., qrs., lbs.)		Rye,		Kre,		Rye,		Rye,		Rye,		—					
			Osteke (cont'n-eake)	—	Plot of, Crop 3.	Plot of, Crop 3.	Plot of, Crop 3.	Plot of, Crop 3.	Plot of, Crop 3.	Plot of, Crop 3.	Plot of, Crop 3.	Plot of, Crop 3.								
1	Cross short born	Aged	April 20	1,180	21 12	13 7	22 11	12 11	21 12	13 5	21 2	13 4	20 12	15 5	21 8	12 6	23 7	10 12	24 2	34 14
2	Cross short born	Aged	May 15	1,222	15 13	8 24	15 15	9 9	13 12	8 24	15 12	9 9	18 4	9 8	15 12	10 11	17 10	16 1	28 7	28 7
3	Cross short born	Aged	May 1	1,270	19 14	11 12	19 19	12 13	13 6	19 6	10 9	9 7	18 7	11 11	20 10	11 4	20 10	11 43	229 4	31 12
4	Cross short born	Aged	April 15	1,185	14 0	8 1	13 18	8 9	13 13	8 13	13 6	8 8	12 0	8 11	13 4	8 2	15 10	8 6	143 10	31 11
5	Cross short born	Aged	Jan. 2	1,185	14 0	8 1	13 18	8 9	13 13	8 13	13 6	8 8	12 0	8 11	13 4	8 2	15 10	8 6	143 10	31 11
	Totals			5,684	89 1	54 2	90 4	53 12	89 1	55 0	87 10	83 15	85 7	65 4	88 9	61 13	92 1	49 10	999 9	—

Table X.—continued.
Detailed Record of Food consumed, and Milk yielded, by Cows fed on Unsewaged and Sewaged Meadow Grass, and on Italian Rye Grass, with Oilcake in addition.
 Third Season, 1883; 7 days.—August 25 to August 31.

		Tuesday.	Wednesday.	Thursday.	Friday.	Saturday.	Sunday.	Monday.	Total in 7 days.	Per head per day.
FIVE COWS.—UNSEWAGED MEADOW GRASS, &										
Food consumed.	Grass { Oilcake (cotton-seed)	From which Field, Plot, and Crop		Rye, * Crop 3.		Rye, * Crop 3.		Rye, * Crop 3.		Total in 7 days.
		Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	
Cows, Nos.	Breed.	Weights (Aug. 17)		A.M. P.M.		A.M. P.M.		A.M. P.M.		Total in 7 days.
		Calving.	Aug. 17	Calving.	Aug. 17	Calving.	Aug. 17	Calving.	Aug. 17	
1	Cross short-horn	May 1	1,170							
2	Cross short-horn	Feb. 15	1,253							
3	Cross short-horn	April 13	878							
4	Cross short-horn	May 20	1,118							
5	Cross short-horn	April 17	1,194							
Totals			5,619							
Means			1,124							
TEN COWS.—SEWAGED MEADOW GRASS.										
Food consumed.	Grass { Oilcake (cotton-seed)	From which Field, Plot, and Crop		Ten acre, Plot 3, Crop 3.		Ten acre, Plot 3, Crop 3.		Ten acre, Plot 3, Crop 3.		Total in 7 days.
		Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	
Cows, Nos.	Breed.	Weights (Aug. 17)		A.M. P.M.		A.M. P.M.		A.M. P.M.		Total in 7 days.
		Calving.	Aug. 17	Calving.	Aug. 17	Calving.	Aug. 17	Calving.	Aug. 17	
1	Cross short-horn	Mar. 10	1,182							
2	Cross short-horn	June 15	1,384							
3	Cross short-horn	Feb. 15	1,170							
4	Cross short-horn	Feb. 15	1,070							
5	Cross short-horn	Feb. 22	1,019							
6	Cross short-horn	April 14	1,048							
7	Cross short-horn	April 12	884							
8	Cross short-horn	April 16	1,140							
9	Cross short-horn	April 13	1,040							
10	Cross short-horn	April 25	1,260							
Totals			11,209							
Means			1,121							
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.										
Food consumed.	Grass { Oilcake (cotton-seed)	From which Field, Plot, and Crop		Rye, * Crop 4.		Rye, * Crop 4.		Rye, * Crop 4.		Total in 7 days.
		Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	Quantities weighed (tons, cwt., lbs.)	(ditto)	
Cows, Nos.	Breed.	Weights (Aug. 17)		A.M. P.M.		A.M. P.M.		A.M. P.M.		Total in 7 days.
		Calving.	Aug. 17	Calving.	Aug. 17	Calving.	Aug. 17	Calving.	Aug. 17	
1	Cross short-horn	Mar. 10	1,182							
2	Cross short-horn	June 15	1,384							
3	Cross short-horn	Feb. 15	1,170							
4	Cross short-horn	Feb. 15	1,070							
5	Cross short-horn	Feb. 22	1,019							
6	Cross short-horn	April 14	1,048							
7	Cross short-horn	April 12	884							
8	Cross short-horn	April 16	1,140							
9	Cross short-horn	April 13	1,040							
10	Cross short-horn	April 25	1,260							
Totals			11,209							
Means			1,121							

FIVE COWS.—UNSEWAGED MEADOW GRASS, a											
Food consumed.	Grass	From which Field, Plot, and Crop		Rye, Crop 3 rd .		Rye, Crop 3 rd .		Rye, Crop 3 rd .		—	
				Quantities weighed (tons, cwt, qrs, lbs.)		Quantities weighed (tons, cwt, qrs, lbs.)		Quantities weighed (tons, cwt, qrs, lbs.)			
Yield of Milk, &c.	Cows Nos.	Breed.	Years old.	Dates of Calving.		Rye, Crop 3 rd .		Rye, Crop 3 rd .		—	
				(Aug. 17)		A.M. P.M.		A.M. P.M.			
	1	Cross short-horn	Aged	May 1	1,176	0 4 3 0	0 2 0 13	0 2 2 22	0 2 3 12	1 2 3 21	
	2	Cross short-horn	6	Feb. 15	1,253	0 0 0 15	0 0 0 15	0 0 0 15	0 0 0 15	0 0 0 3	
	3	Cross short-horn	8	May 20	878	14 2 9	14 14 8	8 15 12 9	10 10 15	0 1 3 21	
	4	Cross short-horn	8	April 13	1,118	22 13 14	20 25 2	10 10 22	3 12 12	0 0 0 3	
	5	Cross short-horn	7	April 17	1,194	16 12 10	16 4 9	9 9 3	16 15 12	0 0 0 3	
	Totals		5,619								
	Means		1,124								
TEX COWS.—SEWAGED MEADOW GRASS.											
Food consumed.	Grass	From which Field, Plot, and Crop		Five-acre, Plot 4, Crop 4.		Ten-acre, Plot 4, Crop 4.		Five-acre, Plot 3, Crop 3.		—	
				Quantities weighed (tons, cwt, qrs, lbs.)		Quantities weighed (tons, cwt, qrs, lbs.)		Quantities weighed (tons, cwt, qrs, lbs.)			
Yield of Milk, &c.	Cows Nos.	Breed.	Years old.	Dates of Calving.		Rye, Crop 3 rd .		Rye, Crop 3 rd .		—	
				(Aug. 17)		A.M. P.M.		A.M. P.M.			
	1	Cross short-horn	Aged	Mar. 10	1,182	0 0 0 10	0 18 1 14	0 12 1 20	0 13 3 9	5 12 1 24	
	2	Cross short-horn	6	Feb. 13	1,182	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	3	Cross short-horn	7	Feb. 13	1,182	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	4	Cross short-horn	7	Feb. 13	1,182	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	5	Cross short-horn	8	Feb. 22	1,019	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	6	Cross short-horn	8	Feb. 22	1,019	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	Totals		5,619								
Means		1,124									
FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.											
Food consumed.	Grass	From which Field, Plot, and Crop		Rye, Plot 3, Crop 5.		Rye, Plot 3, Crop 5.		Rye, Plot 3, Crop 5.		—	
				Quantities weighed (tons, cwt, qrs, lbs.)		Quantities weighed (tons, cwt, qrs, lbs.)		Quantities weighed (tons, cwt, qrs, lbs.)			
Yield of Milk, &c.	Cows Nos.	Breed.	Years old.	Dates of Calving.		Rye, Plot 3, Crop 5.		Rye, Plot 3, Crop 5.		—	
				(Aug. 17)		A.M. P.M.		A.M. P.M.			
	1	Cross short-horn	Aged	Mar. 10	1,182	0 0 0 10	0 18 1 14	0 12 1 20	0 13 3 9	5 12 1 24	
	2	Cross short-horn	6	Feb. 13	1,182	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	3	Cross short-horn	7	Feb. 13	1,182	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	4	Cross short-horn	7	Feb. 13	1,182	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	5	Cross short-horn	8	Feb. 22	1,019	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	6	Cross short-horn	8	Feb. 22	1,019	0 0 1 2	0 0 1 2	0 12 1 20	0 13 3 9	0 1 2 12	
	Totals		5,619								
Means		1,124									

Food consumed.		FIVE COWS.—UNSEWAGED MEADOW GRASS. a														Total in 7 days.		Per head per day.	
Grass	Oleate (cotton-seed)	From which Field, Plot, and Crop		Breed.	Years old.	Dates of Calving.	Weights (tons, cwt., qrs., lbs.)	Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.	Sunday.	Monday.	
		Grass	Oleate					Five-acre, Plot 2, Crop 2.	Five-acre, Plot 2, Crop 2.	Five-acre, Plot 2, Crop 2.	Five-acre, Plot 2, Crop 2.	Five-acre, Plot 2, Crop 2.	Five-acre, Plot 2, Crop 2.	Five-acre, Plot 2, Crop 2.	Five-acre, Plot 2, Crop 2.				
1	2	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
2	3	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
3	4	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
4	5	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
5	6	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
Totals		5,772	83 10 56 5	37 4	53 9	88 2	33 10	57 7	11 9	87 15	22 5	85 1	51 10	30 4	53 14	964 9	—	—	
Means		1,154	16 12 11 4	17 5	10 12	17 10	11 2	17 8	10 5	17 9	10 7	17 0	10 3	18 1	10 13	193 15	28 2	—	

Food consumed.		TEN COWS.—SEWAGED MEADOW GRASS.														Total in 7 days.		Per head per day.	
Grass	Oleate (cotton-seed)	From which Field, Plot, and Crop		Breed.	Years old.	Dates of Calving.	Weights (tons, cwt., qrs., lbs.)	Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.	Sunday.	Monday.	
		Grass	Oleate					Five-acre, Plot 3, Crop 3.	Five-acre, Plot 4, Crop 4.	Five-acre, Plot 3, Crop 3.	Five-acre, Plot 4, Crop 4.	Five-acre, Plot 3, Crop 3.	Five-acre, Plot 4, Crop 4.	Five-acre, Plot 3, Crop 3.	Five-acre, Plot 4, Crop 4.				Five-acre, Plot 3, Crop 3.
1	2	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
2	3	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
3	4	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
4	5	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
5	6	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
6	7	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
7	8	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
8	9	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
9	10	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
Totals		11,444	144 10 57 0	144 10	57 0	144 12	102 6	130 10	57 7	140 1	88 7	130 8	95 13	144 3	93 10	134 6	90 14	1703 13	
Means		1,434	14 7 9 11	14 8	10 4	15 1	9 12	14 14	8 14	15 1	9 9	14 7	9 6	15 7	9 1	170 6	34 5	—	

Food consumed.		FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.														Total in 7 days.		Per head per day.	
Grass	Oleate (cotton-seed)	From which Field, Plot, and Crop		Breed.	Years old.	Dates of Calving.	Weights (tons, cwt., qrs., lbs.)	Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.	Sunday.	Monday.	
		Grass	Oleate					Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.				
1	2	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
2	3	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
3	4	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
4	5	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
5	6	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
6	7	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
7	8	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
8	9	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
9	10	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
Totals		11,444	144 10 57 0	144 10	57 0	144 12	102 6	130 10	57 7	140 1	88 7	130 8	95 13	144 3	93 10	134 6	90 14	1703 13	
Means		1,434	14 7 9 11	14 8	10 4	15 1	9 12	14 14	8 14	15 1	9 9	14 7	9 6	15 7	9 1	170 6	34 5	—	

Food consumed.		FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.														Total in 7 days.		Per head per day.	
Grass	Oleate (cotton-seed)	From which Field, Plot, and Crop		Breed.	Years old.	Dates of Calving.	Weights (tons, cwt., qrs., lbs.)	Tuesday.		Wednesday.		Thursday.		Friday.		Saturday.	Sunday.	Monday.	
		Grass	Oleate					Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.	Five-acre, Plot 1, Crop 4.				
1	2	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
2	3	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
3	4	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
4	5	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
5	6	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
6	7	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
7	8	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
8	9	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
9	10	From which Field, Plot, and Crop	Quantities weighed (tons, cwt., qrs., lbs.)	Grass	Oleate	1897	1897	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	0 5 3 22	
Totals		11,444	144 10 57 0	144 10	57 0	144 12	102 6	130 10	57 7	140 1	88 7	130 8	95 13	144 3	93 10	134 6	90 14	1703 13	
Means		1,434	14 7 9 11	14 8	10 4	15 1	9 12	14 14	8 14	15 1	9 9	14 7	9 6	15 7	9 1	170 6	34 5	—	

[illegible]

Table X—continued.

Detailed Record of Food consumed, and Milk yielded, by Cows fed on Unwashed and Sewaged Meadow Grass, and on Italian Rye Grass, with Oilcake in addition.

[illegible]

FIVE COWS—UNSEWAGED MEADOW GRASS.														
Food consumed.	Grass	From which Field, Plot, and Crop			Five-acre, Plot 3, Crop 4.			Five-acre, Plot 4, Crop 4.			Five-acre, Plot 4, Crop 4.			Yield of Milk, &c.
		{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			
Cows.	No.	Breed.	Years old.	Dates of Calving.	Weights (Sept. 14) lbs.	A.M.			P.M.			A.M.		
						Plot 3, Crop 4.			Plot 4, Crop 4.			Plot 4, Crop 4.		
Yield of Milk, &c.	1	Cross short-horn	6	May 1	1,204	21 0	12 13	22 0	21 0	12 13	22 0	21 0	12 13	22 0
	2	Cross short-horn	6	Feb. 15	1,272	10 12	6 4	11 0	10 12	6 4	11 0	10 12	6 4	11 0
	3	Cross short-horn	3	April 18	1,089	16 12	8 13	24 1	16 12	8 13	24 1	16 12	8 13	24 1
	4	Cross short-horn	5	May 30	1,068	17 2	12 14	18 0	17 2	12 14	18 0	17 2	12 14	18 0
	5	Cross short-horn	7	April 17	1,244	17 2	12 14	18 0	17 2	12 14	18 0	17 2	12 14	18 0
Totals					5,707	86 14	53 15	85 13	86 14	53 15	85 13	86 14	53 15	85 13
Means					1,141	17 6	10 13	17 2	17 6	10 13	17 2	17 6	10 13	17 2
TEN COWS—SEWAGED MEADOW GRASS.														
Food consumed.	Grass	From which Field, Plot, and Crop			Five-acre, Plot 3, Crop 4.			Five-acre, Plot 4, Crop 4.			Five-acre, Plot 4, Crop 4.			Yield of Milk, &c.
		{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			
Cows.	No.	Breed.	Years old.	Dates of Calving.	Weights (Sept. 14) lbs.	A.M.			P.M.			A.M.		
						Plot 3, Crop 4.			Plot 4, Crop 4.			Plot 4, Crop 4.		
Yield of Milk, &c.	1	Cross short-horn	8	June 10	1,208	14 11	10 2	15 2	14 11	10 2	15 2	14 11	10 2	15 2
	2	Cross short-horn	8	June 10	1,204	17 12	11 8	17 6	17 12	11 8	17 6	17 12	11 8	17 6
	3	Cross short-horn	6	Feb. 30	1,174	31 0	8 0	11 0	31 0	8 0	11 0	31 0	8 0	11 0
	4	Cross short-horn	7	Feb. 15	1,090	23 4	8 8	13 0	23 4	8 8	13 0	23 4	8 8	13 0
	5	Ayrshire	8	Feb. 22	1,053	10 0	7 14	10 4	10 0	7 14	10 4	10 0	7 14	10 4
Yield of Milk, &c.	6	Cross short-horn	6	April 14	1,068	19 1	11 9	19 12	19 1	11 9	19 12	19 1	11 9	19 12
	7	Concurel	8	April 12	1,190	10 16	4 8	19 0	10 16	4 8	19 0	10 16	4 8	19 0
	8	Cross short-horn	7	April 12	1,087	14 6	11 2	14 10	14 6	11 2	14 10	14 6	11 2	14 10
	9	Cross short-horn	7	April 12	1,087	14 6	11 2	14 10	14 6	11 2	14 10	14 6	11 2	14 10
	10	Half short-horn	8	April 25	1,278	19 13	12 9	19 13	19 13	12 9	19 13	19 13	12 9	19 13
Totals					11,444	142 8	86 1	142 8	142 8	86 1	142 8	142 8	86 1	142 8
Means					1,144	14 3	9 8	14 3	14 3	9 8	14 3	14 3	9 8	14 3
Food consumed.	Grass	From which Field, Plot, and Crop			Five-acre, Plot 3, Crop 4.			Five-acre, Plot 4, Crop 4.			Five-acre, Plot 4, Crop 4.			Yield of Milk, &c.
		{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			{ Quantities weighed (tons, cwt., qrs., lbs.) Oileake (3 parts cotton & 2 parts rape-cake) (ditto)			
Cows.	No.	Breed.	Years old.	Dates of Calving.	Weights (Sept. 14) lbs.	A.M.			P.M.			A.M.		
						Plot 3, Crop 4.			Plot 4, Crop 4.			Plot 4, Crop 4.		
Yield of Milk, &c.	1	Cross short-horn	6	April 30	1,121	17 14	11 6	18 6	17 14	11 6	18 6	17 14	11 6	18 6
	2	Ayrshire	6	Mar. 15	1,354	17 0	10 7	16 13	17 0	10 7	16 13	17 0	10 7	16 13
	3	Cross short-horn	6	Mar. 1	1,068	15 0	9 2	14 10	15 0	9 2	14 10	15 0	9 2	14 10
	4	Cross short-horn	6	April 16	1,068	11 2	7 0	13 0	11 2	7 0	13 0	11 2	7 0	13 0
	5	Cross short-horn	6	Jan. 2	1,254	11 2	7 0	13 0	11 2	7 0	13 0	11 2	7 0	13 0
Totals					5,700	76 1	46 2	79 8	76 1	46 2	79 8	76 1	46 2	79 8

Third Season, 1883; 7 days.—October 6 to October 12.

FIVE COWS.—UNSEWAGED MEADOW GRASS. a

Food consumed.	Grass	From which Field, Plot, and Crop	Cows, Nos.	Breed.	Years old.	Dates of Calving.	Weights (Oct. 12) lbs.	Five-acre, Plot 3, Crop 4.		Ten-acre, Plot 4, Crop 4.		Ten-acre, Plot 3, Crop 4.		Ten-acre, Plot 4, Crop 4.		Five-acre, Plot 3, Crop 4.		Five-acre, Plot 4, Crop 3.		—
								A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	
Yield of Milk, &c.	1	Cross short-born	6	May 1	1,123	19 0 12	19 0 12	0 9 3 20	0 0 0 25	0 5 2 7	0 7 3 9	0 7 3 20	0 11 3 25	0 2 2 6	0 2 2 6	2 8 1 9	0 1 1 15	—		
	2	Cross short-born	6	Feb. 15	1,346	11 5 9	11 5 9	0 9 3 20	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 1 2 7	0 0 0 25			
	3	Cross short-born	3	April 13	1,666	15 5 9	15 5 9	0 9 3 20	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25			
	4	Cross short-born	3	May 25	1,598	22 3 15	22 3 15	0 9 3 20	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25			
	5	Cross short-born	7	April 17	1,598	15 5 9	15 5 9	0 9 3 20	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25	0 0 0 25			
Totals							5,233	81 14 53 15	88 3 52 4	84 12 52 11	90 11 49 7	85 9 53 4	81 4 46 15	84 0 48 8	953 5	—	—			
Means							1,047	16 6 10 13	17 10 10 7	16 15 10 9	18 2 9 14	17 2 10 11	16 4 9 6	16 13 9 11	190 11	27 4				

TEN COWS.—SEWAGED MEADOW GRASS.

Food consumed.	Grass	From which Field, Plot, and Crop	Quantities weighed (cows, cwt, qrs, lbs.)		Ten-acre, Plot 4, Crop 4.		Ten-acre, Plot 3, Crop 4.		Ten-acre, Plot 3, Crop 4.		Five-acre, Plot 2, Crop 4.		Five-acre, Plot 2, Crop 4.		—	
			Oileake (3 parts cotton & 2 parts rape-cake) (ultra)	(ultra)	Five-acre, Plot 3, Crop 4.	Ten-acre, Plot 4, Crop 4.	Ten-acre, Plot 3, Crop 4.	Ten-acre, Plot 4, Crop 4.	Ten-acre, Plot 3, Crop 4.	Ten-acre, Plot 4, Crop 4.	Five-acre, Plot 2, Crop 4.	Five-acre, Plot 2, Crop 4.	—	—		
Yield of milk, &c.	1	Cross short-born	8	Mar. 10	1,190	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15	—	
	2	Cross short-born	8	June 15	1,290	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	3	Cross short-born	6	Feb. 20	1,167	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	4	Cross short-born	6	Feb. 13	1,077	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	5	Cross short-born	6	Feb. 24	1,085	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	6	Cross short-born	6	April 11	1,061	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	7	Cross short-born	6	April 14	1,088	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	8	Cross short-born	6	April 16	1,179	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	9	Cross short-born	6	April 13	1,060	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
	10	Half short-born	8	April 25	1,290	0 10 3 11	0 11 0 13	0 13 9 18	0 13 3 12	0 13 3 12	0 5 0 12	0 5 0 12	4 15 3 15	0 1 1 15		
Totals			11,398	326 8	95 2	143 2	91 0	140 13	95 0	140 13	83 6	143 1	89 6	1,629 2	—	
Means			1,133	32 0	9 8	14 5	9 2	14 1	9 8	14 10	9 6	14 5	9 2	163 15		

FIVE COWS.—UNSEWAGED OR SEWAGED ITALIAN RYE GRASS.

Food consumed.	Grass	From which Field, Plot, and Crop	Rye,		Rye,		Rye,		Rye,		Rye,		Rye,		—																						
			Crop 4.	•	Crop 4.	•	Crop 4.	•	Crop 4.	•	Crop 4.	•	Crop 4.	•																							
Yield of Milk.	1	Cross short-born	17	4	11	5	19	15	12	0	20	1	11	0	19	12	12	8	20	6	11	4	19	15	10	11	19	6	11	33	216	13	81	0			
	2	Ayrshire	17	6	9	13	17	6	11	5	16	18	9	12	0	17	13	9	8	15	10	9	4	13	9	7	17	10	9	7	180	2	37	0			
	3	Cross short-born	14	0	9	8	13	13	9	5	15	4	9	1	14	12	9	8	17	10	9	4	13	9	7	17	10	9	7	181	6	23	1				
	4	Cross short-born	16	6	10	1	11	10	10	1	18	1	11	4	1	14	12	9	8	17	10	9	4	13	9	7	17	10	9	7	203	4	30	4			
	5	Cross short-born	16	0	7	12	11	5	11	5	10	9	8	8	11	8	6	12	11	5	11	8	7	6	10	13	7	7	12	7	0	159	14	28	11		
Totals			17	4	11	5	19	15	12	0	20	1	11	0	19	12	12	8	20	6	11	4	19	15	10	11	19	6	11	33	216	13	81	0			
Means			3	4	2	2	3	9	2	0	4	0	2	2	3	1	2	6	4	1	1	4	3	1	2	2	3	1	2	6	4	1	1	43	2	16	0

Summary of the Analyses of 19 Samples of Sewage-water collected in the Ten-acre Field.

Third Season 1862-3; November 1862—May 1863 inclusive.

GRAINS PER GALLON.														
1862.										1863.				
November			December			January		February		March		April	May	Mean : 13 Samples.
3-5.	17-19.	1	1-3.	15-17.	3	4	5	6	2-4.	7	8	3	17	
Sample Numbers - - - - -														
Organic matter. { In solution - - - - - In suspension - - - - - Total - - - - -														
Inorganic matter. { In solution - - - - - In suspension - - - - - Total - - - - -														
Total in solution - - - - -														
Total in suspension - - - - -														
Total solid matter - - - - -														
Ammonia { In solution - - - - - In suspension - - - - - Total - - - - -														

TABLE XII.

SUMMARY of Food consumed, and Milk and Increase yielded, by Cows fed respectively on Unsewaged and Sewaged Meadow Grass, with Oilcake in addition.

SECOND SEASON, 862.

Periods.*		3 Cows.—Unsewaged Grass.					12 Cows.—Sewaged Grass.				
Dates.*	No. of Days.	Average per head per day.			Average Increase in weight per head per week.	Average per head per day.			Average Increase (or Loss) in weight per head per week.		
		Food consumed.		Milk yielded.		Food consumed.		Milk yielded.			
		Green Grass.	Oil-cake.		Green Grass.	Oil-cake.					
May 2 to May 8 - -	7	lbs. 152	lbs. 3	lbs. ozs. 31 10	10 2½	lbs. 133	lbs. 3	lbs. ozs. 28 4	13 0		
" 9 to " 15 - -	7	158	3	31 4		175	3	26 7			
" 16 to " 22 - -	7	206	3	29 9		147	3	25 2			
" 23 to " 29 - -	7	158	3	29 7	29 10½	154	3	25 8	9 0½		
" 30 to June 5 - -	7	152	3	28 4		143	3	23 11			
June 6 to " 12 - -	7	142	3	26 6		145	3	22 11			
" 13 to " 19 - -	7	160	3	25 8	9 12	89	3	22 2	1 1½		
" 20 to " 26 - -	7	145	3	24 15		114	3	21 1			
" 27 to July 3 - -	7	104	3½	24 0		136	3½	20 14			
July 4 to " 10 - -	7	125	3½	23 8	5 9½	137	3½	21 12	0 11		
" 11 to " 17 - -	7	79	4½	22 13		145	4½	21 7			
" 18 to " 24 - -	7	74	4½	22 2		125	4½	21 2			
" 25 to " 31 - -	7	161	4	23 4	2 8	113	4	21 3	6 2½		
Aug. 1 to Aug. 7 - -	7	86	0	21 2		112	0	21 0			
" 8 to " 14 - -	7	176	4	21 14		125	4	21 1			
" 15 to " 21 - -	7	90	4	22 2	1 8	139	4	20 13	-1 10½		
" 22 to " 28 - -	7	120	4	22 7		153	4	29 1			
" 29 to Sept. 4 - -	7	99	4	22 0		114	4	19 7			
Sept. 5 to " 11 - -	7	140	4	22 0	4 14½	157	4	18 15	4 11½		
" 12 to " 18 - -	7	140	4	21 0		164	4	18 11			
" 19 to " 25 - -	7	133	4	20 0		136	4	17 8			
" 26 to Oct. 2 - -	7	115	5	18 9	7 6	153	5	17 5	138		
Oct. 3 to " 9 - -	7	98	5	17 8		141	5	16 6			
" 10 to " 16 - -	7	117	5	16 5		160	5	16 5			
May 2 to Oct. 16† -	168	130	3½	23 10	7 6	138	3½	21 3	4 11½		

* The dates given are of the "milk yielded," but the periods of the "food consumed" date one day earlier in each case, as also do those of the "average increase (or loss) in weight per head per week."

† It should be observed, that from May 1 to May 15, the first crop of unsewaged meadow grass not being ready to cut, the three cows had Italian rye-grass, and for 20 days, from August 9 to August 28, in default of unsewaged grass, they had green clover; but the figures given in the bottom line of this Table relate to the whole period, irrespectively of these unavoidable irregularities.

TABLE XIII.

SUMMARY of the WEIGHTS, INCREASE (or LOSS), and YIELD of MILK, of the Cows fed respectively on UNSEWAGED MEADOW GRASS, on SEWAGED MEADOW GRASS, and on ITALIAN RYE GRASS (Unsewaged and Sewaged), each for Twelve Weeks alone, and each for Twelve Weeks with OILCAKE in addition.

TABLE XIII.

SUMMARY of the WEIGHTS, INCREASE (or LOSS), and YIELD of MILK, of the Cows fed ITALIAN RYE GRASS (Unsewaged and Sewaged), each for Twelve THIRY

Cows. Nos.	TWELVE WEEKS ON GRASS ALONE.								
	Weights.					Increase (or Loss) in Weight.	Yield of Milk.		
	Apr. 27.	May 25.	Inter- mediate.	June 22.	July 20.		First Week.	Last Week.	Average.
			June 8.						
FIVE COWS.—UNSEWAGE									
1 *	lbs. 1,020	lbs. 1,064	lbs. 1,062	lbs. 1,078	lbs. 1,114	lbs. 94	lbs. oss. 207 13	lbs. oss. 201 14	lbs. oss. 216 13
2	1,117	1,184	1,177	1,190	1,226	109	235 14	148 4	178 9
3	746	818	821	804	830	84	200 10	144 1	169 12
4 †	1,089	1,096	1,094	1,069	1,09	69	178 0	253 12	231 12
5	1,058	1,112	1,110	1,097	1,142	84	230 0	201 4	222 15
Totals -	5,030	5,274	5,264	5,238	5,404	440	1,052 4	919 3	1,019 13
Averages -	1,006	1,055	1,053	1,048	1,081	88	210 7	189 13	203 15
TEN COWS.—SEWAGE									
1	1,064	1,136	..	1,121	1,154	90	235 2	197 11	210 9
2 ‡	1,036	1,010	..	1,280	1,384	50	263 3	261 3	263 4
3	1,030	1,000	..	1,066	1,146	116	214 7	174 1	192 7
4	1,036	990	..	906	1,040	4	195 3	163 9	182 2
5	1,002	984	..	1,000	1,029	27	185 9	166 5	174 2
6	880	956	..	958	1,016	136	213 5	202 0	216 3
7	808	838	..	894	892	84	192 0	161 9	182 8
8	900	1,026	..	1,068	1,122	162	159 3	148 0	163 10
9	906	954	..	964	1,046	140	225 10	183 2	213 4
10 §	1,320	1,358	..	1,186	1,228	31	164 15	244 9	175 15
Totals -	10,042	10,312	..	10,533	11,055	840	2,047 9	1,902 1	1,973 0
Averages -	1,004	1,031	..	1,053	1,106	84	204 12	190 3	197 5
FIVE COWS.—UNSEWAGED 0									
1	1,066	1,122	..	1,122	1,140	74	236 6	243 14	238 0
2	874	902	..	878	910	36	202 0	192 12	223 13
3 *	1,042	1,106	..	1,087	1,130	88	** 235 8	165 2	212 11
4	1,185	1,176	..	1,180	1,218	33	280 14	215 7	245 12
5 ¶	1,114	1,048	..	1,128	1,184	- 8	210 1	138 15	186 5
Totals -	5,281	5,354	..	5,395	5,582	223	1,324 13	956 2	1,108 9
Averages -	1,056	1,071	..	1,079	1,116	45	264 15	191 4	221 11

* On May 25, No. 1 cow on unsewaged meadow grass, and No. 3 cow on rye-grass were transposed, and the weights entered in the Table are, from the commencement in each case, those of the newly placed cow.

† The No. 4 cow on unsewaged meadow grass put up at the commencement (April 27) increased 73 lb. by May 25, but diminished considerably in yield of milk, and as she was obviously fattening, was then replaced by the cow whose weight is entered for that date; and in the columns "increase in weight" and "yield of milk" the results obtained on the two cows, successively, are given.

‡ The No. 2 cow on sewaged meadow grass put up at the commencement (April 27) had lost 54 lbs. June 22, and being ill was removed June 24, and replaced by the animal whose weight on June 22 is entered for that date; but in the columns "increase in weight" and "yield of milk" the results obtained on the two cows, successively, are given.

§ The No. 10 cow put up at the commencement (April 27) diminished considerably in yield of milk, a shipping her calf about June 22, was then replaced by the cow whose weight is entered under that date but in the columns "increase in weight" and "yield of milk" the results obtained on the two cows, successively, are given.

TABLE XIII.

respectively on UNSEWAGED MEADOW GRASS, on SEWAGED MEADOW GRASS, and on Weeks alone, and each for Twelve Weeks with OILCAKE in addition.

SEASON, 1863.

TWELVE WEEKS ON GRASS AND OILCAKE.									TOTAL 24 WEEKS	
Weights.					In-crease (or Loss) in Weight.	Yield of Milk.			In-crease in Weight.	Average Yield of Milk per Week
Inter-mediate.	Aug. 17.	Inter-mediate.	Sept. 14.	Inter-mediate.		First Week.	Last Week.	Average.		
Aug. 2.		Sept. 8.		Sept. 23.						

MEADOW GRASS.††

lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs.	lbs. ozs.
1,138	1,176	1,186	1,199	1,204	1,123	9	220 15	229 15	235 4	103	226 0
1,234	1,253	1,274	1,268	1,272	1,246	20	156 11	111 0	138 6	129	158 8
854	878	899	897	899	866	36	156 10	171 13	169 4	120	169 8
1,113	1,118	1,146	1,141	1,098	1,090	- 2	274 7	237 6	270 6	67	251 1
1,170	1,194	1,267	1,236	1,244	1,208	66	205 1	183 3	196 2	150	209 0
5,514	5,619	5,772	5,741	5,707	5,533	129	1,013 12	953 5	1,008 6	509	1,014 1
1,103	1,124	1,154	1,143	1,141	1,107	28	202 12	190 11	201 11	114	202 13

MEADOW GRASS.

..	1,182	..	1,206	..	1,190	36	211 11	168 0	189 13	126	200 3
..	1,384	..	1,304	..	1,360	- 24	263 12	196 12	227 7	26	245 5
..	1,182	..	1,174	..	1,163	22	174 5	134 13	153 3	138	172 13
..	1,070	..	1,090	..	1,077	37	173 5	159 11	160 14	41	171 8
..	1,019	..	1,035	..	1,033	4	173 6	126 5	145 1	31	159 9
..	1,048	..	1,096	..	1,101	85	221 6	213 6	211 7	221	213 13
..	894	..	892	..	868	24	175 14	131 11	149 8	60	166 0
..	1,140	..	1,190	..	1,179	57	153 7	123 6	141 7	219	152 1
..	1,010	..	1,067	..	1,080	14	219 13	165 0	193 14	154	203 9
..	1,260	..	1,278	..	1,290	64	260 12	218 2	244 8	95	210 3
..	11,202	..	11,444	..	11,326	271	2,033 11	1,639 2	1,817 2	1,111	1,895 0
..	1,121	..	1,144	..	1,133	27	203 6	163 15	181 11	111	189 8

SEWAGED ITALIAN RYE GRASS.

..	1,153	1,130	1,121	..	1,088	- 53	258 3	216 13	235 11	22	261 14
..	922	952	934	..	914	4	194 2	189 2	195 1	40	200 7
..	1,146	1,169	1,166	..	1,120	- 10	168 8	161 6	171 9	78	192 2
..	1,271	1,299	1,295	..	1,220	2	223 10	201 4	221 2	35	233 7
..	1,195	1,224	1,234	..	1,200	16	147 6	130 14	142 15	8	140 10
..	5,684	5,754	5,780	..	5,542	- 40	901 13	899 7	906 6	183	1,037 8
..	1,137	1,151	1,150	..	1,108	- 8	198 6	179 14	193 4	37	207 8

† The No. 5 cow on rye-grass put up at the commencement (April 27), falling very ill, had lost by May 2 144 lbs., and was replaced by another cow whose weight is given under that date; but in the column "increase in weight" and "yield of milk" the results obtained on the two cows successively, are given.

†† As will be seen by reference to Appendix, Table X. p. 133, the yield of milk of No. 3 cow on rye-grass is only given for four out of the seven days of the first week of the experiment, but in the calculations the yield per diem for the remaining three days is assumed to be the same as that of the average of the ensuing week of the No. 3 cow put up on May 5.

†† From April 27 to June 8, the first crop of unsewaged meadow grass not being ready to cut, the five cow nominally fed on it had Italian rye-grass, and they also received Italian rye-grass from Aug. 2 to Sept. 8 in default of unsewaged grass; but the figures in the columns "increase in weight" and "yield of milk" are irrespective of these unavoidable irregularities.

TABLE XIV.

SUMMARY of Food consumed, and Milk and Increase yielded, by Cows fed respectively on Un Meadow Grass, on Sewaged Meadow Grass, and on Italian Rye Grass (Unsewaged and S each for Twelve Weeks alone, and each for Twelve Weeks with Oilcake in addition.

THIRD SEASON, 1863.

Periods.*		5 Cows.—Unsewaged Meadow Grass.				10 Cows.—Sewaged Meadow Grass.				5 Cows.—Unsewaged Sewaged Italian Rye			
Dates.*	No. of Days.	Average per head per day.		Average Increase (or Loss) in weight per head per week.		Average per head per day.		Average Increase (or Loss) in weight per head per week.		Average per head per day.		Average Increase (or Loss) in weight per head per week.	
		Food consumed.		Milk yielded.		Food consumed.		Milk yielded.		Food consumed.		Milk yielded.	
		Green Grass.	Oil-cake.			Green Grass.	Oil-cake.			Green Grass.	Oil-cake.		

Twelve Weeks. Grass alone.

Period	No. of Days	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs.	lbs.	lbs. ozs.
April 28 to May 4	7	112	..	30 1		113	..	29 4		112	..	37 14
May 5 to " 11	7	124	..	28 0		122	..	31 2		131	..	35 14
" 12 to " 18	7	132	..	28 6	15 8	145	..	31 11	6 12	133	..	32 10
" 19 to " 25	7	126	..	27 8		145	..	29 12		123	..	33 14
" 26 to June 1	7	109	..	34 0		153	..	29 13		150	..	34 8
June 2 to " 8	7	127	..	32 6		137	..	27 15		168	..	33 3
" 9 to " 15	7	160	..	30 5	- 1 12½	132	..	25 8	1 3½	187	..	31 10
" 16 to " 22	7	113	..	29 9		116	..	24 3		175	..	29 13
" 23 to " 29	7	106	..	27 15		140	..	26 3		162	..	27 14
" 30 to July 6	7	103	..	27 1		170	..	27 11		179	..	27 13
July 7 to " 13	7	60	..	27 3	8 4½	184	..	28 0	13 0½	196	..	27 12
" 14 to " 20	7	62	..	27 3		158	..	27 3		178	..	27 5
April 28 to " 20†	84	110	..	29 2	7 5½	143	..	28 3	7 0	157	..	31 11

Twelve Weeks. Grass and Oilcake.

Period	No. of Days	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs.	lbs.	lbs. ozs.
July 21 to July 27	7	67	3	28 15		157	3	29 1		162	3	28 5
" 28 to Aug. 3	7	54	3	29 11		179	3	29 9		155	3	28 3
Aug. 4 to " 10	7	87	3	30 8	10 13	188	3	28 8	8 13½	184	3	28 11
" 11 to " 17	7	64	3	30 7		176	3	27 5		164	3	28 15
" 18 to " 24	7	56	3	30 9		176	3	26 15		95	3	28 6
" 25 to " 31	7	50	3	29 0		180	3	25 4		90	3	28 2
Sept. 1 to Sept. 7	7	73	3	27 10	6 1½	180	3	24 9	5 14	119	3	27 9
" 8 to " 14	7	110	3	28 2		196	3	24 5		92	3	27 5
" 15 to " 21	7	99	5	28 2		149	3	24 13		128	5	27 5
" 22 to " 28	7	113	5	27 9		173	5	24 3		92	5	27 1
" 29 to Oct. 5	7	155	5	27 14	-10 6½	155	5	23 9	- 2 15½	84	5	25 12
Oct. 6 to " 12	7	155	5	27 4		155	5	23 7		108	5	25 11
July 21 to " 12†	84	90	3½	28 13	2 2½	172	3½	25 15	2 4½	112	3½	27 10

Twenty-four Weeks. First 12 Weeks, Grass alone. Second 12 Weeks, Grass and Oilcake.

Period	No. of Days	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs.	lbs.	lbs. ozs.
April 28 to Oct. 12†	168	100	1½	28 15	4 11½	153	1½	27 1	4 10	134	1½	29 10

* The dates given are of the "milk yielded," but the periods of the "food consumed" date one day earlier in as also do those of the "average increase (or loss) in weight per head per week."

† From April 27 to June 8, the first crop of unsewaged meadow grass not being ready to cut, the five cows now on it had Italian rye-grass; and they also received Italian rye-grass from Aug. 2 to Sept. 8 in default of unsewaged grass, but the figures in these lines relate to the whole period, irrespectively of these unavoidable irregularities.

TABLE XV.

RESULTS of the Analyses of 17 Samples of Sewage-water collected in the Five-acre Field.

Second Season 1861-2; November 1861—October 1862 inclusive.

GRAINS PER GALLON.																			
1861.												1862.							
	Sample numbers	Nov.	Dec.	Jan.	Feb.	March	April	May		June	July	August		September		October		Mean: 17 Sam- ples.	
		4-6.	17 & 18.	6-8.	3-5.	3-5.	3-4.	8-10.	22-24.	5-7.	10-12.	21-26.	4-9.	20 & 21.	1 & 2.	22 & 23.	—		13 & 14.
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16		17
Organic matter {	In solution	11.60	10.70	8.90	8.40	11.19	6.20	6.83	10.10	7.55	6.60	6.10	7.90	9.78	9.00	8.69	7.00	8.46	
	In suspension	25.60	39.39	20.50	11.80	16.05	9.25	20.35	10.90	15.10	13.30	19.70	13.70	14.74	17.70	19.86	9.60	7.63	
	Total	37.20	50.00	29.40	20.20	27.24	15.45	27.17	21.00	22.65	19.90	25.80	20.50	24.53	26.70	28.55	16.60	14.33	
Inorganic matter {	In solution	47.20	43.00	33.30	30.60	27.91	34.35	31.30	34.60	28.25	27.60	40.30	32.70	36.53	42.40	38.55	33.75	33.51	
	In suspension	21.00	10.80	12.30	11.40	14.77	9.65	55.15	22.60	33.35	22.40	22.70	19.10	30.03	22.20	23.13	14.70	13.05	
	Total	68.20	53.80	45.60	42.00	41.93	44.00	86.35	57.20	60.60	50.00	63.00	51.80	66.56	64.60	61.68	48.45	46.56	
Total in solution		58.80	53.70	42.20	39.00	38.40	40.55	38.02	44.70	35.80	34.20	46.40	40.50	46.30	51.40	47.84	40.75	40.21	
		46.80	50.10	32.80	23.20	30.53	18.90	75.50	33.50	47.45	35.70	42.40	31.80	44.76	39.90	47.99	24.50	20.68	
	Total solid matter	105.40	103.80	75.00	62.20	69.23	59.45	113.63	78.20	83.25	69.90	88.80	72.30	91.06	91.30	93.23	65.05	60.89	
Ammonia {	In solution	8.71	7.07	3.82	4.18	4.49	2.00	2.16	3.35	2.23	3.00	4.23	5.15	4.63	7.38	5.56	4.39	4.45	
	In suspension	2.20	2.23	1.25	1.38	1.99	0.55	1.82	0.78	1.13	0.75	1.97	1.66	1.69	1.23	2.13	0.96	1.45	
	Total	10.91	9.30	5.07	5.56	6.48	2.55	3.98	4.13	3.36	3.75	6.25	6.81	6.33	8.61	7.69	5.35	5.90	
Phosphoric acid {		2.14	1.53	
		4.95	3.98	
	Total	9.44	8.53	

COMPARATIVE AND AVERAGE COMPOSITION OF THE SEWAGE-WATER

1861.										1862.										Average for the 12 months in the two fields.					
November.		December.		January.		February.		March.		April.		May.		June.		July.		August.		September.		October.			
5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.	5- acre field.	10- acre field.		
1	1	1	1	1	1	1	1	1	1	1	1	1	2	2	1	1	2	2	2	2	2	2	2	34	
Number of Analyses -																									
Organic matter.																									
In solution -		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.	
In suspension -		11.60		10.70		9.40		8.90		8.40		7.00		6.68		6.35		8.10		8.79		8.05		8.41	
In suspension -		25.60		30.40		39.30		25.40		21.20		18.30		9.88		9.43		15.10		13.72		13.40		11.02	
Total -		37.20		41.00		50.00		34.80		29.40		25.30		15.90		27.24		10.30		22.51		26.85		22.71	
In solution -		47.20		48.40		43.00		41.70		33.30		33.70		33.30		34.35		34.60		32.90		28.25		31.90	
In suspension -		21.00		30.65		10.80		19.30		12.30		8.10		5.80		3.02		9.05		6.35		38.88		41.60	
Total -		68.20		79.05		53.80		61.00		45.60		41.80		42.00		44.00		40.55		41.28		67.08		73.50	
In solution -		58.80		59.00		53.70		51.10		42.20		43.70		39.00		39.40		38.40		37.00		40.55		41.30	
In suspension -		46.00		61.05		50.10		44.70		32.80		24.40		23.20		14.70		30.82		12.90		18.90		15.25	
Total in suspension -		104.80		120.05		103.80		95.80		75.00		67.10		62.20		54.10		69.22		50.50		58.45		66.55	
Total solid matter -		163.00		171.05		157.60		146.60		118.60		110.80		102.60		86.30		106.72		57.00		68.35		78.05	
Total solution -		8.71		8.03		7.07		6.25		3.82		4.28		4.18		4.25		4.49		4.56		4.56		4.54	
In suspension -		2.20		2.75		2.23		1.42		1.25		1.51		1.38		0.90		1.99		0.94		0.94		0.93	
Total -		10.91		10.78		9.30		7.67		5.64		5.79		5.56		5.15		6.48		5.50		5.50		5.47	
Number of Analyses -																									
Organic matter.																									
In solution -		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.		Gns.	
In suspension -		11.60		10.70		9.40		8.90		8.40		7.00		6.68		6.35		8.10		8.79		8.05		8.41	
In suspension -		25.60		30.40		39.30		25.40		21.20		18.30		9.88		9.43		15.10		13.72		13.40		11.02	
Total -		37.20		41.00		50.00		34.80		29.40		25.30		15.90		27.24		10.30		22.51		26.85		22.71	
In solution -		47.20		48.40																					

... of Ground-water collected in the Ten-acts Field.

RESULTS of the Analyses of 12 Samples of Sewage-Water.

GRAINS PER GALLON.												
1862.										1863.		
November		December		January		February		March		April	May	Mean; 13 Samples.
3-5.	17-19.	1-3.	15-17.	5-7.	19-21.	2-4.	16-18.	8	17	7.	7 & 8.	
Sample Numbers	1	2	3	4	5	6	7	8	9	10	11	12
Organic matter.	In solution	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.
	In suspension	8.30	8.36	8.00	10.20	4.35	4.35	5.06	7.63	6.85	7.60	7.11
		19.15	25.08	21.35	15.10	16.25	27.55	10.90	7.40	13.75	30.60	17.06
	Total	25.35	32.04	30.45	25.30	30.60	33.00	16.85	15.05	20.10	38.10	24.77
Inorganic matter.	In solution	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.
	In suspension	34.60	31.90	26.30	31.90	29.15	33.50	29.85	31.25	28.20	43.80	31.40
		21.60	30.35	34.55	24.50	23.65	37.00	16.70	15.55	30.40	18.50	24.92
	Total	56.10	63.25	60.85	56.40	51.80	69.50	46.55	47.10	58.60	61.30	56.41
Total in solution	40.70	43.30	34.90	42.10	33.50	36.55	35.80	38.90	33.55	50.30	28.60	
Total in suspension	40.72	43.60	38.65	39.60	38.90	64.65	27.60	23.25	44.15	39.10	43.68	
Total solid matter	81.42	84.90	75.01	81.70	72.40	101.50	63.40	63.15	76.70	89.40	81.18	
Ammonia	In solution	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.
	In suspension	4.95	4.06	1.91	2.53	3.10	4.60	4.23	5.56	3.41	6.40	4.05
		1.72	1.66	1.46	1.13	0.77	2.84	1.12	0.84	1.63	1.28	1.43
	Total	6.67	6.55	3.37	3.66	3.87	7.44	5.34	6.40	5.04	7.68	5.48

TABLE XX.
Results of the Analyses of 10 Mixed Samples of Sewage-water, each collected partly in the Five-acre and partly in the Ten-acre Field.
Third Season 1863; June—October inclusive.

GRAINS PER GALLON.														
1863.														
June			July			August		September		October			Mean	
1-6.		15-19.	6-10.		20-24.	9-7.	17-21.	7-11.	21-25.	5-9.		19-24.	10 Samples.	
1		3	3		4	5		7	8	9		10		
Grains.		Grains.	Grains.		Grains.	Grains.		Grains.	Grains.	Grains.		Grains.	Grains.	
{ Organic matter. {	In solution	8.50	7.15	9.10	9.00	13.00	9.90	9.30	8.50	11.10	11.70	9.73	Grains.	
	In suspension	28.15	18.00	23.00	36.80	24.30	34.40	76.30	31.60	53.10	53.10	35.97	Grains.	
	Total	36.65	25.15	31.10	45.80	37.30	44.30	85.60	40.10	63.20	63.20	45.70	Grains.	
{ Inorganic matter. {	In solution	45.70	38.15	46.20	43.20	40.50	52.10	47.30	54.40	54.10	53.50	47.53	Grains.	
	In suspension	43.50	23.65	36.20	21.00	33.10	25.10	137.00	40.60	61.80	61.80	46.93	Grains.	
	Total	89.20	61.80	82.40	64.20	73.60	77.20	184.30	104.00	115.90	115.90	94.46	Grains.	
Total in solution		54.20	45.30	55.30	52.20	53.50	62.00	56.60	62.90	65.20	65.20	57.25	Grains.	
Total in suspension		71.65	41.65	58.20	57.80	57.40	59.50	213.30	81.20	113.90	113.90	88.90	Grains.	
Total solid matter		125.85	86.95	113.50	110.00	110.90	121.50	269.90	144.10	179.10	179.10	140.15	Grains.	
				7.23	6.46	6.75	7.47	7.40	10.21	9.76	9.76	7.68	Grains.	
				3.79	3.79	3.57	1.84	4.74	2.35	3.05	3.05	3.77	Grains.	

TABLE XXII.
Results of the Analyses of 8 Samples of Drainage-water collected in the Five-acre Field.
Second Season 1862; May—October inclusive.

GRAINS PER GALLON.											
1862.											
	May 8-10.	July 21-22.	August		September		October		Mean; 8 Samples.		
			4-9.	20 and 21.	1 & 2.	22 & 23.	—	12 & 14.			
Sample Numbers	1	2	3	4	5	6	7	8			
Organic matter { In solution In suspension Total	Grains. 5.50 •	Grains. 10.50 •	Grains. 8.50 2.40	Grains. 6.04 2.65	Grains. 6.10 1.90	Grains. 6.70 2.70	Grains. 8.20 1.00	Grains. 8.90 0.25	Grains. 7.18 1.40		
Inorganic matter { In solution In suspension Total	Grains. 25.20 •	Grains. 31.70 •	Grains. 24.21 7.43	Grains. 8.69 2.77	Grains. 39.90 0.40	Grains. 39.00 0.90	Grains. 35.00 0.80	Grains. 34.09 2.09	Grains. 24.50 1.81		
Total in solution	Grains. 30.70 •	Grains. 42.20 •	Grains. 42.71	Grains. 43.98	Grains. 46.00	Grains. 45.70	Grains. 45.20	Grains. 39.99	Grains. 41.68		
Total in suspension	Grains. •	Grains. •	Grains. 9.97	Grains. 5.42	Grains. 2.30	Grains. 3.60	Grains. 1.80	Grains. 2.51	Grains. 3.21		
Total solid matter	Grains. 30.70	Grains. 42.20	Grains. 52.68	Grains. 49.40	Grains. 48.30	Grains. 49.30	Grains. 45.00	Grains. 42.50	Grains. 44.89		
Ammonia { In solution In suspension	Grains. 0.60 —	Grains. 1.03 —	Grains. 8.03 0.76	Grains. 1.21 0.23	Grains. 0.39 0.15	Grains. 0.27 0.38	Grains. 0.25 0.23	Grains. 0.54 0.15	Grains. 0.80 0.24		

Results of the Analyses of 11 Samples of Drainage Water collected in the Ten-acre Field. Second Season 1862; May—October inclusive.

GRAINS PER GALLON.												
1862.												
Sample Numbers	May		June		July		August		September		October	
	8-7.	19-21.	2-4.	7-12.	21-26.	4-9.	16 & 19.	1 & 2.	22 & 23.	—	13 & 14.	Mean; 11 Samples.
1	1	2	3	4	5	6	7	8	9	10	11	
Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.
5.70	7.60	9.30	5.10	9.90	8.16	8.29	8.92	7.40	7.10	11.41	7.50	7.88
•	•	•	•	•	2.13	5.10	0.59	1.80	4.20	2.87	4.20	1.88
5.70	7.60	9.30	5.10	9.90	8.29	5.10	8.92	8.60	11.30	14.28	11.80	9.23
33.30	36.80	37.30	33.70	33.90	31.08	33.70	33.38	46.30	43.00	33.33	38.10	37.10
•	•	•	•	•	10.81	•	5.88	8.70	6.90	6.01	3.80	3.74
33.30	36.80	37.30	33.70	33.90	41.83	33.70	44.26	53.00	46.90	44.24	40.90	40.84
39.00	44.40	46.60	38.80	42.80	37.18	38.80	47.30	53.70	49.10	49.74	45.60	44.98
•	•	•	•	•	13.94	•	6.47	9.90	11.10	8.88	7.10	8.13
39.00	44.40	46.60	38.80	43.80	50.13	38.80	53.77	63.60	60.20	58.62	53.70	50.06
1.45	1.64	1.53	1.66	1.66	3.25	1.66	1.68	3.79	1.87	1.92	1.18	1.35
•	•	•	•	•	0.39	•	0.60	0.60	1.04	0.54	0.45	0.33
1.45	1.64	1.53	1.66	1.66	3.64	1.66	2.28	4.39	3.61	3.46	1.63	2.13
Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.	Grains.
1.45	1.64	1.53	1.66	1.66	3.64	1.66	2.28	4.39	3.61	3.46	1.63	2.13

* Too small for estimation.

TABLE XXIV.

COMPARATIVE and average Composition of the Drainage-water collected in the Two Fields,
Second Season 1862; May—October inclusive.

Second Season 1862; May—October inclusive.

GRAINS PER GALLON.

1862.

Average
for the
6
Months
in the
2 Fields.

—

May.

June.

July.

August.

September.

October.

19

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

Grains.

* Too small for estimation.

TABLE XXV.
Results of the Analyses of 21 Samples of Drainage-water collected in the Five-acre Field,
Third Season 1862-3; November 1862—October 1863 inclusive.

		GRAINS PER GALLON.																								
		1862.						1863.																		
	Sample numbers	November		December		January		February		March		April		May		June		July		August		September		October.		Mean; 21 Samples.
		3-8.	20-22.	4-6.	18-20.	8-10.	22-24.	5-7.	19-21.	2, and 16, 18-21.	9-10.	11-12.	13-14.	15-16.	17-18.	19-20.	21-22.	23-24.	25-26.	27-28.	29-30.	31-32.	33-34.	35-36.	37-38.	
Organic matter {	In solution -	5.05	7.59	9.91	9.25	9.05	9.10	10.60	5.65	4.25	4.20	5.40	10.25	11.00	12.50	10.85	6.55	13.95	11.50	13.00	6.40	7.00	8.30	6.90	8.87	8.57
	In suspension -	0.47	1.36	2.22	1.30	2.30	2.30	0.40	0.20	0.20	0.20	0.20	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
	Total -	6.12	8.95	12.13	10.55	11.35	11.40	11.00	5.85	4.45	4.40	5.60	10.70	11.35	12.85	11.20	6.90	14.30	11.85	13.35	6.75	7.35	8.65	7.25	9.12	8.92
Inorganic matter {	In solution -	39.35	38.89	36.02	35.50	29.25	33.10	32.00	34.25	36.80	36.75	34.50	41.75	35.00	38.50	44.10	37.20	49.10	49.00	46.60	38.55	38.15	38.15	38.15	38.15	38.15
	In suspension -	1.58	10.33	2.10	0.30	0.60	0.60	2.90	1.65	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
	Total -	40.93	49.22	38.12	35.80	29.85	33.10	34.90	35.90	36.80	36.75	34.65	41.95	35.20	38.70	44.30	37.40	49.30	49.20	46.80	38.75	38.35	38.35	38.35	38.35	38.35
Total in solution -		45.00	49.48	45.93	44.75	38.30	42.20	42.00	39.90	41.05	40.95	39.90	49.55	52.05	40.70	55.30	45.00	49.00	53.50	55.10	56.30	51.80	46.01	46.01	46.01	46.01
	Total in suspension -	2.05	11.69	4.32	1.60	2.90	2.90	3.30	1.85	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40
	Total solid matter-	47.05	58.17	50.25	46.35	41.20	45.10	45.30	41.75	41.45	41.35	40.30	50.95	53.45	41.10	55.70	45.40	49.40	53.90	55.50	56.70	52.20	46.41	46.41	46.41	46.41
Ammonia {	In solution -	0.33	0.94	1.55	0.94	0.89	0.17	0.85	1.37	0.40	0.40	0.33	1.70	0.43	0.40	0.50	0.90	0.31	0.31	0.31	0.31	0.31	0.31	0.31	0.31	0.31
	In suspension -	none	0.62	0.04	0.14	0.19	0.19	0.32	0.10	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
	Total -	0.33	1.56	1.59	1.08	1.08	0.36	1.17	1.47	0.60	0.60	0.53	1.90	0.63	0.60	0.70	0.90	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51

* Too small for estimation.

* Too small for estimation.

TABLE XXVI.
Results of the Analyses of 23 Samples of Drainage-water collected in the Ten-acre Field.
Third Season 1862-3; November 1862—October 1863 inclusive.

		GRAINS PER GALLON.												Mean; 23 Sam- ples.											
		1862.						1863.																	
Sample Numbers		November		December		January		February		March		April		May		June		July		August		September		October	
		3-5.	17-19.	1-3.	15-17.	5-7.	19-21.	2-4.	16-18.	3.	7.	7 & 8.	14 & 15.	1-6.	18 & 19.	7 & 9.	23 & 24.	6 & 7.	20 & 21.	10 & 11.	24 & 25.	8 & 9.	21-24.		
1		2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22			
Organic matter.	In solution -	Gns. 10-20	Gns. 6-90	Gns. 7-40	Gns. 10-40	Gns. 5-65	Gns. 5-30	Gns. 6-10	Gns. 5-25	Gns. 8-60	Gns. 9-20	Gns. 7-75	Gns. 8-50	Gns. 9-60	Gns. 3-50	Gns. 8-60	Gns. 10-50	Gns. 7-50	Gns. 6-30	Gns. 8-06	Gns. 6-70	Gns. 7-98			
	In suspension -	Gns. 1-85	Gns. 0-79	Gns. 4-45	Gns. 2-32	Gns. 1-00	Gns. 1-20	Gns. 1-60	Gns. 1-85	Gns. 8-45	Gns. 0-60	Gns. 3-90	Gns. 1-50	Gns. 3-40	Gns. 2-50	Gns. 3-90	Gns. 5-70	Gns. 5-00	Gns. 2-90	Gns. 6-00	Gns. 5-10	Gns. 3-29			
	Total	12-05	10-99	11-35	9-72	8-90	11-60	7-60	11-30	7-70	17-25	9-80	11-65	10-00	13-30	11-00	12-50	16-00	12-50	9-20	14-00	11-80	11-27		
Inorganic matter.	In solution -	Gns. 36-60	Gns. 35-00	Gns. 38-15	Gns. 38-40	Gns. 33-30	Gns. 33-60	Gns. 34-05	Gns. 33-85	Gns. 34-45	Gns. 43-80	Gns. 45-30	Gns. 46-75	Gns. 38-00	Gns. 50-70	Gns. 47-20	Gns. 52-10	Gns. 31-40	Gns. 53-20	Gns. 54-25	Gns. 53-40	Gns. 42-00	Gns. 41-35		
	In suspension -	Gns. 4-20	Gns. 6-11	Gns. 4-35	Gns. 2-10	Gns. 6-50	Gns. 2-40	Gns. 2-85	Gns. 7-25	Gns. 3-35	Gns. 7-45	Gns. 0-95	Gns. 1-20	Gns. 4-35	Gns. 6-25	Gns. 6-00	Gns. 0-50	Gns. 6-80	Gns. 1-90	Gns. 2-65	Gns. 5-10	Gns. 3-94			
	Total	40-80	41-71	42-50	40-50	40-10	36-00	36-90	41-10	37-80	51-25	46-25	47-95	43-25	56-95	53-20	52-60	38-20	55-10	56-90	58-50	44-10	45-29		
Total in solution -		46-80	45-80	45-05	45-80	41-20	44-00	39-70	39-35	40-55	88-65	92-00	94-50	82-00	101-40	94-20	104-20	62-80	109-10	110-45	108-90	86-10	87-33		
Total in suspension -		6-05	6-90	8-80	4-42	7-80	3-60	4-80	13-05	4-95	3-75	15-90	1-55	5-10	8-85	8-50	4-40	12-50	6-90	5-55	11-70	7-20	7-23		
Total solid matter -		52-85	52-70	53-85	50-22	49-00	47-60	44-50	52-40	45-50	108-95	107-90	96-05	87-10	110-25	102-70	108-60	75-30	116-30	116-95	120-60	93-30	94-56		
Ammonia.	In solution -	Gns. 1-14	Gns. 3-40	Gns. 1-47	Gns. 1-43	Gns. 1-08	Gns. 0-80	Gns. 1-17	Gns. 1-26	Gns. 1-12	Gns. 3-87	Gns. 2-11	Gns. 3-11	Gns. 2-00	Gns. 1-64	Gns. 1-67	Gns. 1-05	Gns. 0-50	Gns. 2-37	Gns. 3-06	Gns. 3-75	Gns. 2-09	Gns. 1-83		
	In suspension -	Gns. 0-75	Gns. 0-29	Gns. 0-67	Gns. 0-13	Gns. 0-19	Gns. 0-31	Gns. 0-20	Gns. 0-19	Gns. 0-19	Gns. 0-55	Gns. ..	Gns. 0-10	Gns. 0-07	Gns. 0-57	Gns. 0-13	Gns. 0-19	Gns. 0-37	Gns. 0-43	Gns. 0-20	Gns. 0-89	Gns. 0-12	Gns. 0-31		
	Total	1-89	3-69	2-14	1-56	1-27	1-11	1-37	1-45	1-31	4-42	2-11	3-21	2-07	2-21	1-80	2-14	0-87	2-80	3-26	4-64	2-21	2-16		

COMPARATIVE AND AVERAGE COMPOSITION OF THE DRAINAGE-WATER COLLECTED IN THE TWO FIELDS.

Third Season 1862-3; November 1862—October 1863 inclusive.

GRAINS PER GALLON.												Average for the 12 months in the two fields.											
1862.												1863.											
November.		December.		January.		February.		March.		April.		May.		June.		July.		August.		September.		October.	
5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.	5- acre Field.	10- acre Field.		
Gns. 6-62 Gals. 10-20	Gns. 9-28 Gals. 7-13	Gns. 9-08 Gals. 7-13	Gns. 9-15 Gals. 7-13	Gns. 8-13 Gals. 5-58	Gns. 8-13 Gals. 5-58	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22	Gns. 8-42 Gals. 4-22		
0-92	1-32	1-76	3-39	1-10	3-57	1-60	3-57	1-60	3-57	1-60	3-57	1-60	3-57	1-60	3-57	1-60	3-57	1-60	3-57	1-60	3-57		
7-54	11-32	11-34	10-54	10-23	10-25	8-43	9-45	4-22	7-70	5-40	7-10	12-50	13-53	8-60	10-83	12-15	13-60	14-25	6-70	10-85	7-60	13-20	
In solution - 39-12 In suspension - 5-93	35-76 5-16	38-28 5-16	31-17 3-22	33-45 4-60	33-45 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	36-78 2-25	
Total - 45-07	41-50	41-50	31-47	38-05	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	33-40	
In-organic matter. 45-74	46-30	45-34	45-43	40-25	42-60	41-00	40-55	39-90	38-65	40-55	38-65	40-55	38-65	40-55	38-65	40-55	38-65	40-55	38-65	40-55	38-65	40-55	
Total in solution - 6-87	6-48	2-96	6-61	1-45	5-70	2-58	8-92	4-95	0-15	3-75	4-45	8-73	0-88	5-48	8-00	9-08	10-90	8-45	3-60	6-23	3-25	9-45	
Total in suspension - 52-61	62-78	46-30	52-04	41-70	48-30	43-83	48-45	41-00	45-50	40-05	42-40	54-00	62-28	47-25	56-43	58-15	60-98	60-80	59-65	58-85	57-30	64-50	
Total solid matter - 0-64	2-27	1-25	1-45	0-53	0-94	1-11	1-34	0-52	1-12	0-33	0-46	1-70	2-99	0-41	2-55	0-70	1-66	0-31	1-23	0-84	0-63	2-92	
In solution - 0-31	0-52	0-69	0-40	0-10	0-25	0-21	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	0-19	
In suspension - 0-93	2-79	1-34	1-85	0-63	1-19	1-92	1-53	0-52	1-31	0-33	0-65	1-70	3-27	0-60	2-64	0-90	2-01	0-38	1-51	0-55	2-53	3-43	
Total - 0-93	2-79	1-34	1-85	0-63	1-19	1-92	1-53	0-52	1-31	0-33	0-65	1-70	3-27	0-60	2-64	0-90	2-01	0-38	1-51	0-55	2-53	3-43	

TABLE XXVIII.
 DETAILS of the Sampling, and of the Determinations of Dry Substance and Mineral Matter (Ash), in the Unsewaged and in the Sewaged Meadow Grass.
 Second Season, 1862.

Particulars of Sampling.				Determinations of Dry Substance and Mineral Matter (Ash).													
Fields.	Plots.	Number of Samples taken.	Weights.			Actual Weights.			Per-centages.								
			In Fresh State.		Air-dried.	Equal in Fresh State.	Dry Sub-stance (at 212° Fahr.).	Mineral Matter (Ash).	Dry in Fresh.		Ash in Fresh.		Ash in Dry.				
			Each Sample.	Total.					Each Total.	Mean.	Each Exper.	Mean.		Each Exper.	Mean.		
FIRST CROP.																	
Five-acre	1 (Unsewaged)	-	44	2½	110	35	0	50	ozs.	13.5	30.286	10.474	0.946	28.66	26.67	9.408	9.020
	2 (Sewaged)	-	25	5	125	33	8	50	ozs.	13.5	30.286	10.460	0.946	28.66	26.67	9.408	9.020
	3 (Sewaged)	-	17	5	85	14	10	50	ozs.	12.5	43.077	10.930	0.926	22.73	22.75	1.013	8.443
	4 (Sewaged)	-	23	5	115	20	13	50	ozs.	12.5	72.650	10.460	1.010	14.57	14.41	1.390	9.631
	0* (Unsewaged)	-	27	2½	67½	17	0	50	ozs.	12.5	69.069	10.537	0.974	15.23	15.23	1.410	9.323
	0* (Unsewaged)	-	8	2½	20	5	12	50	ozs.	12.5	49.633	10.750	0.966	21.63	21.64	1.946	8.966
Ten-acre	1 (Unsewaged)	-	34	2½	85	26	8	50	ozs.	12.5	43.478	10.256	1.015	23.59	23.63	2.335	9.929
	2 (Sewaged)	-	20	5	100	22	12	50	ozs.	12.5	43.478	10.256	1.015	23.59	23.63	2.335	9.929
	3 (Sewaged)	-	14	5	70	11	4	50	ozs.	12.5	43.478	10.256	1.024	23.64	23.64	2.355	9.961
Ten-acre	1 (Unsewaged)	-	34	2½	85	26	8	50	ozs.	12.5	40.094	10.780	1.013	26.89	26.86	2.527	9.397
	2 (Sewaged)	-	20	5	100	22	12	50	ozs.	12.5	40.094	10.755	1.007	26.82	26.82	2.513	9.350
	3 (Sewaged)	-	14	5	70	11	4	50	ozs.	12.5	64.045	10.700	1.006	19.47	19.52	1.835	9.368
Ten-acre	1 (Unsewaged)	-	34	2½	85	26	8	50	ozs.	12.5	77.778	10.470	1.190	13.46	13.51	1.537	11.870
	2 (Sewaged)	-	20	5	100	22	12	50	ozs.	12.5	77.778	10.470	1.190	13.46	13.51	1.537	11.870
	3 (Sewaged)	-	14	5	70	11	4	50	ozs.	12.5	77.778	10.470	1.190	13.46	13.51	1.537	11.870

Table XXVIII.—continued.
Details of the Sampling, and of the Determinations of Dry Substance and Mineral Matter (Ash), in the Unsewaged and in the Sewaged Meadow Grass.
 Second Season, 1862.

Particulars of Sampling.				Determinations of Dry Substance and Mineral Matter (Ash).												
Fields.	Plots.	Number of Samples taken.	Weights.			Actual Weights.			Percentages.							
			In Fresh State.		Air-dried.	Equal in Fresh State.	Dry Substance (Fig. Fair.)	Mineral Matter (Ash).	Dry in Fresh.		Ash in Fresh.		Ash in Dry.			
			Each Sample.	Total.	Total.				Each Exper. [†]	Mean.	Each Exper. [†]	Mean.				
SECOND CROP.																
Five-acre	1 (Unsewaged)	-	8	2½	20	5	8	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.
	2 (Sewaged)	-	14	5	70	12	10	45.455	10.338	0.930	22.74	22.75	2.046	8.996	8.996	8.996
	3 (Sewaged)	-	16	5	80	16	0	45.455	10.345	0.930	23.70	14.31	2.046	8.996	11.817	12.004
	4 (Sewaged)	-	10	5	50	12	0	69.307	9.848	1.180	14.41	16.43	1.703	10.411	10.376	10.390
	0* (Unsewaged)	-	30	2½	75	10	6	69.307	9.848	1.200	14.41	19.39	2.003	10.349	12.377	12.349
Ten-acre	1 (Unsewaged)	-	15	2½	37½	8	4	62.500	10.150	1.900	17.87	17.90	2.091	11.823	11.683	11.683
	2 (Sewaged)	-	8	5	40	8	0	56.812	10.188	1.170	17.83	16.15	1.856	11.602	11.513	11.513
	3 (Sewaged)	-	6	5	30	6	6	62.500	10.085	1.160	16.13	18.96	1.861	10.721	10.153	10.115
	4 (Sewaged)	-	13	5	65	12	14	68.419	11.260†	1.128	18.74	16.73	1.907	10.689	10.065	10.071
									68.424	11.046	1.122	18.78	16.73	1.889	10.685	10.065
* Designated Plot "0" when unmeasured land.																

Table XXVIII.—*continued.*
 Details of the Sampling, and of the Determinations of Dry Substance and Mineral Matter (Ash), in the Unsewaged and in the Sewaged Meadow Grass.
 Second Season, 1862.

Particulars of Sampling.					Determinations of Dry Substance and Mineral Matter (Ash).											
Fields.	Plots.	Number of Samples taken.	Weights.			Actual Weights.				Percentages.						
			In Fresh state.		Air-dried.	Equal in Fresh state.	Dry Sub-stance (at 212° Fahr.).	Mineral Matter (Ash).	Dry in Fresh.		Ash in Fresh.		Ash in Dry.			
			Each Sample.	Total.					Total.	Each Exper.	Mean.	Each Exper.	Mean.	Each Exper.	Mean.	
THIRD CROP.*																
Five-acre	2 (Sewaged)	-	5	25	5 10	ozs. 12.5	ozs. 55.556	10.090	1.345	18.16	18.24	2.421	2.417	12.550	13.247	
	3 (Sewaged)	-	8	40	6 8	12.5	55.556	10.180	1.340	18.32	18.32	2.412	2.412	13.163	13.163	
	4 (Sewaged)	-	15	75	13 3	12.5	79.923	9.918	1.300	12.89	12.89	1.807	1.807	14.018	14.017	
						12.5	76.923	9.916	1.290	14.20	14.19	1.793	1.793	12.631	12.631	
Ten-acre	2 (Sewaged)	-	5	25	4 10	12.5	67.568	9.810	1.240	14.52	14.52	1.835	1.835	12.640	12.634	
	3 (Sewaged)	-	10	60	11 0	12.5	68.182	9.800	1.170	14.51	14.51	1.716	1.716	11.580	11.587	
	4 (Sewaged)	-	9	45	8 12	12.5	64.266	10.146	1.290	15.75	15.80	1.867	1.867	11.827	12.078	
						12.5	64.266	10.164	1.253	15.81		1.949	1.908	12.323		
FOURTH CROP.																
Ten-acre	4 (Sewaged)	-	{ 8 1	20	8 8	19.5 12.5	20.412 22.412	9.938 9.940	1.284 1.286	23.79 23.80	23.80	4.387 4.413	4.500	12.990 13.068	13.069	

Third Season, 1863.

Particulars of Sampling.				Weights.				Actual Weights.				Determinations of Dry Substance and Mineral Matter (Ash).						
Fields.	Plots.	Number of Samples taken.	In fresh state.		Air-dried.	Total.	Each Sample.	Total.	Air-dried.	Equal in fresh state.	Dry Substance (at 212° Fahr.).	Mineral Matter (Ash).	Dry in Fresh.		Ash in Fresh.		Ash in Dry.	
			Each.	Taken.									Each Expt ^r .	Mean.	Each Expt ^r .	Mean.	Each Expt ^r .	Mean.
First Crop.																		
Five-acre	1 (Unsewaged)	6	2½	15	6	1	50	12.5	30.925	11.153	0.850	36.06	36.10	2.652	2.878	7.352	7.072	
	2 (Sewaged)	11	5	55	13	8	50	12.5	30.925	10.937	1.006	36.13	21.49	3.104	2.947	8.561	9.187	
	3 (Sewaged)	8	5	40	8	1	50	12.5	30.925	10.942	1.004	17.45	17.59	1.971	1.973	9.198	10.129	
	4 (Sewaged)	10	5	50	9	5	50	12.5	30.925	10.935	1.110	16.28	16.26	1.689	1.682	10.069	10.218	
	0* (Unsewaged)	5	2½	12½	3	2½	50	12.5	30.925	11.303	1.050	22.63	22.75	2.147	2.154	9.371	9.382	
	00* (Sewaged)	1	5	5	1	2½	18½	12.5	30.925	10.906	1.383	30.45	20.49	1.700	1.705	8.331	8.352	
Ten-acre	1 (Unsewaged)	12	2½	30	13	4	50	12.5	38.302	11.974	0.807	39.83	39.79	2.851	2.899	7.158	7.286	
	2 (Sewaged)	8	5	40	8	7	50	12.5	38.302	11.249	0.834	38.75	38.79	2.947	2.947	7.414	7.286	
	3 (Sewaged)	8	5	40	8	7	50	12.5	59.256	10.972	0.900	18.52	18.56	1.534	1.545	8.285	8.325	
	4 (Sewaged)	8	5	40	9	2	50	12.5	54.793	10.966	0.922	20.01	20.03	1.555	1.565	9.354	9.383	
	0* (Unsewaged)	5	2½	12½	2	6	38	4.75	74.375	10.982	1.005	20.04	20.03	2.000	1.999	9.995	9.983	
	0* (Unsewaged)	5	2½	12½	4	14	50	6.25	74.375	10.828	1.144	14.56	14.56	1.538	1.549	10.565	10.659	
Twenty-acre	1 (Unsewaged)	6	2½	15	3	12	50	12.5	25.000	4.269	0.370	17.08	17.09	1.480	1.512	8.697	8.850	
	0* (Unsewaged)	14	2½	35	8	10	50	6.25	25.000	4.273	0.380	17.08	17.09	1.544	1.512	9.053	9.083	
	0* (Unsewaged)	5	2½	12½	4	14	50	6.25	16.026	5.008	0.480	34.96	34.96	2.965	3.014	8.559	8.621	
	0* (Unsewaged)	5	2½	12½	4	14	50	6.25	16.026	5.008	0.480	34.96	34.96	2.965	3.014	8.559	8.621	
Twenty-acre	1 (Unsewaged)	6	2½	15	3	12	50	12.5	50.000	10.659	0.976	21.32	21.30	1.952	1.950	9.157	9.157	
	0* (Unsewaged)	14	2½	35	8	10	50	12.5	50.725	10.638	0.946	21.30	21.26	1.965	1.950	9.156	9.157	
* Unmeasured land; designated Plot 0 when unsewaged, and Plot 00 when sewaged.																		
† No sewage was applied for the first crop of Plots 2 and 3, and no samples were taken from the Produce.																		

Table XXIX.—*continued.*
 Details of the Sampling, and of the Determinations of Dry Substance and Mineral Matter (Ash), in the Unsewaged and in the Sewaged Meadow Grass and Rye-grass.
 Third Season, 1863.

Particulars of Sampling.				Determinations of Dry Substance and Mineral Matter (Ash).												
Fields.	Plots.	Number of Samples taken.	Weights.			Actual Weights.			Per-centages.							
			In Fresh State.		Air-dried.	Air-dried.	Equal in Fresh State.	Dry Sub- stance (at 212° Fahr.).	Mineral Matter (Ash).	Dry in Fresh.		Ash in Fresh.		Ash in Dry.		
			Each Sample.	Total.	Taken.					Each Expt.	Mean.	Each Expt.	Mean.	Each Expt.	Mean.	
SECOND CROP.																
Five-acre	1 (Unsewaged)	-	-	2	1	10	4	2	50	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.
	2 (Sewaged)	-	-	6	5	30	6	5	50	30.303	10.420	1.190	34.89	34.42	3.927	3.818
	3 (Sewaged)	-	-	9	5	45	7	12	50	30.303	10.437	1.124	18.51	18.51	2.007	1.997
	4 (Sewaged)	-	-	4	5	20	4	2	50	30.401	10.993	1.180	18.51	18.51	1.987	1.964
	*0 (Unsewaged)	-	-	2	5	20	5	10	50	72.581	10.814	1.130	14.90	14.90	1.557	1.571
	*00 (Sewaged)	-	-	1	10	10	2	7½	39.75	72.581	10.828	1.140	17.78	17.80	1.571	1.564
					4	5	20	4	2	50	60.698	10.774	1.007	17.82	17.82	1.662
Ten-acre	1 (Unsewaged)	-	-	1	5	7½	1	9½	25.5	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.	ozs.
	2 (Sewaged)	-	-	6	5	30	6	1	50	29.095	5.454	0.675	18.15	18.17	2.250	2.242
	3 (Sewaged)	-	-	7	5	35	6	8	50	29.095	5.444	0.670	18.18	17.67	2.250	1.875
	4 (Sewaged)	-	-	6	5	30	6	7	49.895½	61.866	10.932	1.160	17.67	17.66	1.875	1.929
	*0 (Unsewaged)	-	-	3	5	17½	5	13	50	61.866	10.918	1.255	16.25	16.25	1.887	1.830
	*00 (Sewaged)	-	-	1	10	10	1	15	31	67.308	10.940	1.250	16.25	16.25	1.813	1.830
					6	5	30	6	7	49.895½	67.308	10.937	1.220	18.80	18.82	1.927
Unsewaged	1 (Unsewaged)	-	-	7	2½	17½	5	13	50	58.127	10.927	1.129	26.60	26.63	3.481	3.069
	2 (Sewaged)	-	-	9	5	45	9	10	50	58.127	10.951	1.129	26.60	26.63	2.637	2.637
	3 (Sewaged)	-	-	9	5	45	9	10	50	37.631	10.069	1.310	17.17	17.15	1.865	1.869
	4 (Sewaged)	-	-	7	2½	17½	4	12	50	39.090	6.868	0.746	23.69	23.70	1.902	1.912
	*0 (Unsewaged)	-	-	9	5	45	9	10	50	39.090	6.851	0.749	18.83	18.81	1.578	1.578
	*00 (Sewaged)	-	-	9	5	45	9	10	50	46.053	10.915	0.922	17.48	17.48	1.642	1.642
					9	5	45	9	10	46.053	10.915	0.922	17.55	17.55	1.657	1.657

Table XXIX—continued.

Details of the Sampling, and of the Determinations of Dry Substance and Mineral Matter (Ash), in the Unsewaged and in the Sewaged Meadow Grass and Rye-grass.

Third Season, 1863.

Particulars of Sampling.				Determinations of Dry Substance and Mineral Matter (Ash).												
Fields.	Plots.	Number of Samples taken.	Weights.			Actual Weights.				Per-centages.						
			In fresh state.		Air-dried.	Equal in fresh State.	Dry Sub-stance (at 212° Fahr.)	Mineral Matter (Ash).	Dry in Fresh.		Ash in Fresh.		Ash in Dry.			
			Each Sample.	Total.					Total.	Taken.	Each Exper.	Mean.	Each Exper.	Mean.	Each Exper.	Mean.
THIRD CROP.																
Five-acre	2 (Sewaged)	5	lbs. 5	lbs. 25	ozs. 5	50	ozs. 28.140	10.273	0.25.	17.67	17.67	2.073	2.046	11.730	11.584	
	3 (Sewaged)	5	5	25	6	50	28.140	10.255	0.25.	17.66	17.66	2.019	1.976	11.437	11.437	
	4 (Sewaged)	6	5	25	3	50	28.563	10.139	0.25.	10.95	10.95	1.376	1.374	12.565	12.589	
			5	30	6	50	32.563	10.070	0.25.	10.88	10.88	1.372	1.374	12.612	12.612	
Ten-acre	2 (Sewaged)	5	5	25	3	50	57.602	10.142	0.25.	17.58	17.58	2.018	2.023	11.477	11.500	
	3 (Sewaged)	5	5	25	3	50	57.602	10.143	0.25.	17.58	17.58	2.028	2.028	11.535	11.535	
	4 (Sewaged)	8	5	40	7	50	79.355	9.856	0.25.	13.42	13.42	1.439	1.448	11.587	11.665	
	90* (Sewaged)	1	10	10	2 9½	41½	79.355	9.845	0.25.	13.41	13.41	1.457	1.448	11.742	11.665	
Rye-grass	1 (Unsewaged)	8	2½	20	8	14	69.767	10.140	0.25.	14.53	14.53	1.720	1.734	11.854	11.889	
	2 (Sewaged)	11	5	53	17	6	69.767	10.206	0.25.	14.53	14.53	1.747	1.734	11.944	11.889	
	3 (Sewaged)	14	5	70	14	4	66.667	10.115	0.25.	15.17	15.17	1.761	1.761	11.668	11.668	
	0* (Unsewaged)	9	2½	22½	9	14	66.667	10.115	0.25.	21.12	21.12	2.480	2.478	11.727	11.736	
• Unmeasured land; designated Plot 0 when unsewaged, and Plot 00 when sewaged.																
Rye-grass	1 (Unsewaged)	8	2½	20	8	14	39.996	8.439	0.25.	36.35	36.35	2.503	2.503	6.885	6.910	
	2 (Sewaged)	11	5	53	17	6	28.169	10.238	0.25.	36.34	36.34	2.521	2.521	6.935	6.910	
	3 (Sewaged)	14	5	70	14	4	39.996	10.871	0.25.	27.47	27.47	2.757	2.757	10.036	10.036	
	0* (Unsewaged)	9	2½	22½	9	14	61.404	11.227	0.25.	18.28	18.28	1.871	1.871	10.599	10.235	
• Unmeasured land; designated Plot 0 when unsewaged, and Plot 00 when sewaged.																
Rye-grass	1 (Unsewaged)	8	2½	20	8	14	28.481	10.396	0.25.	36.50	36.50	2.940	2.940	8.080	8.085	
	2 (Sewaged)	11	5	53	17	6	28.481	10.382	0.25.	36.45	36.45	2.956	2.956	8.110	8.085	
	3 (Sewaged)	14	5	70	14	4	61.404	11.225	0.25.	18.28	18.28	1.871	1.871	9.870	9.870	
	0* (Unsewaged)	9	2½	22½	9	14	61.404	11.225	0.25.	21.15	21.15	2.480	2.478	11.727	11.736	

• Unmeasured land; designated Plot 0 when unsewaged, and Plot 00 when sewaged.

and Rye-grass.

Third Season, 1863.

Determinations of Dry Substance and Mineral Matter (Ash).

Determinations of Dry Substance and Mineral Matter (Ash).														
Particulars of Sampling.				Actual Weights.				Per-centages.						
Fields.	Plots.	Number of Samples taken.	Weights.			Air-dried.	Equal in Fresh State.	Dry Sub-stance Matter (at 212° Fahr.)	Dry in Fresh.		Ash in Fresh.		Ash in Dry.	
			In fresh state.	Air-dried.					Each Expert.	Mean.	Each Expert.	Mean.	Each Expert.	Mean.
FIFTH CROP.*														
Five-acre	4 (Sewaged)	1	lbs. 20	lbs. 3 11	ozs. 12.461 { 12.461	ozs. 67.576 10.308	ozs. 1.352 1.352	15.25 15.25	15.25 15.25	2.001 { 1.971	1.986 { 1.986	13.116 { 13.087	13.023	
	{ 1 (Unsewaged) 2 (Sewaged) 3 (Sewaged)	1	20	4 11	50 { 12.5	53.328 53.328	10.620 10.627	1.562 1.570	19.91 19.93	19.92 19.93	2.929 2.944	2.937 2.944	14.708 14.774	
		2	5	10 2 04	324 { 8.063	39.995 39.995	6.717 6.717	0.874 0.884	16.79 16.75	16.75 16.75	2.186 2.186	2.186 2.186	13.012 13.012	
Rye-grass	{ 2 (Sewaged) 3 (Sewaged)	5	5	5 6	50 { 12.5	58.140 58.140	10.845 10.845	1.270 1.270	17.79 17.72	17.76 17.76	2.184 2.187	2.176 2.176	12.956 12.954	
		SIXTH CROP. †												
Rye-grass	{ 2 (Sewaged) 3 (Sewaged)	2	10	4 6	50 { 12.5	57.143 57.143	10.665 10.642	1.466 1.460	18.66 18.62	18.64 18.62	2.618 2.607	2.618 2.607	14.014 14.001	
		3	10	4 134	50 { 12.5	51.610 51.610	10.598 10.598	1.465 1.465	20.36 20.36	20.38 20.38	2.941 2.937	2.941 2.937	14.481 14.481	

* taken from Plots 2 and 3 in the 5-acre field, from Plots 3 and 4 in the 10-acre field, and from Plot 0 in the rye-grass field, but very late in the season, and they and some of them were so wet that the analyses were not proceeded with.

† taken from Plots 2 and 3 in the 5-acre field, from Plots 3 and 4 in the 10-acre field, and from Plot 0 in the rye-grass field, but very late in the season, and they and some of them were so wet that the analyses were not proceeded with.

TABLE XXX.

SUMMARY of the Percentages of Dry Substance (at 212° Fahr.), and of Mineral Matter (Ash), in the Unsewaged and the Sewaged Meadow Grass, Second Season, 1862.

Plots.	PER-CENTAGES.											
	Dry Substance in Fresh Grass.				Mineral Matter in Fresh Grass.				Mineral Matter in Dry Substance.			
	1st Crop.	2d Crop.	3d Crop.	4th Crop.	1st Crop.	2d Crop.	3d Crop.	4th Crop.	1st Crop.	2d Crop.	3d Crop.	4th Crop.
FIVE-ACRE FIELD.												
1 (Unsewaged)	28.7	23.8	2.41	2.03	9.02	8.99
2 (Sewaged)	22.8	14.3	18.2	..	1.92	1.72	2.42	..	8.44	12.00	13.25	..
3 (Sewaged)	14.4	16.4	12.9	..	1.39	1.70	1.81	..	9.65	10.38	14.03	..
4 (Sewaged)	15.3	19.4	14.2	..	1.42	2.02	1.80	..	9.32	10.39	12.67	..
TEN-ACRE FIELD.												
1 (Unsewaged)	26.9	17.9	2.62	2.09	9.38	11.68
2 (Sewaged)	19.5	16.2	14.5	..	1.84	1.86	1.83	..	9.40	11.51	12.62	..
3 (Sewaged)	13.5	19.0	14.4	..	1.54	1.62	1.73	..	11.37	10.12	11.99	..
4 (Sewaged)	13.1	16.7	15.8	33.8	1.63	1.69	1.91	4.39	12.44	10.07	12.08	13.99

**SUMMARY of the Percentages of Dry Substance (at 212° Fahr.), and of Mineral Matter (Ash), in the Unsewaged and the Sewaged Meadow Grass and Rye-grass,
Third Season, 1863.**

Plots.	PER-CENTAGES.																	
	Dry Substance in Fresh Grass.						Mineral Matter in Fresh Grass.						Mineral Matter in Dry Substance.					
	1st Crop.	2d Crop.	3d Crop.	4th Crop.	5th Crop.	6th Crop.	1st Crop.	2d Crop.	3d Crop.	4th Crop.	5th Crop.	6th Crop.	1st Crop.	2d Crop.	3d Crop.	4th Crop.	5th Crop.	6th Crop.
FIVE-ACRE FIELD.																		
1 (Unsewaged)	-	36.1	34.4	2.88	3.82	7.97	11.10
2 (Sewaged)	-	21.5	18.5	17.7	15.8	..	1.97	2.00	2.05	1.88	9.19	10.79	11.58	11.91
3 (Sewaged)	-	17.6	14.9	10.9	13.0	..	1.78	1.56	1.37	1.69	10.13	10.49	12.59	12.96
4 (Sewaged)	-	16.3	17.8	17.0	12.3	15.3	1.60	1.50	2.02	1.45	1.99	..	10.23	8.77	11.51	11.76	13.02	..
TEN-ACRE FIELD.																		
1 (Unsewaged)	-	39.8	18.2	2.90	2.24	7.29	12.34
2 (Sewaged)	-	18.6	17.7	12.4	1.55	1.03	1.45	8.33	10.98	11.67
3 (Sewaged)	-	20.0	16.3	14.6	13.9	..	2.00	1.83	1.73	1.74	9.98	11.20	11.89	12.53
4 (Sewaged)	-	14.6	18.8	15.2	13.6	..	1.53	1.94	1.76	1.70	10.64	10.28	11.67	12.46
RYE-GRASS.																		
1 (Unsewaged)	-	21.3	23.7	36.4	33.2	19.9	1.95	1.91	2.51	3.80	2.94	..	9.16	8.07	6.91	11.44	14.74	..
2 (Sewaged)	-	..	18.8	27.5	13.8	16.8	..	1.55	2.76	1.37	2.20	2.61	..	8.38	10.04	9.91	13.12	14.01
3 (Sewaged)	-	..	17.5	18.3	18.9	17.8	20.4	..	1.67	2.09	2.18	2.94	..	9.51	10.24	11.08	12.25	14.43

* No. Sewason was applied for the 1st Crop of Plots 2 and 3, and no Samples were taken from the produce.

* No Sewage was applied for the 1st Crop of Plots 2 and 3, and no Samples were taken from the produce.

TABLE XXVIII.
Growing the Composition (per cent.) of the Dry Substance of the Unsewaged and the Sewaged Meadow Grass.
Second Season, 1862.

[illegible]

• Designated Plot 0 when unmeasured land.

TABLE XXXIV.
Showing the Composition (per cent.) of the Unsewaged and the Sewaged Meadow Grass in the *Fresh State*, as weighed.
Third Season, 1863.

	First Crop.				Second Crop.				Third Crop.				Fourth Crop.				Fifth Crop.
	Un-sewaged.		Sewaged.		Un-sewaged.		Sewaged.		Sewaged.		Sewaged.		Sewaged.		Sewaged.		Sewaged.
	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 2.	Plot 3.	Plot 4.	Plot 2.	Plot 3.	Plot 4.	Plot 2.	Plot 3.	Plot 4.
FIVE-ACRE FIELD.																	
Nitrogenous substance ($N \times 6.3$) -	2.75	2.93	3.03	3.95	4.44	3.07	3.04	3.15	3.30	2.46	3.82	4.03	3.30	3.15	4.03	3.15	4.01
Fatty matter (ether extract) -	1.14	1.09	0.98	0.80	1.48	0.98	0.56	0.70	0.93	0.58	0.88	0.59	0.60	0.61	0.59	0.61	0.77
Woody fibre -	10.92	5.38	4.10	3.60	8.87	5.24	4.32	5.37	4.41	2.80	4.38	3.41	3.15	3.00	3.41	3.15	3.13
Other non-nitrogenous substances	18.65	10.11	7.72	6.07	15.74	7.04	5.41	6.59	6.93	3.09	6.42	5.27	4.31	3.97	4.31	3.97	4.53
Mineral matter (ash) -	2.64	1.98	1.77	1.78	3.89	2.18	1.58	1.79	2.03	1.39	2.08	1.85	1.59	1.37	1.85	1.59	1.92
Total dry substance -	36.10	21.49	17.59	16.26	34.42	18.51	14.91	17.80	17.67	10.92	17.58	15.75	13.04	12.30	15.75	13.04	15.25
Water -	63.90	78.51	82.41	83.74	65.58	81.49	85.09	82.20	82.33	89.08	82.42	84.25	86.96	87.70	84.25	86.96	84.75
100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
TEN-ACRE FIELD.																	
Nitrogenous substance ($N \times 6.3$) -	3.50	2.40	3.19	2.97	2.17	3.31	3.15	2.92	2.36	2.54	3.02	—	3.54	3.38	—	—	—
Fatty matter (ether extract) -	1.72	0.94	0.87	0.83	0.83	0.96	0.62	0.65	0.66	0.65	0.70	—	0.70	0.75	—	—	—
Woody fibre -	12.07	5.52	5.17	3.43	4.72	4.47	3.24	5.06	3.24	3.90	4.26	—	4.26	3.21	—	—	—
Other non-nitrogenous substances	18.65	8.13	8.81	5.81	8.16	6.66	6.12	7.63	4.90	5.79	5.38	—	4.75	4.55	—	—	—
Mineral matter (ash) -	2.95	1.57	1.99	1.02	2.30	2.11	1.89	1.96	1.50	1.70	1.81	—	1.71	1.73	—	—	—
Total dry substance -	39.79	18.56	20.03	14.56	18.17	17.66	16.25	18.82	12.42	14.58	15.17	—	13.87	13.62	—	—	—
Water -	60.21	81.44	79.97	85.44	81.83	82.34	83.75	81.18	87.58	85.42	84.83	—	86.13	86.38	—	—	—
100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	—	100.00	100.00	—	—	—

TABLE XXXV.
Unsewaged and the Sewaged Meadow Grass.
Showing the Composition (per cent.) of the Dry Substance of the
Third Season, 1868.

	First Crop.				Second Crop.				Third Crop.				Fourth Crop.				Fifth Crop.
	Un-sewaged.		Sewaged.		Un-sewaged.		Sewaged.		Sewaged.		Sewaged.		Sewaged.		Sewaged.		Sewaged.
	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 4.
FIVE-ACRE FIELD.																	
Nitrogenous substance (N x 6.3)	7.62	13.71	17.33	24.31	12.88	16.57	20.38	17.71	18.66	22.00	21.71	29.30	25.27	25.59	32.19		
Fatty matter (ether extract)	8.16	5.09	5.40	5.27	4.31	5.29	3.75	3.91	5.38	5.30	4.98	3.77	5.33	4.96	5.07		
Woody fibre	30.24	25.02	23.34	22.13	25.78	28.31	23.94	31.32	24.97	25.02	24.94	21.68	24.15	24.42	20.51		
Other non-nitrogenous substances	51.66	47.06	43.88	37.35	45.73	38.07	36.31	37.02	39.33	33.76	36.51	33.43	33.02	32.24	29.68		
Mineral matter (ash)	7.32	9.12	10.05	10.94	11.30	11.76	10.62	10.04	11.66	12.72	11.86	11.76	12.23	12.79	12.61		
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00		
TEN-ACRE FIELD.																	
Nitrogenous substance (N x 6.3)	8.81	12.93	15.04	19.75	11.94	18.73	19.42	15.49	18.98	17.46	10.87	—	25.53	24.82	—		
Fatty matter (ether extract)	4.32	5.04	4.33	5.69	4.52	5.46	3.79	3.46	5.28	4.46	4.02	—	5.07	5.53	—		
Woody fibre	32.59	29.74	25.83	23.54	25.95	26.72	27.50	30.08	26.12	26.76	28.07	—	22.85	23.59	—		
Other non-nitrogenous substances	46.87	43.81	43.97	39.92	44.91	37.17	37.66	40.54	37.53	39.63	35.49	—	34.25	33.38	—		
Mineral matter (ash)	7.41	8.46	9.93	11.10	12.68	11.92	11.63	10.43	12.09	11.64	11.95	—	12.32	12.68	—		
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	—	100.00	100.00	—		

TABLE XXXVI.
SHOWING the Composition (per cent.) of the Unsewaged and the Sewaged Italian Rye-grass in the *Fresh State*, as weighed.
Season 1863.

	First Crop.		Second Crop.		Third Crop.		Fourth Crop.		Fifth Crop.		Sixth Crop.	
	Un-sewaged.		Un-sewaged.		Un-sewaged.		Un-sewaged.		Un-sewaged.		Un-sewaged.	
	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.
Nitrogenous substance ($N \times 6.3$) -	2.67	3.26	3.05	3.33	2.32	2.30	4.05	2.30	3.63	3.87	4.60	5.06
Fatty matter (ether extract) -	0.77	0.72	1.10	0.69	0.69	0.77	1.18	0.77	0.87	0.81	0.83	1.19
Woody fibre -	3.79	4.40	10.36	7.70	5.40	3.83	8.33	3.83	4.61	4.20	4.22	5.43
Other non-nitrogenous substances	12.05	8.49	10.24	12.77	8.61	5.46	14.97	5.46	7.87	6.62	6.33	6.68
Mineral matter (ash) -	2.02	1.85	2.56	2.08	1.31	1.41	3.78	1.41	2.94	2.25	2.66	3.09
Total dry substance -	21.30	18.81	30.35	27.47	18.28	13.84	35.21	13.84	19.02	16.76	18.64	20.38
Water -	78.70	81.19	63.65	72.53	81.72	86.16	66.79	86.16	80.08	83.25	81.36	79.62
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

TABLE XXXVII.
SHOWING the Composition (per cent.) of the *Dry Substance* of the Unsewaged and the Sewaged Italian Rye-grass.
Season 1863.

	First Crop.		Second Crop.		Third Crop.		Fourth Crop.		Fifth Crop.		Sixth Crop.	
	Un-sewaged.		Un-sewaged.		Un-sewaged.		Un-sewaged.		Un-sewaged.		Un-sewaged.	
	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.	Plot 1.	Plot 2.
Nitrogenous substance ($N \times 6.3$) -	12.51	8.19	17.33	13.07	8.38	13.12	12.70	14.02	18.22	23.11	24.70	24.83

TABLE XXXVIII.
Average Composition (per cent.) of the Meadow Grass from each Plot, and of each successive Crop.
Second Season, 1862.

		MEAN COMPOSITION, PER CENT.															
		Of the Fresh Grass as weighed.								Of the Dry Substance of the Grass.							
		Without, or with different Quantities of Sewage.				In each successive Crop.				Without, or with different Quantities of Sewage.				In each successive Crop.			
		Un-sewaged.		Sewaged.		1st Crop.		2d Crop.		Un-sewaged.		Sewaged.		1st Crop.		2d Crop.	
		Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Plot 1.	Plot 2.	Plot 3.	Plot 4.
FIVE-ACRE FIELD.																	
Number of analyses	-	2	3	3	3	4	4	4	3	2	3	3	3	4	4	3	1
Nitrogenous substance (N x 6.3)	-	1.97	2.87	2.23	2.66	2.01	2.28	2.01	3.37	7.01	16.36	15.53	16.65	10.65	13.11	22.16	-
Fatty matter (ether extract)	-	0.06	0.65	0.52	0.60	0.53	0.64	0.65	0.65	2.07	3.67	3.55	3.73	2.70	3.62	4.59	-
Woody fibre	-	7.60	5.44	4.34	4.73	6.30	5.44	4.03	-	31.12	29.49	29.72	29.07	31.99	29.81	28.76	-
Other non-nitrogenous substances	-	12.06	7.44	5.88	6.62	9.16	8.00	5.10	-	43.02	39.42	40.02	40.31	45.79	43.10	33.84	-
Mineral matter (ash)	-	2.33	1.99	1.60	1.66	1.78	1.86	1.96	-	9.38	11.06	11.16	10.24	8.97	10.36	12.96	-
Total dry substance	-	24.71	18.43	14.57	16.29	19.78	18.22	15.11	-	-	-	-	-	-	-	-	-
Water	-	75.29	81.57	85.43	83.71	80.22	81.78	84.89	-	100.00	100.00	100.00	100.00	100.00	100.00	100.00	-
	-	100.00	100.00	100.00	100.00	100.00	100.00	100.00	-	100.00	100.00	100.00	100.00	100.00	100.00	100.00	-
TEN-ACRE FIELD.																	
Number of analyses	-	2	3	3	3	4	4	4	3	1	2	3	3	4	4	3	1
Nitrogenous substance (N x 6.3)	-	2.42	2.43	2.41	3.38	2.49	2.10	2.10	2.79	6.16	11.08	14.94	15.85	16.97	14.93	12.06	18.23
Fatty matter (ether extract)	-	0.09	0.64	0.58	0.80	0.55	0.65	0.65	0.65	1.49	3.19	3.95	3.72	3.95	3.14	3.77	4.42
Woody fibre	-	6.47	4.90	4.47	5.74	6.02	4.00	3.98	8.40	28.47	28.92	28.02	28.65	33.69	28.09	28.63	24.86
Other non-nitrogenous substances	-	10.51	7.10	6.49	7.68	7.67	7.91	5.70	13.08	40.77	41.08	40.90	38.03	40.04	45.50	38.16	38.70
Mineral matter (ash)	-	2.29	1.66	1.68	2.27	1.53	1.88	1.80	4.67	10.40	10.21	10.70	10.60	8.50	10.78	12.10	13.60
Total dry substance	-	22.38	16.73	15.63	19.87	18.26	17.44	14.92	33.80	-	-	-	-	-	-	-	-
Water	-	77.62	83.27	84.37	80.13	81.74	82.56	85.03	66.20	-	-	-	-	-	-	-	-
	-	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	-	-	-	-	-	-	-	-

TABLE XXXIX.
Average Composition (per cent.) of the Meadow Grass from each Plot, and of each successive Crop.
Third Season, 1863.

MEAN COMPOSITION, PER CENT.																																					
Of the Fresh Grass as weighed.										Of the Dry Substance of the Grass.																											
Without, or with different Quantities of Sewage.										Without, or with different Quantities of Sewage.																											
Un-sewaged.					Sewaged.					Un-sewaged.					Sewaged.																						
Plot 1.		Plot 2.		Plot 3.		Plot 4.		1st Crop.		2d Crop.		3d Crop.		4th Crop.		5th Crop.		1st Crop.		2d Crop.		3d Crop.		4th Crop.		5th Crop.											
FIVE-ACRE FIELD.																																					
Number of analyses		-		2		4		4		4		4		3		3		1		2		4		4		5		4		3		3		1			
Nitrogenous substance (N x 8.3)		-		3.60		3.49		2.96		3.80		3.17		3.42		3.19		3.09		4.01		10.25		19.53		21.40		24.30		15.74		16.89		20.99		33.19	
Fatty matter (ether extract)		-		1.31		0.90		0.70		0.76		1.01		0.68		0.80		0.63		0.77		3.78		4.88		4.94		4.84		4.73		4.31		5.22		4.68	
Woody fibre		-		9.80		4.61		3.59		3.94		6.00		3.86		3.19		3.13		3.13		29.01		24.99		25.51		24.66		25.18		28.59		25.19		20.51	
Other non-nitrogenous substances		-		17.20		7.34		5.28		5.51		10.64		8.70		5.69		4.52		4.52		43.70		39.47		36.74		34.55		41.90		39.28		36.53		32.90	
Mineral matter (ash)		-		3.26		2.01		1.93		1.83		2.04		2.30		1.85		1.67		1.92		9.31		11.08		11.41		11.65		9.36		10.83		12.06		12.61	
Total dry substance		-		35.26		18.35		14.11		15.84		22.83		21.41		15.39		13.70		15.25		100.00		100.00		100.00		100.00		100.00		100.00		100.00		100.00	
Water		-		64.74		81.65		85.98		84.16		77.14		78.59		84.61		80.30		84.75		100.00		100.00		100.00		100.00		100.00		100.00		100.00		100.00	
TEN-ACRE FIELD.																																					
Number of analyses		-		2		3		4		4		4		4		3		2		-		2		3		4		4		4		3		2		-	
Nitrogenous substance (N x 8.3)		-		2.83		2.69		3.10		3.05		2.99		2.80		2.64		3.46		-		10.33		16.89		19.59		19.08		14.36		16.39		18.77		25.17	
Fatty matter (ether extract)		-		1.27		0.85		0.71		0.73		1.00		0.76		0.67		0.73		-		4.42		5.26		4.41		4.83		4.84		4.31		4.79		5.30	
Woody fibre		-		8.94		4.19		4.18		4.14		6.77		4.89		3.80		3.19		-		29.27		27.53		25.73		24.82		27.08		27.56		26.08		23.21	
Other non-nitrogenous substances		-		13.41		6.46		6.37		5.84		10.35		7.12		5.28		4.65		-		45.89		39.50		38.89		37.83		43.64		40.07		37.57		33.82	
Mineral matter (ash)		-		2.63		1.73		1.98		1.78		2.03		2.07		1.67		1.72		-		10.04		10.82		11.38		11.64		9.23		11.07		11.89		12.60	
Total dry substance		-		28.96		16.21		16.18		15.54		23.23		17.73		14.00		13.76		-		100.00		100.00		100.00		100.00		100.00		100.00		100.00		100.00	
Water		-		71.06		83.79		83.82		84.46		76.77		82.27		85.94		86.24		-		100.00		100.00		100.00		100.00		100.00		100.00		100.00		100.00	

TABLE XL.

AVERAGE Composition (per cent.) of the Italian Rye-grass from each Plot, and of each successive Crop.

SEASON 1863.

		MEAN COMPOSITION, PER CENT.																																			
		Of the Fresh Grass, as weighed.												Of the Dry Substance of the Grass.																							
		Without, or with different Quantities of Sewage.						In each successive Crop.						Without, or with different Quantities of Sewage.						In each successive Crop.																	
		Un-sewaged.		Sewaged.				1st Crop.		2d Crop.		3d Crop.		4th Crop.		5th Crop.		6th Crop.		Un-sewaged.		Sewaged.		1st Crop.		2d Crop.		3d Crop.		4th Crop.		5th Crop.		6th Crop.			
		Plot 1.		Plot 2.		Plot 3.		Plot 1.		Plot 2.		Plot 3.		Plot 1.		Plot 2.		Plot 3.		Plot 1.		Plot 2.		Plot 3.		Plot 1.		Plot 2.		Plot 3.		Plot 1.		Plot 2.		Plot 3.	
Number of Analyses -		5	6	5			1	3	3	3	3	3	3	3	3	3	3	3	5	5	5	1	3	3	3	3	3	3	3	3	3	3	3	3			
Nitrogenous substance } (N x 6.8)		3.25	3.47	3.39			2.67	2.49	2.90	3.58	3.77	4.83	12.44	13.78	13.11	12.51	12.86	11.07	16.70	20.91	24.76																
Fatty matter (ether)		0.93	0.83	0.73			0.77	0.61	0.99	0.89	0.81	1.01	3.53	4.45	3.95	3.61	3.01	3.28	4.31	4.49	5.13																
Woody fibre -		6.79	4.90	4.73			3.79	5.43	7.84	5.98	4.10	4.33	24.08	25.51	25.55	17.79	26.95	23.79	27.16	22.64	23.16																
Other non-nitrogenous } substances -		13.39	7.74	7.63			12.05	9.72	13.55	9.21	6.98	6.47	49.21	39.76	41.47	56.59	46.23	46.88	40.87	33.34	33.24																
Mineral matter (ash) -		2.64	2.17	2.07			2.02	1.77	2.19	2.44	2.46	2.88	10.14	11.50	11.03	9.50	8.90	7.98	10.96	13.63	14.71																
Total dry substance		26.90	19.10	18.57			21.30	20.01	27.37	21.98	18.14	19.51																									
Water -		73.10	80.90	81.43			78.70	79.99	72.63	78.03	81.86	80.49	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00																
		100.00	100.00	100.00			100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00																

Season 1862.

[illegible]

TABLE XLII.
Details of the Sampling, and of the Determinations of Dry Substance, and Mineral Matter (Ash), in the Grass grown in the Fourth Season, 1864,
without Sewage.

without Sewage.

Particulars of Sampling.

Determinations of Dry Substance, and Mineral Matter (Ash).

Fields.	Plots.	How treated in 1861, 1862, and 1863.	Number of Samples taken.	Weights.			Actual Weights.	Per-centages.											
				In fresh state.		Air-dried.		Dry in Fresh.	Ash in Fresh.	Ash in Dry.									
				Each Sample.	Total.														
First Crop.		lbs.	lbs.	lbs. ozs.	ozs.	ozs.	ozs.	Each Exper.	Mean.	Each Exper.	Mean.								
Five-acre	1	Unsewaged	2	10	20	8	6	50	13.5	29.851	10.392	0.764	34.51	34.70	2.559	2.579	2.509	7.353	7.408
	2	Sewaged	4	5	20	7	5	50	13.5	29.851	10.323	0.770	34.53	30.19	2.311	2.339	2.325	7.459	7.708
	3	Sewaged	4	5	20	6	14	50	13.5	34.186	10.312	0.800	30.16	28.19	2.339	2.317	2.131	7.753	7.580
	4	Sewaged	4	5	20	5	0	50	12.5	36.364	10.199	0.760	28.06	20.31	2.145	1.890	1.810	8.945	8.904
Ten-acre	1	Unsewaged	2	10	20	8	1	50	12.5	31.008	10.348	0.714	33.37	33.36	2.395	2.274	2.274	6.900	8.915
	2	Sewaged	2	10	20	7	8	50	12.5	31.008	10.344	0.688	33.38	30.76	2.245	2.370	2.376	6.729	7.705
	3	Sewaged	4	5	20	5	12	50	12.5	33.353	10.232	0.790	30.92	23.63	2.382	1.918	1.907	7.704	8.072
	4	Sewaged	4	5	20	5	5	50	12.5	43.478	10.312	0.824	23.53	21.43	1.985	1.908	1.905	8.055	8.982
Second Crop.		lbs.	lbs.	lbs. ozs.	ozs.	ozs.	ozs.	Each Exper.	Mean.	Each Exper.	Mean.								
Five-acre	4	Sewaged	1	20	20	13	7	50	12.5	18.605	10.337	0.803	53.56	55.53	4.316	4.322	4.322	7.708	7.730
									12.5	18.605	10.325	0.794	55.49		4.268			7.682	

TABLE XLIII.
Results of the Mechanical Analysis of the Rugby Soils.
Composition per Cent.

	Plot 1.	Plot 2.	Plot 3.	Plot 4.	Mean.
FIVE-ACRE FIELD.					
Stones retained by 1 inch sieve -	5.59	6.95	7.49	6.01	6.51
Stones passing 1 inch, and re- tained by $\frac{1}{2}$ inch sieve - - }	9.48	7.33	7.43	6.18	7.60
Stones passing $\frac{1}{2}$ inch, and re- tained by $\frac{1}{4}$ inch sieve - - }	6.04	4.44	3.58	3.36	4.35
Total stones -	21.11	18.72	18.50	15.55	18.46
Mould passing $\frac{1}{4}$ inch sieve -	77.77	80.36	79.80	83.29	80.31
Roots, loss by evaporation, &c. -	1.12	0.92	1.70	1.16	1.23
Total fresh mould -	78.89	81.28	81.50	84.45	81.54
	100.00	100.00	100.00	100.00	100.00
TEN-ACRE FIELD.					
Stones retained by 1 inch sieve -	1.54	2.47	3.66	2.90	2.64
Stones passing 1 inch, and re- tained by $\frac{1}{2}$ inch sieve - - }	1.92	2.18	2.43	3.68	2.55
Stones passing $\frac{1}{2}$ inch, and re- tained by $\frac{1}{4}$ inch sieve - - }	3.04	1.72	1.98	3.43	2.54
Total stones -	6.50	6.37	8.07	10.01	7.73
Mould passing $\frac{1}{4}$ inch sieve -	92.65	91.25	90.36	89.13	90.85
Roots, loss by evaporation, &c. -	0.85	2.38	1.57	0.86	1.42
Total fresh mould -	93.50	93.63	91.93	89.99	92.27
	100.00	100.00	100.00	100.00	100.00

The mode of collecting the samples was as follows:—A strong iron frame, in the form of the four sides of a box without either top or bottom, 12 inches square by 9 inches deep, was driven into the ground as much more than 9 inches as to allow a very thin sod (only just thick enough to insure the removal of the whole of the green matter) to be taken off the top of it. The contents of the frame were then carefully dug out; and four such samples of 12 inches square by 9 inches deep were taken from each plot, and mixed together. The soil of each plot was thus represented by a sample, averaging, in the case of the Five-acre Field rather over, and in that of the Ten-acre Field somewhat under, 300 lbs.

TABLE XLIV.

Results of the Chemical Analysis of the Ragby Soils.

Per-centages of Water, Organic Matter, Nitrogen, and Nitrogen reckoned as Ammonia.

Pots.	Water per Cent.			Organic Matter per Cent.			Nitrogen* per Cent.						Nitrogen reckoned as Ammonia. Mean, per Cent.		
	In Fresh Mould as analysed.	In Total Fresh Soil (including Stones, &c.)	In Fresh Mould.	In Dry Mould.	In Total Fresh Soil (including Stones, &c.)	In Fresh Mould.	In Fresh Mould.			In Dry Mould.	In Total Fresh Soil (including Stones, &c.)	In Fresh Mould.	In Dry Mould.	In Total Fresh Soil (including Stones, &c.)	
							Exp. 1.	Exp. 2.	Exp. 3.	Exp. 4.	Mean.	Mean.	Mean.	Mean.	

FIVE-ACRE FIELD.															
1	14.65	11.39	5.30	6.21	4.12	0.192	0.191	0.191	0.191	0.191	0.224	0.146	0.233	0.273	0.180
2	15.59	13.53	5.80	6.87	4.66	0.195	0.191	0.197	0.204	0.197	0.233	0.166	0.239	0.283	0.192
3	15.22	13.15	6.06	7.15	4.84	0.187	0.183	0.190	0.185	0.185	0.218	0.146	0.235	0.265	0.180
4	14.63	12.23	5.05	5.92	4.21	0.182	0.181	0.186	0.184	0.183	0.214	0.133	0.222	0.260	0.185
Mean -	15.04	12.08	5.55	6.54	4.46	0.189	0.187	0.191	0.190	0.189	0.223	0.152	0.230	0.270	0.184

TEN-ACRE FIELD.															
1	20.66	19.14	9.10	11.47	8.45	0.264	0.258	0.257	0.252	0.258	0.325	0.239	0.313	0.335	0.290
2	18.61	16.96	9.80	12.04	8.94	0.243	0.247	0.236	0.246	0.243	0.299	0.223	0.295	0.303	0.270
3	17.60	15.90	8.20	9.95	7.41	0.240	0.236	0.226	0.231	0.233	0.283	0.211	0.283	0.344	0.266
4	17.56	15.65	9.20	11.16	8.20	0.254	0.247	0.252	0.249	0.251	0.304	0.224	0.305	0.369	0.272
Mean -	18.61	16.93	9.08	11.16	8.25	0.250	0.247	0.245	0.245	0.246	0.303	0.224	0.299	0.368	0.272

* Nitrogen was determined by combustion with soda lime, and estimation as ammonia in the usual volumetric way. The results do not, therefore, represent the total nitrogen, including that in the form of nitric acid, if any, excepting so far as it might be reduced by the organic matter present. Further, as the note at the foot of Table XLIII. will show, the figures only relate to the composition of the layer of 9 inches in thickness taken immediately below a very thin sod.

APPENDIX No. 2.

NOTES ON THE EDINBURGH SEWAGE MEADOWS.

The following statements relating to the sewage-irrigated meadows in the neighbourhood of Edinburgh are based partly upon information obtained on the spot from Mr. Mc Pherson the city surveyor, and from the occupiers or managers of the meadows in the respective localities and partly upon correspondence with the latter gentlemen. Mr. Mc Pherson kindly provided information as to the population, water supply, springs, and area of rainfall and drainage, contributing to each set of meadows as named or classed together in Table I. below ; and from the occupiers or managers, information as to the area under irrigation, and other points, was obtained. As, however, the sewage is in no case passed over the land the whole 365 days and nights of the year, it must be borne in mind that the statements given as to the amount of population contributing to, and the number of tons available for, each acre, do not show the amounts actually utilized in each case, but only approximately the total amounts available, whether used or wasted. The fact that water-closets are not universal must also be taken into account.

TABLE I.

Names of Meadows.	Imperial Acres under Irrigation.	Approximate Population contributing to each Acre.	Approximate Quantity of Sewage available for each Acre
			Tons.
Lochend, Spring Gardens, and Craigentenny	285	337	20,500
Roseburn and Western Dalry - -	80	112	17,000
Quarry Holes - - - -	8	562	65,000
Broughton Burn - - - -	6	1,666	102,000
The Grange - - - -	16½	302	97,000

1. Lochend and Spring Gardens.

The Lochend and Spring Gardens meadows, occupied by Mr. Scott and under the management of Mr. Peter Taylor, comprise about 285 imperial acres, and are irrigated by the sewage of a large portion of Edinburgh on its way to the more extensive Craigentenny meadows. Mr. Taylor estimates that, on the average, each acre gets the flow of a stream of 12 by 8 inches, at a rate of 2 miles per hour, for 10 days 16 hours, annually, which is equal to about 31,000 tons per acre per annum. The sewage is applied from about the beginning of March to the end of November. The herbage consists chiefly of rough meadow grass (*Poa trivialis*), which is the most prevalent where the ground is dry, and couch (*Triticum repens*), which is the second in predominance and grows freely, but is not so early as the rough meadow-grass. There is also a great deal of crow-foot, more particularly where the drainage is imperfect.

On a portion of higher lying land, which is irrigated by the aid of a water-wheel, worked by the sewage stream itself, and where the supply is necessarily more limited, Italian rye-grass is grown, which involves the periodical breaking up of the land. After two years under the rye-grass a crop of potatoes is taken, then Italian rye-grass is sown again and so on. Mr. Taylor stated that if this land could be irrigated by

abundance of sewage by gravitation he should prefer to lay it down as permanent meadow.

2. Craigentinny.

The meadows at Craigentinny, which are the property of Mr. Christie-Miller, are under the management of Mr. Bryce. About 190 acres of permanent meadow are irrigated by gravitation. A portion of this area consists of a good loamy soil, but a part of it was only barren sand before it was laid down for sewage irrigation. During the summer the sewage is applied day and night, and all that is available is then used, excepting during floods. The sewage is also applied through a considerable portion of the winter, but then during the day only. Perhaps it is unused 70 to 80 days in the year. The general plan during the summer is to let the whole of the water go over from 2 to $2\frac{1}{2}$ acres at a time, changing every 3 or 4 hours during the day, but less frequently during the night; and the application is so timed as to get over, on the average, about 60 acres per week, and to give each acre such a dressing about once in three weeks. The applications are, however, less frequent during the winter. The distribution over about 100 acres can be attended to by one man; but the cleaning of the runs, keeping the roads, &c., require additional labour. Four to five crops are obtained annually; though four, cut at the proper times, generally yield better, and leave the herbage in better condition than when five are taken. From good, well managed meadows, with sewage as liberally applied as on the gravitation meadows at Craigentinny, Mr. Bryce thinks about 60 tons of green grass should be obtained per imperial acre annually. The price varies, according to season and other circumstances, from 6*d.* to 14*d.* per cwt., on the ground, standing. The produce consists almost entirely of rough meadow-grass, which is considered the most valuable, couch, which is looked upon as a very good grass and of very rapid growth, and common rye-grass, which is also considered a good grass, but not to give so close a bottom as the others.

Arrangements are also made for irrigating some higher lying land, by raising the sewage about 20 feet, it being first brought into a large tank by means of a deep underground drain from the highest level of the natural flow, and thence pumped into open channels for surface distribution. Only about 60 imperial acres are now so irrigated, but formerly a larger area was under the treatment. The application is continued from April to October inclusive; and each plot gets six dressings, and yields three cuttings, annually. If it were not for the cost of lifting, more of this land would be laid down as permanent meadow, and much more sewage would be put upon it; but the supply being so limited by the cost of application, Mr. Bryce thinks it better to sow Italian rye-grass, break up every two years and grow potatoes, re-sow rye-grass, and so on.

3. Roseburn and Western Dalry.

These meadows, situated to the west of the city, stand second in point of extent and importance to those of Craigentinny. They formerly comprised 100 imperial acres or more, but are now limited to about 80, having been considerably curtailed by the encroachments of railways, and for other purposes; and as much as 1,000*l.* per acre has frequently been paid as compensation. They were laid down for irrigation by Mr. James Thomson, who commenced his operations in 1826, and they still continue under his excellent management. Part of the soil is gravelly, and part loamy, with a subsoil of clay. The sewage

coming to these meadows includes the refuse from extensive slaughter-houses, and also that from a very large distillery. It is used all the year round, both day and night, and on Sundays when it can conveniently be left to flow from Saturday night to Monday morning. In summer the water seldom goes over the same piece of land more than a few hours together—as long as may be necessary thoroughly to soak it. The land is generally watered only twice, but occasionally three times, between the cuttings. In winter the water is allowed to flow for a longer period over a given area, in order to “feed” the land as much as possible. In laying down permanent meadow for irrigation, Mr. Thomson has sometimes sown a great variety of grasses, but finding that even when he has sown 15 or 20 different kinds, most of them have in the course of a few years gradually died out, leaving only those suitable to the soil and the sewage, he sometimes selects only three or four, and in one instance transplanted couch from arable land, and reports that the piece so treated is now as good as any.

4. *The Grange.*

Next in extent to the Roseburn and Dalry sewage meadows come those of the Grange, situated to the south of the city, and farmed by Mr. Mc Gill. The area of these meadows is at present only $16\frac{1}{2}$ imperial acres ; but it is less now than formerly, having been contracted for want, as was stated, of a sufficient supply of sewage. Indeed, the opinion given by Mr. Mc Gill, junior, was, that the greater the amount of water applied the better. The Table (p. 198) shows that the population within the area contributing to these meadows, and the amount of fluid available per acre, are, however, very large. On the other hand, the district is but imperfectly provided with water-closets, and the sewage is probably very dilute. The flow is frequently shifted only once a day, though sometimes three or four times, and in the spring, before the first cutting, generally as much as twice. The water is, as a rule, applied the day the grass is removed. The application is continued through the greater part of the winter, excepting where the last crop has been fed off. Perhaps the sewage is unused about six weeks in the year. The land is heavy, and rather steeply inclined. To a few plots on one side of the valley sewage is applied by gravitation, through a pipe carried across from a higher level of the natural flow on the opposite side.

The herbage contains no clover ; and is, in fact, very soon brought down to a very simple character. It contains much couch, and also in some parts a good deal of crow-foot. When the plots are well managed four or five cuttings are obtained during the season. The crops of 1862 were sold at sums varying from 13*l.* to 38*l.* 5*s.* per imperial acre ; the difference depending much upon the amount of sewage, the character of the land, the aspect, and the previous management of the cuttings, all of which affect the amount and character of the produce ; but the convenience of position for cartage, proximity to other plots held by the purchaser, and other incidental circumstances, also sometimes affect the rates given for individual plots considerably.

Mr. McGill was of opinion that very much depended upon the quantity and strength of the sewage applied, and stated that portions of the land which had been irrigated by comparatively strong sewage were much more productive than others irrigated by almost pure water.

5. *Rose Bank, Broughton Road.*

These meadows, under the management of Mr. William Reid, comprise a little more than six imperial acres, and have been under irrigation

about 30 years. They are watered about 11 months in the year, at a cost of two shillings per week, given to a labourer who does the work of application at his meal-times and in over-time. The runs are, however, cleaned out in the winter by extra labour, paid for by the sale of the refuse, which commands about 2s. 6d. per ton. As the Table shows, the quantity of sewage available is enormous, being about 100,000 tons per acre per annum. Mr. Reid considered that quantity of water was a very important point, but that his supply was sufficient for considerably more land. He had not any more, however, at a convenient level, and did not consider that it would pay to be at any expense in raising the sewage. The irrigation is continued throughout the greater part of the winter as well as in the summer, and on Sundays as well as on other days. The water is applied to the same plot for three or four consecutive days, and the land gets, on the average, two such dressings for each cutting. It is considered better not to apply the sewage immediately after cutting, but to wait a few days until the grass has begun to shoot. The water does not run off the land clear. Four or five cuttings are obtained annually. The plots are from one-quarter to half an acre each, and they let at rates which amount to from 25*l.* to 30*l.* per imperial acre.

Mr. Reid's supply of sewage being so abundant, and having garden ground lying conveniently for its application, he occasionally applies it to about two acres; but he stated that he would go to very little expense in arrangements for the application of such small quantities as could be so used. The garden crops for which the sewage was found to be the most useful are turnips, cabbages, and onions.

6. *Quarry Holes.*

The Quarry-Holes meadows, farmed by Mr. Thomas Skirving, comprise only about eight imperial acres, and the amount of sewage available per acre is very large, being about 65,000 tons per annum. The whole is in permanent meadow. The sewage is applied day and night, and on Sundays, the whole year round, excepting during very hard frost. A plot of an acre, more or less, receives the supply for about two days at a time, and gets three such dressings between the cuttings; but the water is not put on until two or three days after cutting, nor is it considered desirable to apply it when the grass is far advanced. As the Table shows, the amount of population contributing to each acre is greater in the case of these meadows than in that of any other in the neighbourhood of Edinburgh, excepting those of Rose Bank, and the amount of water is also very large. There is no doubt that there is very extravagant expenditure of manurial constituents here, as there is indeed in all the other cases; but it must, at the same time, be admitted that it is under these conditions that a greater amount of produce is obtained per acre under the influence of sewage than anywhere else, and perhaps among all the Edinburgh sewage meadows those of the Quarry Holes stand second to none in point of evenness of herbage, and amount and value of produce per acre. The prevailing grasses are rough meadow-grass and common couch.

APPENDIX No. 3.

NOTES ON THE SEWAGE-IRRIGATED MEADOWS AT BEDDINGTON NEAR CROYDON.

The following particulars were obtained on the spot, partly from Mr. Latham the engineer to the Croydon Local Board of Health, and partly from Mr. Marriage, junior, son of the gentleman who rents and farms the irrigated meadows.

The population of Croydon contributing to the sewage tanks is about 16,000; and the water contributed to them is estimated at about 40 gallons per head per day without rainfall, and to average, the year round, perhaps 60 gallons per head per day with rainfall. These amounts are equal to about 65 tons per head per annum without, and 98 tons with rainfall. About 300 acres are rented by the Board at 4*l.* per acre without sewage, and sub-let to Mr. Marriage at 5*l.* per acre with sewage. Up to Midsummer 1864, 260 acres had been prepared for irrigation, of which about 250 might be considered as actually under irrigation during the year. It was intended to have 90 to 100 acres constantly under Italian rye-grass, but as yet not so large an area was under that crop.

The plan of irrigation is to let the sewage flow over from 20 to 30 acres for about four days and nights, and to give three such dressings between each cutting. As much of the water as can be recovered for the purpose is re-distributed, and in this way a large proportion is always used at least twice, sometimes three, and even four times over, and on the average about two and a half times, by which its utilisation and purification are rendered much more complete than otherwise would be the case. According to the figures given above there are about 6,250 tons of the dilute sewage with rainfall annually available for each of the 250 acres; but as so much of the water is re-used the average amount passing over each acre is very much more. There are also annually available for each acre the excretal matters of about 64 individuals of the mixed population of both sexes and all ages.

The land under sewaged Italian rye-grass is estimated to yield at least four cuttings, and from 30 to 35 tons of green produce per acre per annum. The cuttings commence in April, and last to the end of October, and even into November. This grass sells for about 25*s.* per ton in London, and is estimated to realise from 16 to 17 shillings per ton on the land. The sewaged meadow-grass also yields at least four cuttings annually, but it is much less liked than the Italian rye-grass by the London feeders, and is generally sold on the land by the rod, or grazed, and is estimated to yield several pounds less gross money return per acre per annum than the rye-grass.

Mr. Marriage does not apply the sewage in any systematic way to any other crops than permanent meadow and Italian rye-grass, but was about to try it on a small scale to mangolds.

There was a proposition under consideration by the Board to enlarge the area by 100 to 150 acres, which, notwithstanding the rapid increase of the population and of the sewage of the district, will, if carried out, considerably reduce the number of population contributing, and the amount of sewage available, to each acre annually.

About 180 tons of moist solid matter are annually deposited, or intercepted by the strainers, at the tanks, and are sold by the Board at a very low price per ton.

Before the above arrangements for passing the sewage of Croydon over the land were made, the Board were constantly subjected to legal

proceedings on account of the pollution of the river Wandle by the discharge of the sewage into it; but now those having the right of the fishing in the river have found it worth while to put down gratings to prevent the fish ascending the drainage outfall from the sewage-irrigated land.

In the following Table are collected together the results of some partial analyses of Croydon sewage, drainage, &c., which have been kindly communicated by Mr. Latham. The first four analyses are given on the authority of Messrs. Way, Evans, and R. D. Thomson, and the remainder on that of Messrs. Way and Evans alone.

TABLE I.—Results of the Analysis of the Water of the River Wandle, and of the Sewage of Croydon before and after Irrigation.

	Grains per Gallon.			
	Inorganic Matter.	Organic Matter.	Total Solid Matter.	Ammonia.
Samples collected October 1861.				
River Wandle above sewage outfall . . .	18'56	1'44	20'00	0'18
Do. do. below sewage outfall . . .	20'16	2'08	22'24	0'18
Sewage from an independent sewer . . .	48'30	52'20	100'50	6'70
Drainage after sewage irrigation . . .	23'40	2'40	25'80	0'21
Samples collected November 16, 1861.				
Sewage before reaching the tank, 11 a.m. . .	26'30	12'30	39'10	..
Drainage after sewage irrigation, 12.30 p.m. . .	21'25	6'50	27'75	..
Do. do. do. do. 2.30 p.m. . .	26'30	2'40	28'70	..
Do. do. do. do. 4.30 p.m. . .	25'50	3'45	28'95	..
Samples collected November 18, 1861 (after a sharp frost).				
Drainage after sewage irrigation, 2 p.m. . .	20'65	2'65	23'30	..
Do. do. do. do. 4 p.m. . .	20'95	2'90	23'85	..

The above analyses do not show what proportion of the several constituents existed in suspension and solution respectively; and, of the two samples of sewage, that collected November 16 contained, even before reaching the land, comparatively little solid matter of any kind. So far, however, as can be judged from these few, and little more than initiative results, it would appear that the water was to a considerably greater degree purified by its passage over and through the land than was the case in the Rugby experiments, in which the arrangements did not admit of the fluid being used two or three times over as with the more extended area at Croydon. The subject of the composition of the drainage-water passing from land liberally sewaged, but under arrangements for the re-distribution of the water until it is as far as practicable both utilised and purified, is, however, one of great importance, still requiring careful investigation.

LONDON:
Printed by GEORGE E. EYRE and WILLIAM SPOTTISWOOD,
Printers to the Queen's most Excellent Majesty.

REPORT
OF
EXPERIMENTS UNDERTAKEN BY ORDER OF
THE BOARD OF TRADE
TO DETERMINE THE
RELATIVE VALUES OF UNMALTED AND
MALTED BARLEY
AS FOOD FOR STOCK.

Presented to both Houses of Parliament by Command of Her Majesty.



LONDON:
PRINTED BY GEORGE EDWARD EYRE & WILLIAM SPOTTISWOODE,
PRINTERS TO THE QUEEN'S MOST EXCELLENT MAJESTY.
FOR HER MAJESTY'S STATIONERY OFFICE.

12179.

1866.

REPRINTED, 1868,
BY W. CLOWE AND SONS, DUKE STREET, STAMFORD STREET, AND CHANCING CROSS.



CONTENTS.

PART I.

INTRODUCTORY OBSERVATIONS, AND NOTICE OF PREVIOUS EXPERIMENTS.

	PAGE
Introductory Observations	5
I. The Experiments of Drs. T. and R. D. Thomson, in 1845-6	6
1. The Experiments with milking Cows	6
2. The Experiments with fattening Oxen	8
II. Experiments at Rothamsted, in 1848, 1849, and 1854	9
1. The Experiments in 1848 and 1849, with Sheep	9
2. The Experiments in 1854, with Pigs	13

PART II.

THE EXPERIMENTS MADE IN 1863-4 BY ORDER OF THE BOARD OF TRADE.

I. The Selection of the Barley, the Malting, and the Composition of the Unmalted and the Malted Grain	15
1. The Selection of the Grain, the Malting, &c.	15
2. The Analysis, and the Per-centage Composition of the Unmalted and the Malted Barley	17
3. The loss of Constituents by Malting	22
Summary	27
II. The Feeding Experiments in 1863-4	28
1. The Experiments with milking Cows	29
2. The Experiments with fattening Oxen	33
3. The Experiments with fattening Sheep	38
4. The Experiments with fattening Pigs	44
General Summary	51

APPENDIX.

The Determination of the Sugar in the Unmalted and the Malted Grain ..	56
The Determination of the Woody Fibre in the Unmalted and the Malted Grain	58
TABLES I.-IX., &c., relating to the Malting, and to the Composition of the Unmalted and the Malted Grain	59-67
TABLES X.-XXIV. relating to the Experiments with Cows	68-92
TABLES XXV.-XXXIII. relating to the Experiments with Oxen	93-101
TABLES XXXIV.-XLII. relating to the Experiments with Sheep	102-119
TABLES XLIII.-LI. relating to the Experiments with Pigs	120-137



REPORT OF EXPERIMENTS,

&c.

PART I.

INTRODUCTORY OBSERVATIONS, AND NOTICE OF PREVIOUS EXPERIMENTS.

It seems to have been established by experience that malt, given in moderate proportion, in admixture with other food, is much relished by most of the animals of the farm, and that it frequently exercises a beneficial influence upon their health and progress, more especially in the case of young or weakly animals. There are, however, so far as we are aware, no accounts on record showing either a greater amount of milk, or a greater amount of increase in live-weight, from the use of a given amount of malt than from that of the amount of barley from which it would be produced. On the contrary, all the trials with which we are acquainted, in which weights have been accepted as the measure of the result produced, have shown a better yield from a given amount of barley than from the amount of malt that would be produced from it. At the same time, it must be admitted that the evidence of some observers appears to show that, so far as quality is concerned, the meat of animals receiving a certain amount of malt in their food ranks deservedly high, and in a few cases in which the trials have been comparative it has been pronounced to be higher than when barley instead of malt was employed.

In the trials the results of which it is the object of this Report to record, 20 milking cows, 20 fattening oxen, 60 sheep, and 48 pigs have been experimented upon. Most of the experiments were arranged to compare the effects of a given amount of barley with those of the malt (and its dust) produced from an equal amount of barley taken from the same stock; and although

the quality, both of the milk and the meat produced, has not been overlooked, the *weight*, whether of milk or increase, has been taken as the chief measure of the effect produced.

Before detailing the results of these new experiments it will be well to give a brief summary of the results of any former experiments in which weights have been carefully taken and recorded.

I.—THE EXPERIMENTS OF DRs. T. AND R. D. THOMSON, IN 1845-6.

In 1845-6, the late Drs. Thomas and Robert Dundas Thomson, of Glasgow, made experiments on the relative qualities of barley and malt as food for stock. From June 24 to September 3, 1845, they experimented upon two milking cows, and from October 1, 1845, to January 16, 1846, upon two fattening oxen. The results of their inquiry are recorded in an official report.*

1. *The Experiments with Cows.*

In the experiments with the two cows both animals were kept upon the same description of food for a certain number of days in no case exceeding 16, and then, sometimes with, and sometimes without a short interval, they were put upon the food the results of which were to be compared with those of that previously given.

It is unfortunate that so few as two animals were taken though it would doubtless have been very difficult for the Drs. Thomson to have followed out their elaborate inquiry with larger number. It is also much to be regretted that the duration of each experiment was so short, that there was such a frequent change of food, and that (as was the case with the cows) the periods during which the foods to be compared with one another were given were successive instead of parallel.

The following is a summary of the Drs. Thomson's results with cows, calculated from the records given in the Report, arranged in a convenient form for showing the comparative effects of barley and malt in the different experiments. In experiments 1 and 2 the barley and malt were given entire, in experiments 3, 4, and 5 they were crushed, and in all the were steeped in hot water before being given to the animals.

* Reports to Her Majesty's Government in respect to Feeding of Cattle and Malt, April, 1846.

TABLE I.—CALCULATED SUMMARY of the Drs. THOMSON'S EXPERIMENTS with Cows.

	Food consumed.				Increase (or Loss) in Live-weight.	Milk yielded.		
	Total.	Per Head per Day.	Per 1000 lbs. Live-weight per Week.	To produce 100 lbs. Milk.		Total.	Per Head per Day.	Per 1000 lbs. Live-weight per Week.
Experiment 1.—Two Cows; Special Food—Barley; Period 11 Days.								
Barley	Lbs. 95	Lbs. 4·3	Lbs. 30·1	Lbs. 21·3	} - 41	Lbs.	Lbs.	Lbs.
Grass	1880	85·5	595·4	420·9		447	20·3	141
Experiment 2.—Two Cows; Special Food—Malt; Period 10 Days.								
Malt	Lbs. 112	Lbs. 5·6	Lbs. 38·6	Lbs. 28·4	} - 22	Lbs.	Lbs.	Lbs.
Grass	1700	85·0	585·2	431·5		394	19·7	136
Experiment 3.—Two Cows; Special Food—Barley; Period 16 Days.								
Barley	Lbs. 282	Lbs. 8·8	Lbs. 59·7	Lbs. 40·7	} 89	Lbs.	Lbs.	Lbs.
Malt	6	0·2	1·3	0·9				
Grass	532	16·6	112·6	76·8		693	21·7	147
Hay	763½	23·9	161·6	110·2				
Experiment 4.—Two Cows; Special Food—Malt; Period 16 Days.								
Barley	Lbs. 6	Lbs. 0·2	Lbs. 1·3	Lbs. 0·9	} 7·5	Lbs.	Lbs.	Lbs.
Malt	336	10·5	71·2	51·3		655	20·5	139
Hay	882½	27·6	186·8	134·7				
Experiment 5.—Two Cows; Special Food—Barley; Period 5 Days.								
Barley	Lbs. 116	Lbs. 11·6	Lbs. 78·9	Lbs. 57·2	} - 21·5	Lbs.	Lbs.	Lbs.
Hay	264	26·4	179·5	130·2		203	20·3	138

In the experiments of the Drs. Thomson screened malt, without the malt dust, was employed, and the exact proportion of the malt to the barley from which it was produced was not recorded. In the above Table (I.), therefore, the figures relating to malt represent the actual amounts of malt given, and not the amounts of barley from which the malt would be produced, as is the case in most of the Tables embodied in this Report.

The table shows that in each of the comparative trials a greater weight of screened malt than of barley was consumed

for the production of a given amount of milk ; and if the figures were amended so as to represent the amount of barley from which the malt had been produced, the results would of course appear still more unfavourable to the malted as compared with the unmalted barley.

2. *The Experiments with fattening Oxen.*

In all the experiments of Dr. Thomson with oxen the barley or the malt, as the case might be, was ground and mixed into mash with hot water before being given. Each comparative experiment was parallel as to time, instead of successive as in the case with the cows ; but only one animal was put on each description of food, the one receiving barley being designated as Bullock "A," and the one receiving malt as Bullock "B." There were five periods of experiment, one of 14, one of 15, one of 11, one of 17, and one of 18 days, in some cases with, and in some without, an interval between the periods, which, together, extended from October 1, 1845, to January 16, 1846, making a total period of 108 days, out of which the animals were under exact experiment only 75 days.

The following is a calculated summary of the results obtained during the five separate periods, amounting to 75 days in all.

TABLE II.—CALCULATED SUMMARY OF DR. THOMSON'S EXPERIMENTS WITH BULLOCKS

		Food consumed.				Increase in Live-weight.		
		Total.	Per Head per Day.	Per 1000 lbs. Live-weight per Week.	To produce 100 lbs. Increase.	Total.	Per Head per Week.	Per 1000 lbs. Live-weight.
		Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	
Bullock A.	Barley ..	766	10·2	64·5	411·8	186	17·4	
	Hay ..	876½	11·7	73·9	471·3			
	Cake ..	55½	0·7	4·7	29·8			
	Roots ..	1904	25·4	160·4	1023·7			
Bullock B.	Malt ..	768	10·2	54·0	376·5	204	19·0	
	Hay ..	1008	13·4	70·8	494·1			
	Cake ..	55	0·7	3·9	27·0			
	Roots ..	1904	25·4	133·8	933·8			

It will be obvious that no very satisfactory conclusions can be formed from results obtained with single animals only in each case, and over such short periods as those adopted in the experiments ; but, under the conditions stated, the results were as follow :

During the first 14 days the Bullock A, on barley and hay, gave more increase than Bullock B, on malt and hay, and required less food to yield a given amount of increase.

For two or three weeks from the conclusion of the above experiment the animals suffered from foot disease accompanied with fever, then very prevalent. On recovery the comparative trial was resumed for 15 days, Bullock A having barley, hay, and roots, and Bullock B malt, hay, and roots. The result was again in favour of the unmalted barley, both in regard to the actual amount of increase in live-weight and to the amount of food required to yield a given amount of increase.

The next period was consecutive with the last, and extended over 11 days. During this period the turnips were found to be much diseased, and both animals suffered in consequence, especially the one having barley, which gave much less increase than the other.

The next period was one of 17 days, was consecutive with the preceding one, and the weights were considerably in favour of the bullock having barley.

The fifth and last period commenced 10 days after the conclusion of the preceding one, lasted 18 days, and the result was in favour of the malt.

The Table (II.) gives the results of these five periods collectively. It is seen that Bullock B, receiving malt, gave, under the conditions stated, rather more increase than the Bullock A, receiving barley, and also consumed less malt to yield a given amount of increase than the other did of barley. But if the comparison were made, not between the actual weights of barley and malt, respectively, that were consumed, but between the amount of barley actually consumed in the one experiment, and the amount of barley from which the malt consumed in the other would be produced, the result would then be found to be in favour of the unmalted instead of the malted barley.

With fattening bullocks, then, as with milking cows, the results obtained by the Drs. Thomson were unfavourable to the use of malted as compared with unmalted barley, and the conclusions they drew on the subject were consistent with the result here stated.

II.—EXPERIMENTS AT ROTHAMSTED, IN 1848, 1849, AND 1854.

1. *The Experiments in 1848 and 1849, with Sheep.**

A summary of the results obtained under this head is given in the following Table (III.).

* 'Agricultural Chemistry: Sheep Feeding and Manure.' ('Journal of the Royal Agricultural Society of England,' Vol. X., Part I.)

TABLE III.—SUMMARY OF RESULTS obtained at ROTHAMSTED, in 1848 and with SHEEP.

	Food consumed.				Increase in Live-weight.		
	Total.	Per Head per Day.	Per 100 lbs. Live-weight per Week.	To produce 100 lbs. Increase.	Total.	Per Head per Week.	Per Live-weight.
Experiment 1.—From June 5 to October 17, 1848 = 19 Weeks and 1 Day; under							
5 Sheep { Barley (ground)	Lbs. 665	Lbs. 1·0	Lbs. 5·2	Lbs. 479	} 139	Lbs. 1·47	
5 Sheep { Clover Hay ..	1986	3·0	15·6	1429			
5 Sheep { Malt (ground) ..	625	0·94	5·0	517	} 121	Lbs. 1·28	
5 Sheep { Clover Hay ..	1974	2·9	15·7	1631			
Experiment 2.—From March 20 to May 29, 1849 = 10 Weeks; under Cover							
4 Sheep { Barley (ground)	Lbs. 280	Lbs. 1·0	Lbs. 5·0	Lbs. 346	} 81	Lbs. 2·03	
4 Sheep { Mangolds ..	3867	13·8	69·1	4774			
5 Sheep { Malt (and Dust) ground	283	0·81	4·0	270	} 105	Lbs. 2·09	
5 Sheep { Mangolds ..	4694	15·2	66·5	4470			
Experiment 3.—From March 20 to May 29, 1849 = 10 Weeks; under Cover							
4 Sheep { Barley (ground) and steeped	Lbs. 280	Lbs. 1·0	Lbs. 4·8	Lbs. 276	} 101½	Lbs. 2·53	
4 Sheep { Mangolds	5322	16·8	90·5	5243			
4 Sheep { Malt (and Dust) ground and steeped	227	0·81	3·8	291	} 78	Lbs. 1·97	
4 Sheep { Mangolds	4458	15·9	74·5	5715			
5 Sheep { Malt (and Dust) ground	350	1·0	4·9	324	} 108	Lbs. 2·16	
5 Sheep { Mangolds	5404	15·4	75·0	5004			
Experiment 4.—From March 12 to May 14, 1849 = 9 Weeks; folded in the							
27 Sheep { Barley	Lbs. 851	Lbs. 0·5	Lbs. 2·5	Lbs. 182	} 468	Lbs. 1·92	
27 Sheep { Clover Chaff ..	851	0·5	2·5	182			
27 Sheep { Swedes (ad lib.)			
27 Sheep { Malt (and Dust) ..	680	0·4	2·0	156	} 437½	Lbs. 1·80	
27 Sheep { Clover Chaff ..	851	0·5	2·5	194			
27 Sheep { Swedes (ad lib.)			

In experiment 1 the design was to give equal weights of barley and of malt, 1 lb. per sheep per day, but not quite so much of the malt was consumed. The complementary food was in both cases clover-chaff, which, with water, was given *ad libitum*, but weighed, and the experiment was continued for 19 weeks. The Table shows, not only that the sheep having unmalted barley gave a greater actual increase in live-weight, but also that it required a greater weight of malt than of barley to yield a given amount of increase. The result was in reality still more unfavourable to the malt than the figures show, for the 517 lbs. of screened malt consumed to produce 100 lbs. of increase would have required about 645 lbs. of barley to produce it, whereas only 479 lbs. of raw barley were consumed to produce 100 lbs. of animal increase.

In the second experiment, as in the first, the sheep were fed under cover. The plan was to give to one lot 1 lb. of barley per head per day, to another the amount of malt and malt-dust produced from 1 lb. of barley, and to both mangolds *ad libitum*, but weighed. The barley and the malt were ground; and the experiment was continued for ten weeks, with four sheep on the barley, and five on the malt diet. As the figures stand, the result appears considerably in favour of the malt; and even when the malt and malt-dust consumed are calculated up to the amount of barley from which they would be produced, the result is still slightly in favour of the malt as regards the amount of food required to produce a given amount of increase in live-weight.

The plan of experiment 3 differed little from that of experiment 2. In the first pen the amount of barley, and in the second the amount of malt, with its dust, were the same as before, but each was now steeped for some time as well as ground before being given. In the third pen each sheep received 1 lb. of malt (with its dust) per head per day, that is, the same weight as those in pen 1 received of barley, instead of, as in pen 2, only as much as would be produced from 1 lb. of barley, and instead of being both ground and steeped, the malt was now only ground. Comparing the result of pen 1 with barley ground and steeped, with that of pen 2 with malt both ground and steeped, it is decidedly in favour of the unmalted barley, and of course more so than the figures show if the amount of malt consumed be reckoned up to the amount of barley from which it would be produced. The result of pen 3 with the increased amount of malt, but only ground, and not afterwards steeped, is still more unfavourable to its use.

As a control to the above experiments, in all of which the

sheep were fed under cover, in some with clover-hay without any succulent food, and in others with mangolds (themselves highly saccharine) in addition to the barley or the malt, another experiment, No. 4, was made, in which there was a larger number of sheep (twenty-seven) in each lot, the animals were folded in the open field, they received only about half as much barley or malt per head per day, $\frac{1}{4}$ lb. clover-chaff per head per day, and swedes *ad libitum* and unweighed. To lot 1 the daily allowance of barley was $\frac{1}{2}$ lb. per head, and to lot 2 as much malt with its dust as was produced from $\frac{1}{2}$ lb. of barley, and the experiment was continued for nine weeks.

The rate of increase upon a given live-weight within a given time was somewhat under the average obtained when sheep are liberally fed on fattening food, and was not equal to that in experiments 2 and 3, in which the animals had mangolds, and were fed under cover. The total increase of the lot having barley was 468 lbs. against 437 $\frac{1}{2}$ lbs. on the lot having a corresponding amount of malt; and reckoning the malt (and dust) consumed to produce 100 lbs. of increase up to the amount of barley which it represented, the result was decidedly in favour of the unmalted as compared with malted barley.

It is seen, then, that in only one of these earlier comparative trials with sheep was the result in favour of malt as compared with the amount of barley from which it would be produced. The results of the Drs. Thomson, with milking cows and fattening oxen, also afforded preponderating evidence in favour of the use of unmalted rather than malted barley, as a staple food for stock.

So far as we are aware there are no other comparative trials of barley and malt recorded, in which the weights of the food consumed, and of the milk or increase in live-weight yield, have been taken as the measure of the effects produced. It is true that there are records enough of trials in which malt has been shown to be a good food for stock, and no further evidence on that point can be needed. But whether or not the favourable result, so far as the health and progress of the animals are concerned, has been economical compared with what it would have been had raw barley been used instead, is quite another matter, and it is the question of the comparative economy of unmalted and malted barley as food that the experiments above quoted, and those to which this report has special reference, have been undertaken and arranged to throw light upon.

2. The Experiments in 1854 with Pigs.*

Independently of certain theoretical considerations, one object of these experiments was to determine whether any or what benefit would be likely to accrue, either to the growers of sugar, or to the agricultural interest of this country, if the lower qualities of cane-sugar were admitted duty-free for the feeding of animals; another was to provide data bearing upon the question whether or not any advantage would result from the conversion of a portion of the starch of the food of animals into sugar by subjecting it to the malting process.

The plan adopted was, to give to each of several lots of pigs a fixed amount of lentil-meal and bran, enough to supply all the nitrogenous matter that they would require, but leaving them deficient in the necessary amount of the non-nitrogenous constituents of food, which were supplemented in one experiment by as much starch, in another by as much sugar, and in another by as much of both starch and sugar, as the animals chose to eat; whilst in a fourth experiment they were allowed to take lentil-meal, bran, starch, and sugar, each separately and *ad libitum*.

The results given in the following Table (IV.) of the first two of the experiments above referred to, in which both lots of animals had the same amount of lentil-meal and bran, but the one starch and the other sugar, in addition and *ad libitum*, bear the most directly upon the questions here under consideration.

TABLE IV.—EXPERIMENTS with PIGS, on the Equivalency of STARCH and SUGAR in FOOD.

Period of Experiment 10 weeks; February 27 to May 8, 1854.

Food consumed.					Increase in Live-weight.		
	Total.	Per Head per Day	Per 100 lbs. Live-weight per Week.	To produce 100 lbs. Increase.	Total.	Per Head per Week.	Per 100 lbs. Live-weight per Week.
	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.
Pigs { Lentil-meal	672	3·20	16·4	272	} 247	8·23	6·03
{ Bran	126	0·60	3·1	51			
{ Sugar, <i>ad libitum</i> (1)	388½	1·85	9·5	157			
) Containing dry substance	(363½)	(1·73)	(8·9)	(147)
Pigs { Lentil-meal	672	3·20	16·4	271	} 248	8·27	6·06
{ Bran	126	0·60	3·1	51			
{ Starch, <i>ad libitum</i> (2)	450½	2·15	11·0	182			
) Containing dry substance	(362½)	(1·73)	(8·9)	(146)			

* * On the Equivalency of Starch and Sugar in Food.—Report of the British Association for the Advancement of Science for 1854.

It is seen that the two lots consumed almost identical amounts of lentil-meal and bran, whether reckoned per head per 100 lbs. live-weight per week, or to produce 100 increase; but that the one lot consumed in addition considerably less of the raw sugar than the other did of the raw starch; however, the amounts of water which the sugar and the starch contained in the condition in which they were respectively fed to the animals be deducted from the actual weights consumed, is found, as shown by the figures given in parenthesis in Table, that the amounts of real dry or solid substance consumed were all but absolutely identical in the two cases. Indeed it is borne in mind that the animals were dependent on the sugar or the starch supplied to them for about one-third of the solid matter of their food, and that in each case they were able to take as much as they chose, the coincidence in the amounts of dry or solid substance consumed in the two cases is striking.

The amounts of increase, whether reckoned per head per 100 lbs. live-weight per week, were also all but identical.

The results of these experiments indicate an almost equivalency of the dry or solid substance of cane-sugar and starch in the food of the pig, provided, of course, that neither is given in undue proportion, and that other constituents be at the same time supplied in sufficient amount; indeed, the two probably only differ in point of fact in relative food capacity, a very slight proportion in which they are known to differ in percentage amount of carbon.

If, then, cane-sugar have no higher, and perhaps even a lower, value as a constituent of food than starch, a consideration of the relative prices of sugar duty-free, and of the starch generally used for feeding, and also of the fact that the latter at the same time supply a considerable amount of the needed nitrogenous constituents of food, will afford an easy means of estimating the relative economy of the two.

As cane-sugar contains somewhat less carbon than an equal weight of starch, so does the saccharine matter of malt in its turn somewhat less than an equal weight of the sugar of the cane; hence, so far as composition is concerned, it would be concluded, *à priori*, that a given weight of cane-sugar would have a higher value as a constituent of food than an equal weight of the sugar of malt; on the other hand, the sugar of malt is nearly allied to that which is the product of the transformation of starch within the animal body than is that of the cane.

The most obvious conclusion from these experiments is that there would be no economy in substituting a

of the starchy grains usually used as food by a corresponding amount of the sugar of the cane; and a very probable conclusion was, that there would be as little or less advantage from the conversion of a certain amount of the starch of food into the sugar of malt by subjecting the starchy grains to the process of malting.

We proceed now to consider the experiments which form the special subject of the present Report.

PART II.

THE EXPERIMENTS MADE IN 1863-4 BY ORDER OF THE BOARD OF TRADE.

1. THE SELECTION OF THE BARLEY, THE MALTING, AND THE COM- POSITION OF THE UNMALTED AND THE MALTED GRAIN.

1. *The Selection of the Grain, the Malting, &c.*

In order to determine fairly the comparative feeding quality, and the comparative economy as food, of unmalted and malted barley, it was decided to employ malt produced from barley taken from the same stock as that given for comparison in the unmalted state; and to give to parallel lots of animals, in one case a given amount of barley, and in another the amount of malt (with its dust) which was found in actual working to be produced from that amount of raw barley. It was also thought desirable to employ in some of the experiments barley of a quality admitted to be well adapted for the malting process, and in others such as would not be considered suitable for making brewer's malt, but which would be considered well adapted for feeding, and to determine the amount and composition, not only of the final products (malt and malt-dust), but also of some of the intermediate products of the malting of each of the descriptions of grain selected.

In order to accomplish these ends it was necessary to have access to, and liberty to disturb, and take samples from, the "floors," during growth. The Board of Excise having been communicated with, arrangements were made for conducting the operations in one of the maltings of the Messrs. Gripper, at Hertford, and ample facilities were afforded for doing all that might be deemed desirable, subject only to the observation of

an officer appointed for the purpose by the Surveyor of district.

These matters being arranged, 70 quarters of barley, of but not first-class, malting quality, were purchased. Half reserved to be consumed in the raw state, and half was m and screened in the usual way. For distinction, this lot wi designated No. 1.

70 quarters of barley of fair feeding quality were also chased, half reserved to be consumed in the raw state, and other half malted, &c., as No. 1. This lot will be design No. 2.

The grain was measured over, every eighth bushel weigh and samples were taken in the raw state, at different stag growth, and when completely malted, in order to have the m of determining, by analysis and calculation, the amoun chemical change, and of loss, during the malting process.

The following Table (IV.) shows the quantity, both by mea and weight, of each lot of barley malted, and also the qua and proportion of malt and malt-dust produced in each case, correction having been made for the quantities remove samples during the process.

TABLE IV.—SHOWING the QUANTITIES of BARLEY MALTED, and of MALT-DUST produced.

		Measure.	Weight per Bushel.	Total Weight.	Malt to 100 Barley; by Measure.	Malt and Dust to 100 Barley; by Weight.
		Bushels.	Lbs. oz.	Lbs.		
Barley No. 1.	{ Unmalted ..	280	54 10	15,295
	{ Malted ..	290	41 10	12,072½	103·6	78·9 }
	{ Malt-dust	336	..	2·2 }
Barley No. 2.	{ Unmalted ..	280	50 10	14,175
	{ Malted ..	284	37 4	10,578	101·4	74·6 }
	{ Malt-dust	454½	..	3·2 }

It is seen that the malt from the good malting barley, No. 1, measured about $3\frac{1}{2}$ per cent., but that from the feeding barley No. 2, scarcely $1\frac{1}{2}$ per cent. more than the barley from which was produced. In weight, however, both lost considerably. No. 1 about 21, and No. 2 about $25\frac{1}{2}$ per cent., reckoning the screened malt as a product of the process; but, even including the malt-dust as a product, the loss in weight was, with barley No. 1 about 19, and with barley No. 2 a little over 22 cent.

The question arises—of what does this loss in weight by lting consist?

Before fully considering this point it will be well to call attention to the difference in the per-centage composition of the ley, and its products at different stages of growth, as shown the results of the analyses, without the aid of which that question could not be answered.

The Analysis and the Per-centage Composition of the unmalted and the malted Barley.

For the purpose of these determinations a sample of 25 lbs. s taken from the grain in the dry state before steeping, after eeping when thrown out from the couch, at intervals of several ys during growth, and when completely malted and screened. As soon as the samples were taken they were sent to the thamsted Laboratory, and there at once stove-dried for pre- vation. The analyses were also all made at Rothamsted; but should be observed that the determinations of the sugar were de by Mr. Hugh Morris, of the Inland Revenue Laboratory, merset House, who was kindly sent down by Mr. George illips, the able Director of that laboratory, to assist in the ytical work.

Some particulars relating to the analyses will be found at . 56-58, and the detailed results are given in Tables I.-V., . 59-63, in the Appendix.

Table V. (p. 18) summarises, for barley No. 1, the composi- n of the grain at the different stages; the upper division owing the composition of the specimens in the condition of isture in which they were sampled; the middle division the nposition of the dry or solid substance, that is, exclusive of all isture in each case; and the lower division, that of the barley its natural state of dryness, of the malt and malt-dust as kiln- ed, and of the intermediate products of growth assuming each have been brought to the same condition of dryness as the n-dried malt.

Table VI. (page 19). shows the same particulars for barley . 2 as does Table V. for barley No. 1.

For comparison with the results in these two tables, Table II., p. 66 in the Appendix, gives similar particulars, so far as y were obtained, relating to the barley, and the preparation of malt, used in some of the experiments with sheep at Roth- sted in 1849.

Owing to the very great difference in the amount of moisture the different specimens in the condition in which they were mpled for analysis (the barley being in the usual marketable

TABLE V.—SHOWING the COMPOSITION of BARLEY No. 1, before Malting, at different Stages of the Process, and when completely Malting.

	Barley, before steeping.	Growing.						Screened Malt.	M d
		Nov. 2. As thrown from the Couch.	Nov. 6. 4½ Days on the Floor.	Nov. 10. 8 Days on the Floor.	Nov. 12. 10½ Days on Floor (1st third to Kiln).	Nov. 14. 12½ Days on Floor (2d third to Kiln).	Nov. 16. 14½ Days on Floor (3d third to Kiln.)		

1.—In the condition of Moisture as sampled.

Sugar	2.11	0.89	4.74	5.95	No sample taken.	6.92	7.23	10.28	1
Starch (and dextrine)	66.24	46.48	43.44	42.10		41.94	41.75	67.23	1
Woody-fibre	3.86	3.00	2.88	3.02		2.95	2.99	4.51	
Albuminous (or "flesh-form- ing") matters*	8.09	5.63	5.75	5.92		6.09	6.19	9.29	2
Mineral matter (ash)	2.06	1.30	1.32	1.36		1.39	1.40	2.63	
Total solid matter	82.36	57.30	58.13	58.35		59.29	59.56	93.34	8
Moisture	17.64	42.70	41.87	41.65		40.71	40.44	6.66	1
Total	100.00	100.00	100.00	100.00		100.00	100.00	100.00	10
* Containing nitrogen	1.28	0.89	0.91	0.94		0.97	0.98	1.47	

2.—Exclusive of Moisture.

Sugar	2.56	1.56	8.16	10.19	No sample taken.	11.67	12.14	11.01	1
Starch (and dextrine)	80.42	81.12	74.72	72.16		70.73	70.09	72.03	4
Woody-fibre	4.69	5.22	4.96	5.18		4.98	5.03	4.84	
Albuminous (or "flesh-form- ing") matters*	9.83	9.83	9.89	10.14		10.27	10.39	9.99	1
Mineral matter (ash)	2.50	2.27	2.27	2.33		2.35	2.35	2.17	
Total solid matter	100.00	100.00	100.00	100.00		100.00	100.00	100.00	11
* Containing nitrogen	1.56	1.56	1.57	1.61		1.63	1.65	1.58	

3.—Barley, Malt, and Malt-dust, as sampled; intermediate Products in the same condition Dryness as Malt.

Sugar	2.11	1.46	7.61	9.51	No sample taken.	10.90	11.33	10.28	
Starch (and dextrine)	66.24	75.72	69.75	67.35		66.01	65.42	67.23	
Woody-fibre	3.86	4.88	4.63	4.84		4.65	4.70	4.51	
Albuminous (or "flesh-form- ing") matters*	8.09	9.17	9.23	9.47		9.59	9.70	9.29	
Mineral matter (ash)	2.06	2.11	2.12	2.17		2.19	2.19	2.63	
Total solid matter	82.36	93.34	93.34	93.34		93.34	93.34	93.34	
Moisture	17.64	6.66	6.66	6.66		6.66	6.66	6.66	
Total	100.00	100.00	100.00	100.00		100.00	100.00	100.00	
* Containing nitrogen	1.28	1.46	1.47	1.50		1.52	1.54	1.48	

condition,* the intermediate products in the wet state in after steeping they are spread on the floor to grow, and then (and malt-dust kiln-dried) the results as given in the upper portion of Tables V. and VI. do not at once bring to view variation in the composition of the solid substance of the so clearly as those in the middle and lower divisions.

In the middle division, which shows the composition of grain exclusive of all moisture, the variation in the composition of the solid substance of the grain as the sprouting proceeds is the most strikingly brought out; but, inasmuch as in practice grain malted for feeding purposes would, at whatever stage of growth were stopped, (if dried at all) probably be kiln-dried about the same extent as brewer's malt, it will be more to illustrate the variation in composition by reference to figures given in the bottom division of the tables.

The most striking change which the figures show is the increase in the amount of sugar as the malting proceeds. It may be said that the most characteristic effect of the malting process is the conversion of a portion of the starch of the grain into dextrine, and then a portion of the dextrine into sugar, changes which are effected by the agency of a substance called diastase, which is itself a product of the transformation of a portion of the nitrogenous substance of the grain. It is to be observed, however, that the first stage of the malting process, that of steeping the grain, dissolves a considerable proportion of the small amount of sugar already existing in it; and by analysis of the steep water as well as the grain, it is found that a certain amount both of nitrogenous and mineral matters is also extracted at this stage. It is seen from the experiments to which the tables refer, the amount of sugar was increased in one case from about $2\frac{1}{2}$, and in the other about $3\frac{1}{2}$ per cent. in the dry substance of the raw grain, from 11 to 12 per cent. in that of the finished malt.

It is the substance diastase, developed during germination, above referred to, which, in the process of "mashing," induces the conversion, first into dextrine and then into sugar, of the starch which has remained unchanged by the malting. It is the same substance which gives to an infusion of malt the property of inducing a like change in the starch of ungerminated grain, or other starchy substances mixed with it. This conversion takes place the most rapidly at a temperature of 170° F. It proceeds pretty actively, however, at ordinary

* The amount of moisture in grain varies according to season and other circumstances. Barley in the bulk may be reckoned to contain generally between 12 and 18 per cent. of moisture, but samples kept in dry places, or much malted, may contain several per cent. less.

peratures, provided the starch have previously been gelatinized by means of hot water; and Mr. George Phillips informs us that it had been found in experiments made at the Inland Revenue Laboratory, that diastase even acts upon starch when ground malt is mixed with water at ordinary temperatures.

As the sugar produced in the process of malting is formed at the expense of the starch, which is first converted into dextrine, the amount of starch and dextrine is necessarily reduced in about the same degree that that of the sugar is increased. A portion of the sugar, however, undergoes further change during the growth, the result of which is an actual loss of some of the original non-nitrogenous substance of the grain, as will be further illustrated presently.

The proportion of woody-fibre seems to increase slightly as the growth proceeds; though it is not much higher than in the barley in any of the products of the malting process excepting the malt-dust, the dry substance of which contains about twice as much as that of either the raw barley or any of the intermediate products.

The proportion of nitrogenous or so called "flesh-forming" substances is seen to be somewhat higher in the dry or solid substance of the intermediate products of growth than in the raw barley, and in that of most of them higher than in that of the finished and screened malt. That the malt contains a somewhat lower per-centage of nitrogenous substances than the intermediate products is, of course, explained by the fact that by the screening process it has been deprived of the malt-dust, which has withdrawn nitrogen from the grain during growth, and itself contains a very high per-centage of it.

Within certain limits, the higher the per-centage of nitrogenous matters, the greater will be the value of an article of food, either as such, or on account of the richness of the manure obtained from the animals consuming it. But inasmuch as the increased proportion of nitrogenous substances in the grown grain is not due to any gain of them during the process, but, as will presently be seen, to a loss of other substances, by which it is only their proportion, and not their actual amount, that is increased, it would obviously be a delusion to conclude that there is any advantage or economical gain in the increased per-centage of those important constituents which is obtained by submitting the grain to germination; indeed, as will be seen further on, there will always be a greater or less loss of the nitrogenous substances of grain in converting it into malt, so that, so far as the feeding value is dependent on the amount of nitrogenous constituents, a given amount of grain will provide more of them in the raw state than after it has been malted. It is true that a given weight

of malt will contain a larger amount of nitrogenous substance than an equal weight of raw barley, but that given weight of malt will have required $1\frac{1}{2}$ th to $1\frac{1}{4}$ th of its weight of raw grain besides the cost of the process, to produce it.

From the high percentage of nitrogenous substances in malt-dust, it is obvious that there would be a loss, both of feed and manurial matter, if it were not employed along with the grain whenever grain were malted for feeding purposes.

It will be observed that the feeding barley (No. 2) contains more nitrogenous matter, and more woody-fibre, but less starch than the better malting sample No. 1; and this difference in composition was pretty uniformly maintained between the corresponding products of growth of the two descriptions of grain. Indeed, the samples selected as the best for malting for brewing purposes are the plumper, the more starchy, and the less nitrogenous grains; whilst those employed for feeding are generally thinner, more nitrogenous, and less starchy.

3. *The Loss of Constituents by Malting.*

In the following tables (in Table VII. for barley No. 1, and Table VIII. for barley No. 2) the foregoing analytical results are applied so as to show the actual and percentage loss of constituents which the barley had suffered at certain stages, and at conclusion of the malting process. At the periods selected these estimates, not only were samples taken for analysis, but the whole quantity of grain on the floors was measured over, and the weight of every eighth bushel taken, in order to determine the total weight as already referred to. For the estimates founded on the data so obtained absolute accuracy cannot be claimed since every individual bushel was not weighed, but it is believed that the results sufficiently nearly represent the truth for all practical purposes.

It appears that, in the steeping, No. 1 barley absorbed about 43, and No. 2 barley rather more than 47 per cent. of the weight of water. Each lost at the same time a certain amount of solid matter, which consisted mainly of saccharine, nitrogenous, and mineral* substances, more of which were abstracted from the less perfectly matured second quality grain than from the other.

The greatest amount of loss of solid substance during growth on the floor was of non-nitrogenous organic matter

* The figures do not indeed show a loss of mineral matter by steeping in the case of the second quality barley; but, inasmuch as the so-called "mineral matter" was the crude ash determined by burning, and the samples were not free from more or less of incombustible impurity, this would account for some discrepancy in the figures in relation to mineral matter.

TABLE VII.—SHOWING THE LOSS OF CONSTITUENTS OF BARLEY “No. 1” at certain Stages, and at the Conclusion of the Malting Process.

	Barley before steeping.	Nov. 2. As thrown from the Couch.	Nov. 6. 44 Days on the Floor.	Nov. 10. 8 Days on the Floor.	Final Products. Sent to the Kiln Nov. 12, 14, and 16 = 10, 12, and 14 Days on the Floor.			Loss.
					Screened Malt.	Malt dust.	Total.	
Actual Weights, lbs.								
As sampled	15,295	21,898	21,459	20,705	12,072½	336	12,408½	2,886½
Total dry or solid matter	12,597	12,548	12,474	12,081	11,268½	296½	11,565	1,032
Non-nitrogenous organic matter	11,044	11,030	10,957	10,574	9,902½	191½	10,094	950
Nitrogenous matter	1238.0	1233.2	1233.8	1225.4	1121.7	79.8	1201.5	36.5
Mineral matter	314.8	284.3	283.5	281.5	244.5	25.0	269.5	45.3
Proportion to 100 before steeping.								
As sampled	100	143.2	140.3	135.4	78.93	2.20	81.13	18.87
Total dry or solid matter	100	99.6	99.0	95.9	89.45	2.35	91.80	8.20
Non-nitrogenous organic matter	100	99.9	99.2	95.7	89.66	1.74	91.40	8.60
Nitrogenous matter	100	99.6	99.7	99.0	90.61	6.45	97.06	2.94
Mineral matter	100	90.3	90.1	89.4	77.68	7.94	85.62	14.38

TABLE VIII.—SHOWING THE LOSS OF CONSTITUENTS OF BARLEY "No. 2" at certain Stages, and at the Conclusion of the Malting Process.

	Barley before steeping.	Nov. 6. As thrown from the Couch.	Nov. 14. 8 Days on the Floor.	Final Products. Sent to the Kiln Nov. 17, 19, and 21 = 11, 13, and 15 Days on the Floor.			Loss.
				Screened Malt.	Malt-dust.	Total.	
Actual Weights, lbs.							
As sampled	14,175	20,870	19,654	10,578	454½	11,032½	3,142½
Total dry or solid matter	11,488	11,412	11,044	10,084	392	10,476	1,012
Non-nitrogenous organic matter	9,789	9,736	9,378	8,633	256½	8,889½	899½
Nitrogenous matter	1404.0	1380.3	1363.6	1200.7	100.3	1301.0	103.0
Mineral matter	295.3	295.3	302.5	250.3	35.5	285.8	9.5
Proportion to 100 before steeping.							
As sampled	100	147.2	138.7	74.62	3.21	77.83	22.17
Total dry or solid matter	100	99.3	96.1	87.78	3.41	91.19	8.81
Non-nitrogenous organic matter	100	99.5	95.8	88.19	2.62	90.81	9.19
Nitrogenous matter	100	98.3	97.1	85.52	7.14	92.66	7.34
Mineral matter	100	100.0	102.5	84.76	12.02	96.78	3.22

doubtless chiefly sugar, destroyed by decomposition, as the growth advanced.

Of nitrogenous substances, a certain but apparently not very material amount is extracted by the steep water, the partial analysis of which is given in Table VII. p. 65, in the Appendix.

During the actual processes of germination and growth, there is, so far as existing knowledge goes, no necessary loss of nitrogen unless the moist grain be allowed to get foul. Nor, perhaps, is there in the drying any really unavoidable loss; though in practice, as at present conducted, a certain amount of the young shoots or "malt-dust" (which is highly nitrogenous), always passes through the wires during the drying, and this portion would be lost as food if not as manure also.

According to the figures in the lower divisions of Tables VII. and VIII., the total loss of nitrogen or nitrogenous substances amounted in the case of barley No. 1 to about 3, and in that of barley No. 2 to about 7 per cent. of the original quantity in the grain; and whilst the screened malt from the barley No. 1 retained 90½, that from barley No. 2 only retained 85½ per cent. of the original nitrogen. Barley No. 2 was, indeed, as already stated, the thinner and the more nitrogenous. It was, moreover, one day longer on the floor, and was pronounced to be relatively to the other rather further grown, in fact rather too far, and to be somewhat exhausted and not very fresh. Hence, doubtless, the larger proportion of the nitrogen of the grain extracted by the sprouts, the larger amount in the malt-dust, and the larger amount of the latter (and so of nitrogenous substance) detached and lost in the manipulations in the kiln-drying.

It may be remarked that in the experiments made at Rothamsted on this subject in 1849 (the results of which are given in Table IX. p. 67 in the Appendix), the proportion of the nitrogen of the original grain retained by the screened malt was 86½ per cent., that recovered in the malt and kiln-dust was nearly 10 per cent., and that totally lost was about 3½ per cent. In this case, however, the grain was rather longer (about 16 days) on the floor.

Of mineral matter the necessary loss will be confined to that extracted by the steep water, and to the amount contained in that portion of the "dust" which passes through the wires during the drying in the kiln.

It may, perhaps, be safely concluded, that if malt were manufactured for feeding, the growth would not be carried so far as it is for brewing purposes; that, if screened, the malt would retain over 90, and perhaps sometimes nearer 95 per cent. of the original nitrogen of the grain; and that, including the malt-dust as a product of the process, about 98 per cent. of the total nitrogen

of the grain might be recovered, leaving only about 2 per cent as the probable average loss by extraction by the steep-water and by the loss of a portion of the malt-dust.

The following Table (IX.) gives a condensed view of the products and loss from 100 lbs. of barley in each of the two experiments.

TABLE IX.—Showing the PRODUCTS and LOSS on MALTING 100 lbs. BARLEY.

		Barley No. 1.	Barley No. 2.
Malt		78·93	74·62
Malt-dust		2·20	3·21
Total Products		81·13	77·83
Loss ..	{ Moisture	12·12	15·03
	{ Non-nitrogenous vegetable substance ..	6·21	6·34
	{ Nitrogenous substance	0·24	0·73
	{ Mineral matter	0·30	0·07
		100·00	100·00

Thus, the fair malting sample (No. 1) gave about 79 per cent of its weight, and the feeding sample (No. 2) only about 74½ per cent. of its weight of screened malt. Adding the malt-dust, No. 1 gave rather more than 81, and No. 2 rather less than 80 per cent. of total products. The result is, then, that even reckoning in the malt-dust as a product of the process, there was a loss on the original weight of the barley of about 19 per cent in the one case, and of about 22 per cent. in the other. In each case about two-thirds of the loss in weight (in the one rather under and in the other rather over) was only moisture driven off in the kiln-drying; but there was, besides, in each case, a loss of about 7 per cent. of real solid substance or food material, of which about 6½ was non-nitrogenous vegetable substance, from ¼ to ¾ nitrogenous substance, and the remainder mineral matter.

From the above facts it will be readily understood how it is that the products of the malting process have a higher percentage of nitrogenous or "flesh-forming" substances than the barley from which they were produced. There is, indeed, an actual loss of these substances; but inasmuch as there is a much greater loss of the non-nitrogenous matters, and a considerable dissipation of water in the kiln-drying, which together reduce the weight of the products to about 20 per cent. (less or more) below that of the barley from which they were produced, this diminishes

* See foot note on p. 22.

weight, though containing even a less actual amount of nitrogenous substances, nevertheless contains a higher proportion of them in 100 parts. It is obvious, therefore, as before observed, that it would be a delusion to represent the higher per-centage of nitrogenous substances in the malted grain as indicating any gain by the process, so far as the amount of those important constituents of food is concerned. There is, in fact, a loss, not a gain; but as there is a much greater loss of other matters the proportion of them in what remains is somewhat the greater.

The loss and chemical changes which barley undergoes by malting may be summarily enumerated as follow :

1. The weight of the malt, together with the malt-dust, produced from a given quantity of barley malted in the usual way, is little more than four-fifths that of the unmalted grain; about two-thirds of the loss being water, and one-third solid substance or food material.
2. The loss of solid substance consists chiefly of non-nitrogenous matters, but includes also a small amount of nitrogenous or "flesh-forming," and mineral matters.
3. A portion of the starch of the grain is converted into dextrine, and a portion of this is further converted into sugar, the amount of which is thus raised from 2 or 3 per cent. in the raw barley, to about 10 or 12 per cent. in the finished malt; and there is, besides, an actual loss of a portion of the changed starch by further decomposition as the growth proceeds.
4. The per-centage of nitrogenous or "flesh-forming" substances is higher in the diminished weight of the malted products, though the actual amount of them is less than in the raw grain from which the malt has been produced.
5. A portion of the nitrogenous substance of the grain undergoes changes by malting, by virtue of which, when the malt is digested with water, not only the previously unchanged starch of the malt itself, but the starch of a considerable amount of unmalted grain or other starchy substances mixed with it, may become converted into the more soluble forms of dextrine and sugar; the conversion taking place but slowly if cold water be employed, and the most rapidly at a temperature of about 170° F.
6. If grain were largely malted for feeding, it is probable that the growth would not be carried so far as in malting for brewing purposes, and in that case the loss of food material would be less.

The question arises—whether, by the conversion of a portion of the starch of grain into the more soluble forms of dextrine and

sugar by malting, or by the property which, under certain conditions, malted grain possesses of furthering this change, not only in its own remaining starch, but also in the starch of unmalted grain or other starchy food mixed with it, a given amount of food is thereby rendered of so much more feeding value as to compensate for the loss of food substance and the expense of the process, and leave a profit besides?

This question must be answered by means of direct feeding experiments, and we now proceed to give the results of those which have been made in the course of the inquiry to which this Report has special reference.

II.—THE FEEDING EXPERIMENTS IN 1863-4.

From the facts brought out in the foregoing section, it is obvious that, in experiments on the comparative feeding value of unmalted and malted grain, it would be quite fallacious to compare the effects of a given amount of the raw with those of an equal weight of the malted grain, without taking into account the great loss of weight in the malting. In most of the experiments to which this Report refers, therefore, a certain quantity of unmalted barley (with other appropriate food) was given to one lot of animals, and to another similar lot (having the same additional food) there was given, instead of the whole or a part of the unmalted barley, not an equal weight of the malted but only so much malt and malt-dust as were found to be produced from the quantity of raw barley for which the malted was to be substituted. In some cases, however, the barley and the malt (with its dust) were each given separately, *ad libitum*, and the amount of barley which the consumed malt and dust represented was calculated.

It was so arranged that, in some of the experiments the malt should constitute a considerable, and in others a comparatively small proportion of the total food. The data were thus provided for judging, both of the value of malt itself as a staple food compared with raw barley, and also whether or not the admixture of a certain amount of malt probably served to render other food more digestible and assimilable.

From what has been said of the properties of an infusion of malt in converting starch into the more soluble forms of dextrine and sugar, it is obvious that for the full development of the change in the food of malt-fed animals before giving it to them, it should be submitted to a process of cooking; but, if it had been decided to cook the food of the malt-fed animals, it would have been necessary, for the sake of fair comparison, to have cooked that of the barley-fed ones also; and this would have

required a duplication of most, if not all, of the feeding experiments—an extension of the inquiry which could not easily be undertaken. It was understood that the main question at issue was that of the advantage, or otherwise, of feeding animals with grain malted instead of unmalted, assuming it to be in other respects in the same condition as the unmalted grain would usually be given; and it was considered that to introduce the question of cooking would be to go beyond both the legitimate, and the most important object of the inquiry. Furthermore, as starch is known rapidly to change into sugar within the animal body, it was obviously a question whether the use of malt, if beneficial at all, might not be so by accelerating this change within the body, as, when mixed with water, it induces it without the body, especially when aided by a certain amount of heat.

The description and number of animals subjected to experiment, and the duration of each experiment, were as follow:—

20 cows, in two lots of 10 each	10 weeks.
20 oxen, in two lots of 10 each	20 "
60 sheep, in five lots of 12 each	20 "
48 pigs, in six lots of 8 each	10 "

The following is the description of the plan and results of each set of experiments.

1. *The Experiments with Milking Cows.*

Application being made to James Archibald Campbell, Esquire, of Rugby, he kindly placed his numerous herd of milking cows at disposal for the purposes of this experiment, and allowed it to be conducted in his byres, at Newlands Farm, near the town.

Two lots of 10 each having been selected, weighed, and placed apart December 1863, all received for a preliminary period of about a fortnight (during which time the milk was daily weighed) exactly the same description and amount of food, consisting of 5 lbs. rape cake and 2 lbs. bean-meal per head per day, and a mixture of clover-chaff, straw-chaff, and swedes, which averaged, of clover-chaff about 14 lbs., of straw-chaff between 7 and 8 lbs., and of swedes about 50 lbs. per head per day. The swedes were pulped, mixed with the clover and straw-chaff, the mixture slightly heated by fermentation, sprinkled with water, the meal then added, and the allowance given in three feeds daily.

On December 21, 1863, the animals were re-weighed, and with a view to greater equality in the lots, some few changes in the allotment were made; the exact experiment was then commenced, and it was continued for 10 weeks, to February 29, 1864. Both lots continued to receive the same food as previously,

excepting that Lot 1 had 3 lbs. of rape cake (per head per day), replaced by 3 lbs. of barley No. 1, and Lot 2. had 3lbs. of cake replaced by the amount of malt and malt-dust produced from 3 lbs. of barley No. 1.

The milk of each cow was weighed morning and afternoon daily. A mixed sample of the milk of each lot was tested once a week, morning and afternoon, by the lactometer; and the animals themselves were weighed at the beginning, and at the end of the fourth, eighth, and tenth weeks.

The results of all the weighings, both of the milk and of the cows, are recorded in detail in Tables X.-XIX., pp. 68-87 in the Appendix; those obtained on testing the milk in Table XX. p. 88; and the results relating both to the food consumed and to the weights, the increase in live weight, and the milk yielded, during the first four, the second four, the last two weeks, and the total period of the experiment, are given in Tables XXI.-XXIV., pp. 89-92, also in the Appendix.

The results relating to the whole period of 10 weeks are summarized in Table X., which now follows, and with slight reference to the details in the Appendix the figures there given will bring to view most of the points to which it will be necessary here to call attention.

It should be particularly observed, that in all the tables relating to these new experiments—whether with cows, oxen, sheep, or pigs, and both in the Appendix and in the body of the Report—the figures represent, wherever malt was given, not the actual weight of malt and malt-dust consumed, but the weight of barley from which (according to Table IV. p. 16) they would be produced.

In the experiments with cows it was designed to give to one lot an uniform amount of barley per head per day throughout the whole period, and to the other, also uniformly throughout the period, exactly as much malt and malt-dust as was produced from that amount of barley. Owing, however, to the gain of weight of the kiln-dried malt and dust by the absorption of moisture as the feeding experiment proceeded, some modification in the actual amounts given had to be made from time to time by way of compensation, and it was found by careful calculation at the conclusion that the malt-fed cows had in fact consumed somewhat less than the allotted amount when reckoned in the condition of dryness in which the produced malt was weighed, sampled, and its relation to the raw barley settled as in Table IV. p. 16. The slight irregularities in the actual weights given at the different periods as here explained are not recognised in the tables, but the total quantities consumed are equalized over the several periods. Due correction is, however, made for the slight deficiency in the total consumption of the

	Total in Ten Weeks.		Per Head per Day.	Per 1000 lbs. Live-weight per Week.	Cows.	Weight.		Increase in Live-weight.		Milk yielded.	
	Total in Ten Weeks.	At Commencement.				At Conclusion.	Total in Ten Weeks.	Per 1000 lbs. Live-weight per week.	Total in Ten Weeks.	Per Head per Day.	
Barley, unmalted	Lbs.	Lbs.			{ Nos 1 2 3 4 5 6 7 8 9 10 Totals Averages	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.
Barley, malted	2,100	18.4	370	12.6		1,056	1,136	50	1,311	187.8	
Cake	1,400	2.0	20	8.4		1,044	1,114	70	2,717½	31.3	
Bean-Meal	1,400	12.3	123	8.4		1,065	1,150	85	4,751½	25.1	
Clover-Chaff	9,800	12.3	123	8.4		1,052	1,150	68	1,540½	22.0	
Clover-Chaff	9,800	14.0	140	85.8		1,029	1,018	-2	1,766	20.4	146
Straw-Chaff	5,350	47.3	473	32.3		1,164	1,312	148	1,425	20.7	
Straw-Chaff	5,350	7.7	77	32.3		1,260	1,308	48	1,450	23.3	
Straw	35,258	309.2	3092	211.7		1,156	1,252	96	1,362	18.6	
Pulped Swedes		50.4				1,236	1,347	111	2,307½	32.9	

Lot 2.—10 Cows; Special Food—Malted Barley (with Malt-dust).

[illegible]

• These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

malt and dust. Thus, as the tables show, the 10 malt-fed cows consumed in the 10 weeks malt and dust equivalent to 2072 lbs. of barley, whereas they should have consumed the amount produced from 2100 lbs. of barley, a difference which it will be admitted is of little or no practical importance.

The first point to remark in regard to the results obtained with the cows is, that the barley, or the malt, constituted a comparatively small proportion of the total food consumed; certainly by no means enough to nauseate or disturb the digestion by virtue of the amount of sugar supplied when malt was given, but, on the other hand, fully sufficient to increase the digestibility of associated food materials, provided, as has been assumed, it produces that effect by virtue of its property of aiding the conversion of their starch into the more soluble conditions of dextrin and sugar.

According to the figures in the tables the 10 barley-fed cows gave, during the first period of four weeks, rather less increase in live-weight, but rather more milk, and consumed rather less food to produce a given amount of milk, than the 10 malt-fed cows; and during the second period of four, and the third period of two weeks, they gave rather more both of increase and milk, and consumed rather less food to produce a given amount of milk.

Taking the whole period of 10 weeks the 10 barley-fed cows gave 10 lbs. less increase in live-weight, but 1,067 lbs. (more than 100 gallons) more milk, and consumed less of every description of food for the production of 100 lbs. of milk than the malt-fed cows. It should be observed, however, that one of the malt-fed animals (No. 3), being in calf, increased in live-weight very considerably, and fell off very much in yield of milk, giving one third the average total amount of the rest, and that her increase, yield of milk, and proportionate amount of food, excluded by calculation (which is perhaps the fairer course) there would then be rather less increase, but rather more milk yielded, both per head and per 1000 lbs. live-weight per week, and slightly less food consumed to produce a given amount of milk by the malt-fed than by the barley-fed cows.

Thus, taken in the one way the results are in favour of the unmalted, and in the other of the malted grain. But a careful consideration of the facts will lead to the conclusion that the effects were so nearly equal as not to indicate any decided advantage of either over the other; and if there be no decided advantage from the use of a given amount of malted grain over that from the use of the amount of raw grain from which it would be produced, the economical advantage is of course with the unmalted, on account of the cost of the malting process.

On testing the milk of each lot of cows once a week, morning and afternoon, by the lactometer, that from the barley-fed animals invariably showed the higher proportion of cream, which again throws the balance somewhat in favour of the unmalted grain.

2. *The Experiments with fattening Oxen.*

These experiments, as well as those with sheep and pigs, were conducted at Rothamsted, Herts. Twenty-two three-year old polled Scots were purchased, and for about a fortnight the whole were turned out to graze during the day, and were brought into the yards at night, and supplied with clover-chaff and swedes *ad libitum*. They were then, on November 28, 1863, all weighed, two were thrown out, and the remainder were divided into two lots of 10 each, by selecting two as nearly alike as possible taking one for each lot, and so on, giving apparently the best first to one lot and then to the other, until the whole 20 were divided. The two lots were then put, two and two, into separate compartments (each communicating with an open yard) of a shed in which the experiment was to be conducted, and they were all fed for a few days longer on clover-chaff and swedes, *ad libitum*.

On December 1, 1863, the animals were re-weighed, the exact experiment was then commenced, and it was continued for 20 weeks, to April 19, 1864.

During the whole of the 20 weeks Lot 1 had 4 lbs. of barley "No. 2" per head per day, and Lot 2 the amount of malt and malt-dust produced from 4 lbs. of barley "No. 2." The barley and malt were crushed. Each lot had, besides, during the first four weeks 8 lbs. of clover-chaff, during the second four weeks 10 lbs. of clover-chaff, during the next eight weeks 2 lbs. of oilcake and 12 lbs. of clover-chaff, and during the last four weeks 4 lbs. of oilcake and 12 lbs. of clover-chaff, per head per day; also cut swedes, *ad libitum* (but weighed), throughout the whole 20 weeks. The oxen themselves were weighed every four weeks. At the conclusion of the experiment they were killed, and the weights of the carcasses, &c., ascertained.

The details of the food consumed, of the weights, and of the increase in live-weight, during each of the five periods of four weeks each, are recorded in the five Tables XXV.—XXIX., pp. 93—97, in the Appendix; the particulars relating to the weights and increase during each separate and the total period are summarised in Table XXX., p. 98; those relating to the food consumed and increase yielded in Table XXXI., p. 99; and those relating to the dead weights in Tables XXXII. and XXXIII., pp. 100, 101, also in the Appendix.

The results for the whole period of 20 weeks of feeding are further conveniently summarised in Table XI., which now follows:

TABLE XI.—RESULTS obtained with OXEN, on BARLEY and MALT “No. 2.”
 Period of Experiment 20 Weeks; from December 1, 1863, to April 19, 1864.

	Food consumed.				Oxen.	Weights, and Increase in Live-weight.			
	Total in 20 Weeks.	Per Head per Day.	Per 1000 lbs. Live-weight per Week.	To produce 100 lbs. Increase.		Weights.		Increase in Live-weight.	
						At Commencement.	At Conclusion.	Total in 20 Weeks.	Per Head per Week.
Lot 1.—10 Oxen; Special Food—Unmalted Barley.									
Barley, unmalted.	Lbs.	Lbs.	Lbs.	Lbs.	No.	Lbs.	Lbs.	Lbs.	Lbs.
Oilcake.	5,600	4.0	22.2	165.1	1	1,151	1,404	253	12.7
Clover-chaff.	2,240	1.6	8.9	66.0	2	1,079	1,386	307	15.4
Roots,* <i>ad lib.</i>	15,120	10.8	59.9	445.6	3	1,183	1,591	408	20.4
	108,146	77.2	428.6	3187.3	4	1,097	1,404	307	15.4
					5	1,095	1,422	327	16.4
					6	1,039	1,344	305	15.3
					7	1,152	1,566	414	20.7
					8	1,015	1,421	409	20.5
					9	1,102	1,452	350	19.0
					10	1,008	1,291	283	14.2
Totals.					Totals.	10,921	14,314	3,393	..
Averages.					Averages.	1,092	1,431	339.1	17.0
									134
Lot 2.—10 Oxen; Special Food—Malted Barley (with Malt-dust).									
Barley, malted with its dust.	Lbs.	Lbs.	Lbs.	Lbs.	No.	Lbs.	Lbs.	Lbs.	Lbs.
Oilcake.	5,600†	4.0†	22.5†	187.6†	1	1,204	1,525	321	16.1
Clover-chaff.	2,240	1.6	9.0	75.0	2	1,057	1,422	365	18.3
Roots,* <i>ad lib.</i>	15,120	10.8	60.7	506.5	3	1,148	1,525	377	18.9
	108,905	77.8	437.4	3643.4	4	1,088	1,396	308	15.4
					5	1,176	1,444	268	13.4
					6	1,092	1,256	164	8.2
					7	1,078	1,346	268	13.4
					8	1,029	1,252	223	11.2
					9	1,074	1,395	307	15.4
					10	1,008	1,292	284	19.2
Totals.					Totals.	10,921	14,314	3,393	..
Averages.					Averages.	1,092	1,431	339.1	17.0

in the case of the cows, so in that of the oxen (and indeed the sheep also), some allowance was made during the course of the experiment to compensate for the progressive gain of stature by the malt; but in the tables no account is taken of the extra weight given for this purpose, the figures there representing the quantities of the malt reckoned in the state of dryness in which it was originally sampled and allotted, the total quantity consumed being equalized over the several periods, as was in reality very nearly the case.

As already stated the quantity of malt and dust given per day to the malt-fed oxen was that which would be produced from 4 lbs. of barley. This, though not enough to sustain life, was more than enough on the assumption of its acting as a substitute for the digestion of other food, and it was also quite as good as any other food of any cereal grain, it being generally preferable to give a mixed food, including such things as cake or pulse.

Little need be said on the progress of the animals during each separate monthly period. As is always the case, it varied considerably among the individuals of each lot; though the average increase per head was, with each lot, comparatively uniform during each of the first three months; with both, and most with Lot 1 having the barley, it fell off considerably during the fourth period, owing, it was believed, to deterioration of the swedes, but with both it recovered considerably, more especially with Lot 1, during the last month, when the roots consisted in part of mangolds.

Directing attention to the total and average results over the whole period, as given in Table XI., it is worthy of remark that although the roots were given *ad libitum*, the two lots consumed in the 20 weeks within 7 cwts. of the same amount; the malt-fed oxen taking about that amount more than the 10 dry-fed ones. The fact is, that, if not the actual weight of food consumed, but, as in the Table, the amount of barley from which it would be produced be reckoned against the malt-fed animals, more food was expended upon them per 1,000 lbs. live-weight per week, that is in relation to a given live-weight in a given time, than upon the others; though in reality they received even slightly less solid matter, that is, reckoning malt as such instead of as representing so much barley.

Both lots of oxen gave more than an average amount of increase, whether reckoned in proportion to a given live-weight in a given time, or to a given amount of food consumed; the 10 having the unmalted barley gave during the 20 weeks 408 lbs. more increase in live-weight than those having an equal amount of the same barley malted. The result was,

that, whether only the actual weight of malt, or the weight of barley from which it would be produced, be reckoned against the malt-fed oxen, they required more total food, and more dry substance of food, to produce a given amount of increase in live-weight than the barley-fed oxen.

The difference in the amount of increase produced from a given amount of food in the two cases was indeed not great, such as it was it was unfavourable to the malted grain; and it may be observed that, on several occasions during the course of the experiments, experienced breeders and feeders who inspected the animals, whilst approving of the selection and allotment, expressed a decided opinion in favour of the appearance and handling of the barley-fed oxen, and frequently so in opposition to their own preconceived views on the subject.

At the conclusion of the 20 weeks of feeding the beasts only moderately fat; but as it was desirable to close the experiment then, they were at once sent to the butcher, in order, if possible, to gain further information as to the comparative character and ripeness of the two lots than that afforded by increase in live-weight alone.

Mr. Slater, of Kensington, who purchased the whole, was good enough to afford every facility for weighing the several parts of each animal as might be desired, and also favoured with his opinion of the quality of the meat.

The feeding experiment was concluded on April 19, 1881. The first five of the barley-fed oxen were killed on April 19, the first five of the malt-fed on April 21, the second five of the barley-fed on April 22, and the second five of the malt-fed on April 23.

The actual weights of the animals, and of some of the separated parts, are recorded in Table XXXII., and the proportion of the carcass, and of the several offal parts, in 100 weight, in Table XXXIII., in the Appendix.

The percentage of the dead or carcass weight to the live weight of each animal—the most important item to consider—is given in Table XII., which now follows :—(See next page.)

The proportion of dead or carcass weight to live-weight was to be expected, ranged rather low throughout; but it was on the average, and pretty uniformly with each pair of animals, in favour of the barley-fed over the malt-fed oxen.

Mr. Slater's report on the meat of the different lots was, that the beef of one (No. 3) of the first five of the barley-fed was very good, but that that of the remaining four was of a useful quality, and cut too pale in the lean; and that that of the second five of the same lot was not quite so good; that that of the first five of the malt-fed animals cut a very good cut

TABLE XII.—PER-CENTAGES OF DEAD OR CARCASS WEIGHTS (cold) in UNFASTED LIVE-WEIGHTS.

OXEN.

Nos.	Per Cent. Carcass in unfasted Live-weight.	
	Lot 1. Unmalted Barley (No. 2).	Lot 2. Malted Barley (No. 2).
1	53·6	51·1
2	58·0	56·4
3	57·7	53·0
4	57·7	56·0
5	56·5	57·1
6	55·7	55·3
7	54·8	53·1
8	56·5	57·7
9	58·3	56·0
10	57·6	56·6
Mean	56·6	55·6

both in the fat and the lean, and was decidedly preferable to that of the first five of the barley-fed, but that the carcasses of the other five of the malt-fed oxen were very indifferent in point of ripeness; indeed, he concluded that all required six weeks or two months more feeding to rank as first quality beef.

The result was, then, that none of the oxen were ripe enough; that the barley-fed animals were the most even in condition and quality; but that among the malt-fed ones the beef of some was decidedly superior, and that of others decidedly inferior, to that of any of the barley-fed animals. From this it would seem as if the result had been more influenced by the constitution and condition of the individual animals in the case of the malt than of the barley-fed oxen; and it is worthy of remark that of the 10 malt-fed ones it was the five that were the heaviest, and in the best condition at the commencement, and not the more backward or weakly animals, that gave the best result upon the malt diet.

To conclude, the results of the experiments with oxen may be briefly summarised as follows:—the barley-fed animals gave slightly more increase in live-weight, and a slightly higher proportion of dead-weight to live-weight than the malt-fed ones; none were fully ripe; the progress and condition of the barley-fed oxen were the more uniform, but some of the malt-fed were in better and others in worse condition than any of the barley-fed animals.

3. *The Experiments with fattening Sheep.*

From a flock of about 90 Down Wethers, about 10 months old, five lots of 12 each were selected by picking out five of nearly equal weight and character, allotting one for each pen, and so on until there were 12 in each. Each of the lots was put into a separate compartment, on rafters, under cover. For a preliminary period of about a fortnight all were fed alike on about $\frac{3}{4}$ lb. cotton cake, and $\frac{3}{4}$ lb. clover-chaff, per head per day, with cut swedes *ad libitum*.

On December 2, 1863, all were re-weighed, and the exact experiment was then commenced. From that date, for 20 weeks (to April 20, 1864), all had 1 lb. of clover-chaff per head per day, and cut swedes (or mangolds) *ad libitum*; and the respective lots had, besides, as their special food, unmalted or malted barley, as under :—

- Lot 1. Unmalted barley No. 1; for 16 weeks $\frac{3}{4}$ lb., and for the remaining four weeks 1 lb. per head per day.
- Lot 2. Malt and malt-dust from barley No. 1; for 16 weeks an amount per head per day equal to that produced from $\frac{3}{4}$ lb., and for the remaining four weeks an amount equal to that produced from 1 lb. of the barley.
- Lot 3. Unmalted barley No. 2; for 16 weeks $\frac{3}{4}$ lb., and for the remaining four weeks 1 lb. per head per day.
- Lot 4. Malt and malt-dust from barley No. 2; for 16 weeks an amount per head per day equal to that produced from $\frac{3}{4}$ lb., and for the remaining four weeks an amount equal to that produced from 1 lb. of the barley.
- Lot 5. A mixture of two parts unmalted barley No. 2, and of the malt and dust from one part of barley No. 2; for 16 weeks an amount of the mixture per head per day representing $\frac{3}{4}$ lb., and for the remaining four weeks an amount representing 1 lb. of the barley.

Both barley and malt were crushed. The sheep were weighed every four weeks; and at the conclusion of the 20 weeks they were all killed, and the weights of their carcasses, &c., ascertained.

The results relating to each of the five separate periods of four weeks each are recorded in the five Tables XXXIV.-XXXVIII. respectively; a summary of the weights, and of the increase during each and the total period, is given in Table XXXIX., and a summary relating to both food and increase in Table XL., all in the Appendix (pp. 102-115).

The results relating to the whole period of 20 weeks are summarised in Tables XIII. and XIV., which now follow. (See pp. 40, 41.)

From the plan given above it will be seen that these experiments with sheep comprised three directly comparative trials:—

1. Between Lot 1, with a given amount of barley No. 1, and Lot 2, with an equal amount of the same barley malted.
2. Between Lot 3, with a given amount of the feeding-barley No. 2, and Lot 4, with an equal amount of the same barley malted.
3. Between Lot 3, with a given amount of barley No. 2, and Lot 5, with an equal amount of the same barley, two-thirds unmalted and one-third malted.

The only deviation from this plan was, as found by calculation at the conclusion of the experiment, that Lot 2 had, during the 20 weeks, received malt equal to only 1326, instead of 1344 lbs. of barley; that is, 18 lbs. = $1\frac{1}{2}$ lb. per head too little, which, spread over 20 weeks, would be of no consequence, and was partly compensated by the extra amount of swedes consumed.

The tables of detail (in the Appendix) show that all five lots have more increase in live-weight during the first than during any of the subsequent monthly periods; and they pretty uniformly have the least in the fourth month, when the swedes were failing; did they give much more in the fifth and last month, when the allowance of barley and malt were increased. But, as will be afterwards seen, the sheep were very fairly ripe, and as fattening animals approach that condition they do not show so much increase in live-weight in proportion to their real progress in accumulation of solid matter as in the earlier stages of feeding. There was, moreover, considerable variation in the rate and amount of increase among the individual animals of each lot, but by no means more than is usual when as many as a dozen animals are fed together on the same food. In fact, the average result obtained from the twelve animals in each case, and each fed for a period of 20 weeks (see Tables XIII. and XIV.), may be taken as very fairly indicating the comparative qualities of the different foods.

All five lots of sheep gave about an average amount of increase, reckoned both in relation to a given live-weight within a given time, and to the amount of food consumed.

The amount of malt given in experiments 2 and 4, represented three times as much barley as that given in the malted state in experiment 5, and constituted a much larger proportion of the total food than in the case of either the cows or the oxen.

There was, however, apparently no advantage gained from the

Table XIII.—Results obtained with SHEEP, on BARLEY and MALT “No. 1.”
Period of Experiment 20 weeks; from December 2, 1863, to April 20, 1864.

	Food consumed.			Sheep.	Weights, and Increase in Live-weight.			
	Total in 20 Weeks.	Per Head per Day.	Per 100 lbs. Live-weight per Week.		Weights.		Increase in Live-weight.	
					At Com- mencement.	At Conclusion.	Total in 20 Weeks.	Per Head per Week.
Lot 1.—12 Sheep; Special Food—Unmalted Barley.								
				Nos.	Lbs.	Lbs.	Lbs.	Lbs.
Barley, unmalted	lbs.	lbs.	lbs.	1	112	142	30	1.60
				2	101	150	49	2.45
				3	101	154	53	2.65
				4	107	137	30	1.60
				5	100	140	40	2.00
Clover-chaff				6	92	134	42	2.10
	1,680	1.0	5.77	7	100	140	40	2.00
Roots,* <i>ad lib.</i>				8	105	162	57	2.85
				9	88	133	45	2.25
				10	119	161	42	2.10
				11	88	119	31	1.55
				12	94	133	39	1.95
				Totals.	1,207	1,705	498	..
				Averages.	101	142	41½	2.08
								1.71
Lot 2.—12 Sheep; Special food—Malted Barley (with Malt-dust).								
				Nos.	Lbs.	Lbs.	Lbs.	Lbs.
Barley, malted, with its dust .	lbs.	lbs.	lbs.	1	109	154	45	2.25
				2	94	136	42	2.10
				3	96	136	40	2.00
				4	100	140	40	2.00
				5	119	171	52	2.60
Clover-chaff				6	112	161	49	2.45
	1,660	1.0	5.75	7	94	126	32	1.60
Roots,* <i>ad lib.</i>				8	102	137	35	1.75
				9	105	160	55	2.75
				10	94	128	34	1.70
				11	96	140	44	2.20
				12	88	124	36	1.80
								1.72

TABLE XIV.—RESULTS obtained with SHEEP, on BARLEY and MALT “No. 2.”
 Period of Experiment 20 weeks; from December 2, 1863, to April 20, 1864.

Food consumed.					Sheep.	Weights, and Increase in Live-weight.				
				Weights.		Increase in Live-weight.				
Total in 20 Weeks.	Per Head per Day.	Per 100 lbs. Live- weight per Week.	To produce 100 lbs. Increase.	At Com- mence- ment.		At Con- clusion.	Total in 20 Weeks.	Per Head per Week.	Per 100 lbs. Live- weight per Week.	
Lot 3.—12 Sheep; Special Food—Unmalted Barley.										
salted . . “ “ “ “ hd. . . .	Lbs. 1,344 1,680 18,712	Lbs. 0.75 1.0 11.1	Lbs. 4.61 5.76 64.19	Lbs. 267.2 334.0 3720.1	Nos.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.
					1	112	163	51	2.55	
					2	112	158	46	2.30	
					3	114	150	36	1.80	
					4	105	147	42	2.10	
					5	105	164	59	2.95	
					6	94	130	36	1.80	
					7	98	117	19	0.95	
					8	84	120	36	1.80	
					9	104	140	36	1.80	
					10	104	152	48	2.40	
					11	88	142	54	2.70	
					12	86	126	40	2.00	
Totals .					1,206	1,709	503	
Averages					100	142	42	2.10	1.73	
Lot 4.—12 Sheep; Special Food—Malted Barley (with malt-dust).										
fed, with } “ “ “ “ } hd. . . .	Lbs. 1,232† 1,540 16,854	Lbs. 0.75† 1.0 10.9	Lbs. 4.61† 5.77 63.12	Lbs. 275.0† 343.9 3762.1	Nos.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.
					1	119	150	31	1.55	
					2	114	164	50	2.50	
					3	94	134	40	2.00	
					4†	(102)	(95)	(-7)	—	
					5	108	157	49	2.45	
					6	91	128	37	1.85	
					7	107	144	37	1.85	
					8	91	140	49	2.45	
					9	102	140	38	1.90	
					10	94	126	32	1.60	
					11	100	146	46	2.30	
					12	91	130	39	1.95	
Totals .					1,111	1,559	448	
Averages					101	142	40‡	2.04	1.68	
Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley (with malt-dust).										
salted . . fed, with } “ “ “ “ } hd. . . .	Lbs. 896 448† 1,680 18,598	Lbs. 0.5 0.25† 1.0 11.1	Lbs. 3.14 1.57† 5.88 65.10	Lbs. 188.6 94.3† 353.7 3915.4	Nos.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.
					1	128	186	58	2.90	
					2	114	126	12	0.60	
					3	91	130	39	1.95	
					4	104	142	38	1.90	
					5	94	140	46	2.30	
					6	96	145	49	2.45	
					7	94	136	42	2.10	
					8	102	146	44	2.20	
					9	91	134	43	2.15	
					10	98	133	35	1.75	
					11	88	116	28	1.40	
					12	91	132	41	2.05	
Totals .					1,191	1,666	475	
Averages					99	139	39‡	1.95	1.66	

† the first four months; about one-fourth swedes and three-fourths mangolds the last month.
 ‡ Figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they
 † reduced.
 ‡ Sheep of this lot was killed 63 days after the commencement of the experiment, and its weights, and the pro-
 † portion of food, are excluded.

larger proportion of malt in the food ; for, although the sheep gave about an average, the oxen gave more than an average amount of increase.

There was, too, with the sheep, as with the oxen, comparatively little difference in the result obtained with the unmalted and malted grain. But such as it was it was more in favour of the unmalted than of the malted barley. Thus, in the comparative experiment with barley No. 1 (Lot 1 unmalted and Lot 2 malted) there was slightly more average increase per head, and, reckoning only the actual weight of the malt consumed and the amount of barley from which it would be produced, again for the animals, there was slightly less both of fresh food and of solid matter of food consumed to produce a given amount of increase with the malted barley. But in the similar comparative experiment with barley No. 2 (Lot 3 unmalted and Lot 4 malted) the average increase per head was in a somewhat greater degree in favour of the unmalted, and there was also with it less food expended for the production of a given amount of increase. With sheep, was, too, somewhat less average increase per head, and a larger quantity of food required to produce a given amount of increase, when two-thirds of the barley was unmalted and one-third malted (Lot 5) than when an equal amount of the barley was given entirely unmalted (Lot 3). There were, with both, both a less average increase per head, and a greater amount of food required to produce a given amount of increase when one-third (Lot 5), than where the whole of the barley (Lot 3) was malted ; still, the fact is that, in both respects, both lots of malted barley (4 and 5) gave worse results than Lot 3, which was the same description of barley entirely unmalted.

With sheep, therefore, as with oxen, there was, upon the whole, less food expended to produce a given amount of increase in weight when unmalted than when a corresponding amount of malted barley was given. The differences were, indeed, slight, but the economical advantage is, of course, more in favour of the unmalted grain than the figures show, when the cost of the malting process is taken into the calculation.

At the conclusion of the feeding experiment all five sheep were sold to Mr. Slater, and they were all killed within a few days of each other. The feeding experiment ended on April 20. On April 21 five of Lot 1, five of Lot 2, and five of Lot 3, were killed ; on April 22 six of Lot 4, six of Lot 5, and one each of Lots 1, 2, and 3, were killed ; and on April 23 the remaining six of Lot 1, six of Lot 2, six of Lot 3, five of Lot 4, and six of Lot 5, were killed.

The actual dead weights are recorded in Table XLI., and the percentages of the carcasses and other parts, in the live-weights, in Table XLII., pp. 116-119, in the Appendix.

In the following Table (XV.) are summarised the per-centages the dead or carcass weights in the unfasted live-weights of the sheep of each lot:—

TABLE XV.—PER-CENTAGES OF DEAD OR CARCASS WEIGHTS (cold) in UNFASTED LIVE-WEIGHTS.

SHEEP.

Sheep. Nos.	Lot 1.	Lot 2.	Lot 3.	Lot 4.	Lot 5.
	Barley No. 1.		Barley No. 2.		
	Unmalted.	Malted.	Unmalted.	Malted.	$\frac{2}{3}$ Unmalted, $\frac{1}{3}$ Malted.
1	51·4	52·0	53·4	48·7	52·2
2	49·3	47·8	51·3	48·8	52·4
3	51·3	50·0	56·0	50·8	51·5
4	53·3	51·4	55·1	*	50·7
5	49·3	52·6	51·8	51·6	47·1
6	47·8	47·2	54·6	51·6	51·7
7	52·9	54·0	54·7	49·3	50·7
8	51·9	54·0	52·5	49·3	54·1
9	54·9	53·1	54·3	52·9	52·2
10	50·9	50·8	50·0	55·6	49·6
11	50·4	53·6	50·0	51·4	50·0
12	55·6	51·6	53·2	51·5	52·3
Means ..	51·6	51·5	53·1	51·0	51·2

The sheep being sold and killed without being shorn, their e-weights of course included the wool, and hence the low proportion of dead or carcass weight to live-weight in all cases. Nor is the average proportion very different with the different lots. Such as it was, however, it was slightly in favour of the unmalted barley. The highest proportion was given by Lot 3 fed on the malted barley No. 2. That barley, indeed, contained, as will be remembered, considerably the higher per-centage of nitrogen, and was doubtless of very good feeding quality, though much inferior to the other for the manufacture of malt for brewing purposes.

Mr. Slater reported that all the sheep were very good both as regards size and quality; and he did not see reason to draw any distinction between the different lots further than was indicated by the weights.

Upon the whole the experiments with sheep showed very little difference between the effects of a given amount of barley, and of the malt and malt-dust produced from an equal weight of the

* No. 4. Sheep of this lot was killed 63 days after the commencement of the experiment.

same barley. Such as it was the advantage was with the unmalted barley, both in regard to the amount of increase in live weight obtained, and to the proportion of dead-weight to live weight.

4. *The Experiments with fattening Pigs.*

Fifty-two pigs were purchased for the purpose of the experiments, and all were fed for a preliminary period of about a fortnight on a mixture of equal parts barley-meal and bran, given *ad libitum*. They were then weighed, and six lots, of eight each were selected, in the same manner as for the sheep, and all again fed for a few days on the same mixture of barley-meal and bran.

On December 24, 1863, all were re-weighed, and the experiment was commenced, and it was continued for 10 weeks to March 3, 1864. Throughout the whole period all had 1 lb. of pea-meal per head per day, and the respective lots had, in addition, special food, as under:—

Lot 1. Unmalted barley No. 1, *ad libitum*.

Lot 2. The malt (with its dust) from barley No. 1, *ad libitum*.

Lot 3. Unmalted barley No. 1, and the malt (with its dust) from barley No. 1, each separately, *ad libitum*.

Lot 4. Unmalted barley No. 2, *ad libitum*.

Lot 5. The malt (with its dust) from barley No. 2, *ad libitum*.

Lot 6. A mixture of four parts unmalted barley No. 2 with the malt and malt-dust from one part of barley No. 2, *ad libitum*.

The barley and the malt were each coarsely ground; and enough to last for several days being weighed out for each lot, and was mixed with water, and so given as needed. The daily allowance of pea-meal was mixed with a portion of the other food. The animals were weighed fortnightly, and at the conclusion of the 10 weeks they were all killed, and the weights of their carcasses ascertained.

Barley-meal, with or without a small proportion of pea-meal, is recognised as perhaps the most appropriate fattening food for the pig. Such a diet contains a very much larger proportion of starch than that of either cows, oxen, or sheep; and if malt given as food acted in any material or striking degree by reason of its own remaining starch, or that of accompanying food, becoming more readily digestible and assimilable, it might be assumed that it would be particularly advantageous in the characteristically starchy food of the fattening pig.

Analysis has shown that kiln-dried malt may contain nearly two-thirds its weight of unaltered starch and starch changed only into the more soluble form of dextrine, and not more than from

10 to 12 per cent. of ready-formed sugar. The question arises, what proportion of the food, to act the most advantageously, should consist of malted grain, provided it were advantageous to employ it at all?

Considering that so small a proportion of the starch of raw barley is converted into sugar by malting, it seemed desirable, at any rate in some of the experiments, to give nearly the whole of the meal in the state of malt, and in others to give much less, in admixture with a considerable amount of raw grain, or other starchy food. Accordingly, as shown by the plan of the experiments with pigs, given above, beside the 1 lb. of pea-meal given per head per day to each lot of pigs, Lot 1 had barley No. 1, *ad libitum*, and Lot 2 the malt (and dust) from the same barley, *ad libitum*; thus substituting the whole of the unmalted grain of Lot 1 by malted grain, whilst the animals of Lot 3 were allowed to take of the unmalted or the malted grain at pleasure. Again, Lot 4 had barley No. 2, *ad libitum*; Lot 5 the malt (and dust) from barley No. 2, *ad libitum*; and Lot 6 a mixture of four-fifths unmalted and one-fifth malted barley No. 2, *ad libitum*.

The result of this arrangement was, that in the cases of Lots 2 and 5, the one with the first and the other with the second quality malt, *ad libitum*, the malt contributed from 85 to 90 per cent. of the total dry or solid matter consumed; that in experiment 3, in which the pigs took the unmalted or the malted grain at pleasure, the dry or solid matter of the malt consumed amounted, taking the average of the whole period, to only about 13 per cent. of the total solid matter of the food; and that in experiment 6, in which only one-fifth of the barley was given in the malted state, the malt contributed only between 16 and 17 per cent. of the dry or solid substance consumed.

For the sake of comparison with the above statements, it may be observed that in the experiment with cows the malt contributed only about $7\frac{1}{2}$, in that with oxen between 13 and 14, in experiments 2 and 4 with sheep between 22 and 23, and in experiment 5 with sheep only about $7\frac{1}{2}$ per cent. of the total solid matter of the food consumed. Taking all the experiments together, therefore, the proportion of the total food given as malted grain varied very considerably, contributing in several cases only from 7 to 8, and in others nearly 90 per cent. of the total solid substance of the food of the animals.

It may be further remarked on this point, that although taking the average of the whole period the pigs of Lot 3, which settled for themselves the proportion of malt in their food, took only about 13 per cent. of the total solid substance in the form of malt, yet during the first fortnight they took $31\frac{1}{2}$, during the

second 19½, during the third only 3½, during the four during the fifth 6½ per cent., giving over the total 10 weeks the average of about 13 per cent., as above stated. It would thus appear that, pig-like, they took of the sweet grain to nauseation at first; they then reduced their consumption of it to a minimum, and afterwards gradually increased to a slight degree.

The details of the feeding experiments with pigs are Tables XLIII.-XLIX., and those relating to their deaths in Tables L.-LI. in the Appendix, pp. 120-137.

The results of the feeding experiments are summarized in Tables XVI. and XVII., which now follow. (See pp.

Leaving the details given in the Appendix for reference, will suffice to call attention here to the average results over the total period of ten weeks in each experiment.

The pigs of Lots 1, 3, 4, and 6, which, besides the pea-meal per head per day given to all, had, as their additional food, either unmalted barley entirely, or only a small portion of malted barley, gave fully an average amount of increase in relation to their weight within a given time an amount of food consumed. Those of Lots 2 and 5, on the other hand, which had as their additional food malted barley gave a defective result in both respects; those of Lot 2, the malt from the barley No. 1, being the worst.

There is no doubt that in the two last-mentioned experiments (2 and 5), the proportion of saccharine matter in the food was unnaturally large. It was probably somewhat so in experiment 6, in which the additional food consisted of both unmalted barley and the malt and dust from one part of malt for when, as in experiment 3, the unmalted and malted were each given separately, *ad libitum*, although at first the animals took more than one-third from the malted, they then took very much in their consumption of it, until, averaged over the whole ten weeks of the experiment, they had not taken more than one-sixth in the malted state. Even, however, in the experiments in which not more than one-fifth or one-sixth of the additional food, or not more than one-sixth or one-seventh of the additional food was in the form of malted barley, rather more food was required to produce a given amount of increase than when the additional food was unmalted barley.

The pigs were not sent to London, but were sold at auction in the neighbourhood. The animals of pens 1, 2, 4, and 5 were killed in the evening of March 3, the feeding experiments concluded, and the last live-weights having been taken in the morning of that day; but the pigs of pen 6 were not killed until March 7, nor those of pen 3 until March 9, so that

TABLE XVI.—RESULTS obtained with PIGS, on BARLEY and MALT "No 1."

Period of Experiment 10 weeks; from December 24, 1863, to March 3, 1864.

Food consumed.					Pigs.	Weights, and Increase in Live-weight.				
Total in 10 Weeks.	Per Head per Day.	Per 100 lbs. Live- weight per Week.	To produce 100 lbs. increase.	Weights.		Increase in Live-weight.				
				At Commencement.		At Conclusion.	Total in 10 Weeks.	Per Head per Week.	Per 100 lbs. Live- weight per Week.	

Lot 1.—8 Pigs; Food—Pea-meal, and Unmalted Barley.

	lbs.	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.	lbs.
mal . . .	560	1.00	3.4	47.6	1	158	287	129	12.9	7.63
mal, unmalted, & malt-dust, and	4,902	8.75	29.3	416.8	2	160	336	176	17.6	
					3	133	261	131	13.1	
					4	134	287	153	15.3	
					5	135	270	135	13.5	
					6	122	297	175	17.5	
					7	122	241	119	11.9	
					8	122	250	158	15.8	
Totals .						1,086	2,262	1,176
Averages						136	283	147	14.7	7.03

Lot 2.—8 Pigs; Food—Pea-meal, and Malted Barley.

	lbs.	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.	lbs.
mal . . .	560	1.00	3.7	67.9	1	172	290	118	11.8	5.50
mal, malted, & malt-dust, and	4,331*	7.82*	29.2*	531.0*	2	144	278	134	13.4	
					3	142	242	100	10.0	
					4	133	240	107	10.7	
					5	130	224	91	9.4	
					6	124	228	104	10.4	
					7	119	214	95	9.5	
					8	123	196	73	7.3	
Totals .						1,087	1,912	825
Averages						136	239	103	10.3	5.50

Lot 3.—8 Pigs; Food—Pea-meal, Unmalted and Malted Barley (with malt-dust).

	lbs.	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.	lbs.
mal . . .	490	1.00	3.4	50.8	1	147	250	133	13.3	6.73
mal, unmalted, & malt-dust, and	3,491	7.12	24.3	361.7	2	150	287	137	13.7	
					3†	(150)	(163)	(13)	—	
					4	144	282	138	13.8	
					5	129	273	144	14.4	
					6	130	280	150	15.0	
					7	121	284	160	16.0	
					8	128	231	103	10.3	
Totals .						952	1,917	965
Averages						136	274	138	13.8	6.73

These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which could be produced.
No. 3 pig of this lot was killed 21 days after the commencement of the experiment; and its weights, and the total amount of food, are excluded.

TABLE XVII.—RESULTS obtained with PIGS, on BARLEY and MALT “No.
Period of Experiment 10 weeks from December 24, 1863, to March 3, 1864.

Food consumed.					Pigs.	Weights, and Increase in Live-w			
Total in 10 Weeks.	Per Head per Day.	Per 100 lbs. Live- weight per Week.	To produce 100 lbs. Increase.	Weights.		Increase in Live-			
				At Com- mence- ment.		At Con- clusion.	Total in 10 Weeks.	Per Head per Week.	

Lot 4.—8 Pigs; Food—Pea-meal, and Unmalted Barley.

	lbs.	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.
Pea-meal . . .	560	1.00	3.5	52.2	1	150	273	123	12.3
Barley, unmalted, <i>ad lib.</i> . . . }	4,527	8.08	27.9	421.9	2	147	282	135	13.5
					3	142	306	164	16.4
					4	140	262	122	12.2
					5	136	259	123	12.3
					6	126	245	119	11.9
					7	133	280	147	14.7
					8	112	252	140	14.0
Totals .						1,086	2,159	1,070	..
Averages						136	270	134	13.4

Lot 5.—8 Pigs; Food—Pea-meal, and Malted Barley (with Malt-dust).

	lbs.	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.*	lbs.
Pea-meal . . .	560	1.00	3.5	55.5	1	166	284	128	12.8
Barley, malted, with its dust, <i>ad lib.</i> . . . }	5,289*	9.44*	32.7*	524.2*	2	152	294	142	14.2
					3	150	298	148	14.8
					4	130	234	104	10.4
					5	135	262	127	12.7
					6	126	270	144	14.4
					7	134	228	94	9.4
					8	130	252	122	12.2
Totals .						1,113	2,122	1,009	..
Averages						139	265	126	12.6

Lot 6.—8 Pigs; Food—Pea-meal, and mixture of four-fifths Unmalted and one-fifth malted Barley (with Malt-dust) *ad lib.*

	lbs.	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.
Pea-meal . . .	420	1.00	3.4	50.2	1	168	350	182	18.2
Barley, unmalted .	2,929	6.97	24.0	350.4	2†	(157)	(262)	(165)	—
					3	133	250	117	11.7
					4	128	270	142	14.2
					5†	(134)	(134)	(0)	—
					6	136	283	147	14.7
					7	128	250	122	12.2
					8	112	238	126	12.6
Barley, malted, with its dust . . }	732*	1.74*	6.0*	87.6*					
Totals .						805	1,641	836	..
Averages						134	273	139	13.9

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which would be produced.

† No. 3 pig of this lot died 43 days after the experiment commenced, and No. 5 pig was killed on the 10th of the experiment. Their weights, and the proportional amount of food, are excluded.

cases the live-weights upon which the per-centages of the carcasses are calculated were taken some days later than the conclusion of the exact feeding experiment.

The actual live and dead weights are given in Table L., and the per-centages of the dead weights in the live-weights in Table LL., pp. 134-137 in the Appendix.

The per-centages of carcass or dead weight in unfasted live-weight are summarised in Table XVIII., which now follows:—

TABLE XVIII.—PER-CENTAGES OF DEAD OR CARCASS-WEIGHTS (cold) in UNFASTED LIVE-WEIGHTS.

PIGS.

Nos.	Lot 1.	Lot 2.	Lot 3.	Lot 4.	Lot 5.	Lot 6.
	Barley No. 1.			Barley No. 2.		
	Unmalted.	Malted.	Unmalted and Malted; each, <i>ad lib.</i>	Unmalted.	Malted.	Mixture: $\frac{4}{5}$ Unmalted, $\frac{1}{5}$ Malted.
1	79.1	78.4	76.5	78.3	80.2	78.7
2	80.7	78.2	76.2	80.0	78.0	*
3	77.3	77.7	*	77.4	78.9	76.6
4	78.1	82.9	78.2	79.9	77.1	77.9
5	77.0	77.5	77.4	78.8	78.5	*
6	80.1	79.0	78.0	80.0	77.8	75.0
7	82.6	75.0	78.1	78.7	77.9	80.5
8	79.1	80.1	77.5	76.2	75.0	78.3
Means.	79.3	78.6	77.4	78.6	77.9	77.8

The pigs had increased in frame a good deal whilst feeding on bran and barley-meal before the commencement of the experiments, and, for their size, none of them were very fat at the conclusion. The table shows that the average proportion of carcass was greater in pen 1, with barley No. 1 unmalted, than in either pens 2 or 3, with the same barley either entirely or in part malted. It was also greater in pen 4, with barley No. 2 unmalted, than in either pens 5 or 6, with the same barley wholly or in part malted. With both qualities of grain, however, there was a slightly better average proportion of dead weight where the whole than where only a part of the additional food consisted of malt. But it will be remembered that the amount of increase and its proportion to food consumed were notably greater with the smaller proportion of malt, and as the

* No. 3 pig of Lot 3 was killed 21 days after the commencement of the experiment; No. 2 pig of Lot 6 died 43 days after the experiment commenced; and No. 5 pig (Lot 6) was killed on the 15th day of the experiment.

two Lots 3 and 6, which had the smaller proportion, were those which were not killed until several days after the conclusion of the feeding experiment, there is not the same reliance to be placed in the proportion of the carcass to the unfasted live weights, the latter being at any rate different from those taken at the conclusion of the feeding experiment, after which the feeding was possibly not quite so regular, and the animals perhaps fuller and, therefore, heavier in proportion to the carcass, at the time their final unfasted weights were taken.

An inspection after killing showed the comparative character of the carcasses to be as follows:—

- Pen 1, with barley No. 1 unmalted;—much alike, of very good quality, but not fully ripe.
- Pen 2, with barley No. 1 malted;—uneven, pig No. 1 prettily ripe, of good colour, and had a good flare, as also had pig 2, whilst Nos. 4, 5, 6, 7, and 8 appeared to have been wasting, and, as will be seen by reference to Table XLVIII. p. 130 in the Appendix, those animals, especially pig No. 4 (which, however, gave a high proportion of dead weight), fell off very much in rate of increase during the last few weeks of feeding.
- Pen 3, with barley No. 1 unmalted and malted each *ad libitum*;—pork very firm, and of good quality; flesh very white.
- Pen 4, barley No. 2 unmalted;—pork very good, and but little difference among the individuals of the pen.
- Pen 5, barley No. 2 malted;—pork of very good quality.
- Pen 6, barley No. 2, four-fifths unmalted and one-fifth malted;—pork very white in the flesh, and of very good quality.

The general result with the pigs was:—that more increase in live-weight, and more meat, were obtained from a given amount of barley unmalted than from an equal amount of the same barley malted; that the progress of the animals, and the quality of the meat, were less uniform when a considerable quantity of malted barley was given; that when only a moderate quantity of malt was given the quality of the meat was very good; and lastly, that, considering the difference in the amount of increase in live-weight yielded for a given amount of food consumed there was comparatively little difference in the proportion of dead weight to live with the unmalted and the malted barley; but such as it was it was in favour of the unmalted.

SUMMARY.

The results of the whole inquiry may be briefly enumerated as follows:—

The Loss and Chemical Changes which the Grain undergoes by malting.

1. In malting barley of fair malting quality, in the usual way, there was a loss of nearly 19 per cent. of its weight, about 12 of which were water, the remaining 7 being solid matter or food-material.

2. In malting barley of good feeding but inferior malting quality, there was a loss of about 22 per cent. of its weight, of which 15 were water, and 7 solid matter or food material.

3. The loss of solid matter consisted chiefly of starch, or other non-nitrogenous substances, but comprised also a small amount of nitrogenous or "flesh-forming," and mineral matters.

4. The most characteristic change which the grain undergoes by malting is the conversion of a portion of its starch into dextrine, and the further conversion of a portion of the latter, amounting to from 8 to 10 per cent. of the grain, into sugar.

5. By malting, the grain acquires properties, by virtue of which, when the malt is digested with water, much of its own remaining starch gradually changes into dextrine and sugar; and if the digestion be aided by heat, not only the whole of the remaining starch of the malt itself, but the starch of a considerable quantity of unmalted grain or other starchy substances mixed with it, may become so converted.

6. Owing to the great loss of moisture and non-nitrogenous substances—in fact, of total weight—which grain undergoes by malting, a given weight of the malted grain contains a larger quantity of nitrogenous or "flesh-forming" substances than an equal weight of the unmalted grain; but as there is an actual loss of those substances by malting, a given weight of malt will, of course, contain less of them than the amount of barley from which it was produced.

Malting and the use of Malt for feeding.

7. It is probable that if grain were malted extensively for feeding purposes, the growth would not be carried so far as in

the manufacture of malt for brewing, and the loss of solid material or food material would, of course, be less accordingly.

8. As the "malt-dust" contains a considerable amount of food material, abstracted from the grain during growth, when malt is used for feeding the "dust" should either not be separated, or, if separated, should be given to the animals also with the screened malt.

9. Owing to the loss of weight which grain undergoes in malting, equal weights of malted and unmalted grain should not be employed in comparative feeding experiments, but only such much malt (with the dust) as would be produced from the same amount of raw grain given, or to be substituted, in the parallel experiment.

10. Malt given as food to animals may be supposed to be equivalent simply by supplying more or less of the starch of the grain from which it was produced in the more soluble and, perhaps, therefore, more easily digestible conditions of dextrine and sugar, also by aiding the conversion into dextrine and sugar of the starch of other foods given with it.

The Experiments with milking Cows.

11. A comparative experiment was made in which, besides other appropriate food, 10 cows received, for a period of 10 weeks, 3 lbs. of fair malting barley per head per day, and another 10 received the amount of malt (with its dust) produced from 3 lbs. of barley from the same stock.

12. In the experiment in which the malt was given it contributed about $7\frac{1}{2}$ per cent. of the solid matter of the total food.

13. The result was, that almost exactly the same amount of milk was yielded for a given amount of food with the unmalted barley and with the malted barley, but that the milk from the cows having the unmalted barley contained the higher proportion of cream.

The Experiments with fattening Oxen.

14. A comparative feeding experiment was made for a period of 20 weeks, in which, with other appropriate food, 10 oxen received 4 lbs. of good feeding barley per head per day, and another 10 the amount of malt (with its dust) produced from 4 lbs. of barley from the same stock.

15. In the experiment with malt it contributed about $13\frac{1}{2}$ per cent. of the dry or solid substance of the food.

16. Both lots of oxen gave more than an average amount of increase, whether reckoned in proportion to a given live-weight within a given time, or to a given amount of food consumed.

but the 10 having the unmalted barley gave rather more than those having the malted.

17. The barley-fed oxen also gave rather the higher proportion of dead-weight to live, and, although neither lot was fully ripe, the barley-fed animals were more even in condition and quality than the others; but the beef of some of the malt-fed ones was decidedly superior in point of ripeness and quality, and that of others decidedly inferior to that of any of the barley-fed oxen.

18. It would seem, therefore, that the effect of the malt as food was more dependent on the constitution and condition of the individual animals than was that of the barley; and it should be remarked that the oxen which fattened the best upon the malt were not the most backward or weakly animals, but those which were the heaviest and in the best condition at the commencement.

The Experiments with fattening Sheep.

19. Comparative experiments were made for a period of 20 weeks with five lots of sheep, of 12 each. Besides other appropriate food given equally to all, the allowance per head per day was—to Lot 1 from $\frac{3}{4}$ to 1 lb. of fair malting barley; to Lot 2 the malt (with its dust) from an equal amount of the same barley; to Lot 3 from $\frac{3}{4}$ to 1 lb. of good feeding barley; to Lot 4 the malt (with its dust) from an equal amount of the same barley; and to Lot 5 an equal amount of the same barley, two-thirds unmalted and one-third malted.

20. In experiments 2 and 4, the malt contributed about $22\frac{1}{2}$ per cent., and in experiment 5 about $7\frac{1}{2}$ per cent., of the dry or solid substance of the food.

21. All five lots of sheep gave about an average amount of increase; there was very little difference in the result obtained with the unmalted and the malted grain; but such as it was it was rather in favour of the unmalted.

22. The mutton of all five lots was of very good quality; there was no appreciable difference between the lots in this respect, but the barley-fed animals gave slightly the higher proportion of dead-weight to live-weight.

The Experiments with fattening Pigs.

23. The appropriate food of the fattening pig contains a larger proportion of starch than does that of either cows, oxen, or sheep. If, therefore, the starch of food be rendered more digestible and assimilable by its artificial conversion into the more soluble forms of dextrine and sugar, it might be supposed

that it would be peculiarly advantageous to malt a part, or whole, of the characteristically starchy food of the fattening

24. Experiments were made for a period of 10 weeks with lots of pigs of eight each. Besides 1 lb. of pea-meal per pig per day given to all—Lot 1 had crushed malting barley, Lot 2 the crushed malt (with its dust) from the same barley, and Lot 3 the unmalted and the malted barley, each separately, *ad libitum*. Lot 4 had crushed feeding barley, Lot 5 the crushed malt (with its dust) from the same barley, Lot 6 the same barley, four-fifths unmalted and one-fifth malted, *ad libitum*.

25. In experiment 2 the malt contributed 87½, in experiment 3 about 13, in experiment 5 about 89, and in experiment 6 at 16½ per cent., of the dry or solid substance of the food.

26. The pigs having pea-meal and entirely unmalted barley (Lots 1 and 4) gave a full average amount of increase, both in relation to a given live-weight within a given time, and to a given amount of food consumed; those having only a small portion of malted barley (Lots 3 and 6) increased in weight in these respects, nearly, but not quite, as well; but those having the pea-meal and entirely malted barley (Lots 2 and 5, more especially Lot 2), gave less increase in relation to a given live-weight within a given time, and required the expenditure of considerably more barley to produce a given amount of increase.

27. The pigs having the unmalted barley (Lots 1 and 4) gave the best average proportion of dead-weight to live-weight and their pork was of very good quality; and with the exception of Lot 2 having (besides the pea-meal) entirely malted barley No. 1, the pork of the other lots was also of very good quality, but the more evenly so where only a small proportion of malt was given (Lots 3 and 6).

General Conclusions.

28. The general conclusion from the results of the different experiments with cows, oxen, sheep, and pigs, is, that a given weight of barley is more productive, both of the milk of cows and of the increase in live-weight of fattening animals, than an equal amount of malt and malt-dust that would be produced from it.

29. The results of these new experiments, as here stated, are consistent with those obtained in an official inquiry conducted in 1845-6, by the Drs. Thomas and Robert Dundas Thomson with cows and with oxen. They are consistent with the results of experiments made at Rothamsted, in 1848 and 1849, with sheep; and also with those of others made in 1854 with pigs, in which some were fed on sugar.

30. Wherever weights have been taken as a measure of effects produced, experience hitherto has failed to show

advantage in malt over the amount of barley from which it would be produced, as a staple food for healthy milking cows or fattening animals ; and, if no advantage, there must, in point of economy, be a loss, on account of the cost of the malting process.

31. Irrespectively of economy, malt is undoubtedly a very good food for stock ; and common experience seems to show that a certain amount of it is beneficial when given in admixture, and in change, with other food, to young or weakly animals, or in "making up" or "finishing" for exhibition or sale ; that is, when the object is to produce a particular result irrespective of the economy required in ordinary feeding.

JOHN BENNET LAWES.

Rothensted, Herts, August, 1865.

APPENDIX.

The Determination of the Sugar in the Barley, and in the Products of the Malting Process.

As each of the several methods for the quantitative estimation of sugar seemed possibly open to some objection when applied to complex solutions such as the extract of malted grain, or malt-dust, it was decided, for the sake of confirmation, in all cases to adopt two methods, and always to make at least two experiments with each. The following is an outline of the methods adopted; and in Table 1 (p. 59) are given the results of each individual experiment.

1. Determination by the Fermentation and Alcohol Method.

The grain, unmalted or malted as the case might be, being finely ground, 630 grains were stirred up with 200 septems* of cold water, and the mixture allowed to stand for about an hour. It was then well rubbed in a mortar, and transferred to a bottle, 1000 septems of water being used in rinsing the mixture into the bottle, and then 500 septems more were added, making in all 1700 septems of water and 630 grains of substance. The mixtures were generally made about the middle of the day, well shaken at intervals throughout the afternoon, and then allowed to stand to settle till the next morning, when as much as possible of the supernatant liquid was removed by means of a syphon. The solutions were turbid, but did not react with iodine. To 1000 septems 100 septems of lime-water were added, and the mixture was very slightly warmed to expedite precipitation, after which there remained a perfectly clear, but coloured supernatant liquid. Of this 700 septems were taken for fermentation, and the remainder was left for the determination of the sugar by the copper method.

In the preparation of the extract from the malt-dust some deviation from the above mode of procedure was made. In its case, 200 septems of milk of lime were added to 800 septems of the original infusion, and, after filtration, only 600 septems of the liquid were taken for fermentation, the remainder, as before, being left for the copper method.

From the above figures it results that the extract submitted to fermentation represented in the case of the malt-dust, 177.9 grains, and in that of the barley and of the other products of the malting process, 235.8 grains of original substance.

The yeast employed was pressed in a cloth, or between blotting paper, and then well mixed before being weighed out for use. Of the

* 1 septem measure = 7 grains, or $\frac{1}{1000}$ th of a pound avoirdupois, of water.

Two prepared yeast 90 grains were employed for each fermentation ; and two lots, of 90 grains each, were always mixed with water, and left to ferment side by side with the fermenting extracts, the whole being maintained as nearly as possible at a temperature of 78° F.

At the conclusion of the fermentation, each fluid was submitted to distillation, and the distillate was weighed in a 1000-grain bottle, in successive quantities as it was collected, until the specific gravity showed that only pure water came over. The sum of the attenuation of the several separately weighed lots of the distillate, less that of the distillate from the yeast fermented with pure water, gave the total attenuation in 1000 grain measures due to the alcohol formed from the sugar of the substance experimented upon. The amount of proof spirit which 1000 grain measures of spirit of the attenuation thus found being ascertained by reference to Bate's Tables (and interpolation), it only remained to calculate the amount of alcohol which that amount of proof spirit represented, and then the amount of sugar (dry malt- or grape-sugar = $C_{12} H_{12} O_{12}$) to which the amount of alcohol was equivalent, thus :—

$$\frac{x \times 180}{92} = \text{malt-sugar.}$$

2. Determination by the Copper Method.

A standard solution was made by dissolving, separately, in water :—

245 grains of crystallized sulphate of copper,

700 grains of crystallized tartaric acid,

840 grains of fused caustic soda,

making the solutions, and making up with water to 1000 septems at 52° F.

According to calculation, on the assumption that 1 equivalent of grape sugar would reduce 10 equivalents of oxide of copper, 100 septems of this solution should indicate 3.535 grains, or 1 septem .03535 grain of grape-sugar.

Another standard solution was made by dissolving—

242.98 grains of crystallized sulphate of copper,

700 grains of crystallized tartaric acid,

840 grains of fused caustic soda,

and making up to 1000 septems. Of this solution 100 septems represented, by calculation, 3.506 grains, or 1 septem .03506 grain of grape-sugar.

The actual value of each of these solutions was determined from time to time by means of a solution made by dissolving 10 grains of pure cane sugar in 2 or 3 ounces of water, adding 20 or 30 drops of strong sulphuric acid (previously diluted), boiling for a short time to convert the cane sugar into grape-sugar, and when cold making up to 500 septem measures with water. Each septem of this solution represented, therefore, .02 grain of cane-sugar, or

$$\frac{.02 \times 180}{171} = .02105 \text{ grain dry grape-sugar.}$$

In testing the value of the solutions, or in actual working, 50 or 100 septem measures of the copper solution were put into a small flask and heated by means of a water bath. The solution of sugar, or prepared grain-extract, was then allowed to flow in by degrees from a burette, until the point of saturation was attained. Even with the solutions of pure sugar, it was difficult to determine by the eye when the whole of the copper was precipitated, nearer than by one or two septems of the solution; and with the coloured grain-extracts the difficulty, and range of error in reading, were, of course, increased. In practice it was found necessary, as the point of saturation was approached, from time to time to remove a few drops, filter, and test by a solution of yellow prussiate of potass.

The Determination of the Woody-fibre, in the Barley, and in the Products of the Malting Process.

For each determination 12 grammes (185·2 grains) of the finely ground substance were taken. 1200 septems of dilute sulphuric acid made by mixing 1 volume of oil of vitriol and 16 volumes of water were put into a flask with a wide mouth closed by means of a small funnel to prevent loss by evaporation, and then heated to the boiling point. The substance was then added, the mixture well shaken, kept at the boiling point for precisely 15 minutes, filtered immediately, and washed with hot water until the washings no longer reddened blue litmus paper. The solid matter was then washed from the filter by means of hot water, and more hot water added to make up to the same volume as before, namely, 1200 septems. The mixture being brought to the boiling point, in the flask as before, there was then added exactly sufficient of a concentrated solution of caustic potash of known strength to bring the whole to 1 per cent. alkali. This was then boiled for precisely 15 minutes, the supernatant fluid filtered rapidly as possible (the filtration being much expedited by pouring boiling water into the filter along with the fluid), and finally the solid matter transferred to the filter, and washed as rapidly as possible with boiling water. The washing was continued until the filtrate no longer turned reddened litmus paper blue; then two or three drops of dilute sulphuric acid were added, and the washing continued with boiling water until the washings no longer turned blue litmus paper red. The residue on the filter was then transferred to a small covered crucible, dried at 212°, and weighed, the result representing crude woody-fibre. Of this, a weighed portion was burnt, to determine the amount of mineral matter; another portion was taken for the determination of nitrogen, which, multiplied by 6·3 gave the amount of nitrogenous substance. The amount of mineral and nitrogenous matters so determined were then deducted, by calculation, from the crude woody-fibre, the remainder representing the woody-fibre as entered in Table II., p. 60.

In the Condition of Dryness as analysed.

Dates of sampling.	Number of Days growing.	Description.	Per Cent. Dry Substance in the Condition of Dryness as analysed.	By Fermentation & Alcohol Method.				By Copper Method.				Mean, by the Two Methods.					
				By Fermentation & Alcohol Method.		By Copper Method.		By Fermentation & Alcohol Method.		By Copper Method.		By Fermentation & Alcohol Method.		By Copper Method.			
				Expt. 1.	Expt. 2.	Expt. 3.	Mean.	Expt. 1.	Expt. 2.	Expt. 3.	Mean.	Expt. 1.	Expt. 2.	Expt. 3.	Mean.	In the Condition of Dryness as analysed.	In the Condition of Dryness as analysed.
Barley "No. 1."																	
Oct. 30	—	Barley	92.1	2.53	2.42	2.34	2.43	2.32	2.23	2.26	2.27	2.11	2.55	2.11	2.11	2.11	2.11
Nov. 2	6	As thrown from the couch	93.9	1.56	1.56	—	1.56	1.37	—	1.37	—	0.99	1.56	0.99	1.56	1.46	1.46
" 10	8	On the floor	93.6	7.59	7.78	—	7.68	7.58	—	7.58	—	4.74	8.16	4.74	8.16	7.61	7.61
" 12	10	Do.	92.7	9.35	9.54	—	9.44	9.46	—	9.46	—	5.93	10.19	5.93	10.19	9.51	9.51
" 14	12	Do. (1st third to kiln)	Not sampled.	11.23	—	—	—	10.94	—	10.94	—	6.92	11.67	6.92	11.67	10.90	10.90
" 16	14	Do. (2nd third to kiln)	94.5	11.01	11.01	—	11.12	10.94	—	10.94	—	7.23	12.14	7.23	12.14	11.33	11.33
" 18	16	Do. (3rd third to kiln)	91.6	10.89	10.70	10.27	10.45	10.20	10.66	10.35	10.40	10.62	11.38	10.62	11.38	10.62	10.62
" 20	18	Screened malt	91.4	10.38	10.43	10.17	10.43	13.12	13.02	13.08	13.08	10.01*	11.35*	10.01*	11.35*	10.01*	10.01*
" 22	20	Malt-dust	91.9	10.64	—	—	—	—	—	—	—	—	—	—	—	—	—

Barley "No. 2."

Nov. 3	—	Barley	93.8	3.11	3.20	3.03	3.11	3.64	3.17	3.43	3.27	2.83	3.49	2.83	3.49
" 6	6	As thrown from the couch	93.8	2.03	2.13	—	2.08	1.83	—	1.83	—	1.14	2.08	1.14	2.08
" 10	10	On the floor	93.6	6.22	6.51	—	6.36	6.41	—	6.41	—	3.73	6.80	3.73	6.80
" 12	12	Do.	93.6	10.27	10.38	—	10.32	10.10	—	10.10	—	6.13	10.91	6.13	10.91
" 14	14	Do. (1st third to kiln)	94.4	9.88	9.74	—	9.81	10.28	—	10.22	—	8.95	10.60	8.95	10.60
" 16	16	Do. (2nd third to kiln)	92.1	9.88	9.54	—	9.71	10.28	—	10.17	—	9.94	10.79	9.94	10.79
" 18	18	Do. (3rd third to kiln)	93.5	10.70	10.48	—	10.59	10.95	—	10.95	—	10.86	11.52	10.86	11.52
" 20	20	Screened malt	91.4	11.01	10.70	11.11	10.94	10.59	11.01	10.80	10.87	11.34	11.89	11.34	11.89
" 22	22	Malt-dust	91.4	11.44	11.44	11.76	11.54	13.56	13.81	13.64	13.64	10.89*	12.62*	10.89*	12.62*

* In the cases of the malt-dust, the results by the fermentation and alcohol method are alone adopted; those by the copper method being supposed to be too high owing to reduction by other matters than sugar in the malt-dust extract. The probability of this was confirmed by determinations made by extraction of the malt-dust with spirit, expansion of the alcohol by evaporation, and submitting the residue to fermentation in the ordinary way, when the results obtained were much below those by the copper method, and corresponded fairly with those obtained by the fermentation of the usual aqueous extract.

TABLE II.—SHOWING the DETERMINATIONS of WOODY-FIBRE, in the BARLEY, and in the PRODUCTS of the MALTING PROCESS.

Particulars of the Sampling.			Per Cent. Woody Fibre.				
Dates of Sampling.	Number of Days growing.	Description.	Per Cent. Dry Matter in Condition as sampled.	In the Dry Substance.			In the Barley, Malt, and Dust, as sampled; in others in the same Dryness as Malt.
				Expt. 1.	Expt. 2.	Mean.	
Barley "No. 1."							
Oct. 30	—	Barley	82.4	4.75	4.62	4.69	3.86
Nov. 2	44	As thrown from the couch	57.3	5.22	5.23	5.23	2.99
" 6	8	On the floor	58.1	4.97	4.95	4.96	2.88
" 10	8	Ditto	58.4	5.22	5.14	5.18	3.02
" 12	104	Ditto (1st third to kiln)	Not sampled.	—	—	—	—
" 14	124	Ditto (2nd third to kiln)	59.3	4.98	5.07	4.98	4.55
" 16	144	Ditto (3rd third to kiln)	59.6	5.00	5.06	5.03	4.69
" 16		Screened Malt	93.3	4.89	4.78	4.84	4.52
" 16		Malt-dust	88.3	9.61	9.73	9.67	8.54
Barley "No. 2."							
Nov. 3		Barley	81.0	5.55	5.47	5.51	4.47
" 6		As thrown from the couch	54.7	Lost	5.61	5.61	5.34
" 10	4	On the floor	54.9	5.25	5.20	5.22	2.87
" 14	8	Ditto	56.2	5.61	5.60	5.60	3.15
" 16	11	Ditto (1st third to kiln)	59.1	5.52	5.40	5.46	3.32
" 18	13	Ditto (2nd third to kiln)	59.3	5.55	5.43	5.49	3.47
" 21	16	Ditto (3rd third to kiln)	59.6	5.12	5.78	5.45	5.73
" 21		Screened Malt	95.3	6.73	6.78	6.75	5.49
" 21		Malt-dust	88.3	9.54	9.60	9.55	8.41

Particulars of the Sampling.			Per Cent. Nitrogen.					Per Cent. Nitrogenous Substance (N : X * 8.3).		
Dates of Sampling.	Number of Days growing.	Description.	Per Cent. Dry Matter, in the Condition of Moisture as sampled.	In the Dry Substance.				In the Condition of Moisture as sampled.	In the Barley Malt, and Dust, as sampled; in others in the same Dryness as Malt.	
				Expt. 1.	Expt. 2	Expt. 3.	Expt. 4.			Mean.
Barley "No. 1."										
Oct. 30		Barley	82.4	1.59	1.54	1.56	1.54	1.56	8.09	
Nov. 2		As thrown from the couch	57.3	1.58	1.54	1.56	1.56	5.63	9.17	
" 6	4½	On the floor	58.1	1.51	1.59	1.59	1.60	5.75	9.23	
" 10	8	Ditto	58.4	1.53	1.61	1.62	1.66	5.92	9.47	
" 12	10½	Ditto (1st third to kiln)	59.3	—	—	—	—	—	—	
" 14	12½	Ditto (2nd third to kiln)	59.3	1.53	1.69	1.63	1.68	6.09	9.59	
" 16	14½	Ditto (3rd third to kiln)	59.6	1.67	1.66	1.57	1.63	6.19	9.70	
" 18		Screened Malt	93.3	1.61	1.58	1.59	1.54	9.29	9.29	
" 20		Malt-dust	88.3	4.23	4.18	4.30	4.38	23.75	23.75	
Barley "No. 2."										
Nov. 3		Barley	81.0	1.83	1.92	1.96	1.96	1.94	9.91	
" 6		As thrown from the couch	64.7	1.92	1.90	1.92	1.93	6.61	12.22	
" 10	4	On the floor	64.9	1.96	1.96	1.96	1.99	6.71	12.10	
" 14	8	Ditto	66.2	1.99	1.97	1.95	1.94	6.94	12.35	
" 17	11	Ditto (1st third to kiln)	66.1	1.88	1.99	2.05	1.96	6.98	12.44	
" 19	13	Ditto (2nd third to kiln)	68.3	1.96	2.06	1.97	1.99	7.31	12.54	
" 21	15	Ditto (3rd third to kiln)	69.6	2.04	2.03	2.07	1.98	7.62	12.19	
" 21		Screened Malt	95.3	1.95	1.86	1.87	1.89	11.35	11.35	
" 21		Malt-dust	86.2	4.07	4.13	4.03	4.00	22.06	22.06	

TABLE IV.—DETAILS of the SAMPLING, and of the DETERMINATIONS of the DRY SUBSTANCE and MINERAL MATTER (ASH) in the BARLEY, and in the PRODUCTS of the MALTING PROCESS.

Barley "No. 1."

Particulars of Sampling.			Actual Weights.				Per-centages.			
Dates of Sampling.	Number of Days growing.	Description.	As sampled.	Taken for drying and burning.	Dried at 212° F.	Burnt (Ash).	Dry in Fresh.	Ash in Fresh.	Ash in Dry.	
			lbs.	ozs.	ozs.	ozs.	Each Experiment.	Mean.	Each Experiment.	
Oct. 30	—	Barley	25 ¹	99.96 99.96	82.31 82.34	2.038 2.076	82.34 82.37	2.030 2.077	2.476 2.521	
Nov. 2	—	As thrown from the couch	25	100.00 100.00	57.23 57.36	1.306 1.290	57.23 57.36	1.306 1.290	2.282 2.249	
„ 6	4½	On the floor	25 ²	99.98 99.98	58.05 58.18	1.315 1.327	58.06 58.19	1.315 1.327	2.265 2.281	
„ 10	8	Do.	25 ³	99.97 99.97	58.29 58.38	1.365 1.352	58.31 58.39	1.365 1.352	2.342 2.316	
„ 12	10½	Do. (1st third to kiln)	(Not sampled).	99.97 99.97	59.24 59.29	1.392 1.392	59.26 59.31	1.392 1.392	2.350 2.349	
„ 14	12½	Do. (2nd third to kiln)	25 ⁴	99.96 99.96	59.56 59.51	1.396 1.402	59.58 59.53	1.397 1.403	2.344 2.350	
„ 16	14½	Do. (3rd third to kiln)	25 ⁵	99.92 99.92	92.89 93.63	2.015 2.032	92.97 93.71	2.017 2.034	2.169 2.170	
		Screened Malt	25 ⁶	49.93 49.93	44.04 44.11	3.690 3.716	88.20 88.33	7.390 7.442	8.379 8.425	
		Malt-dust	12.57							

30.130 tons of malted malt.

5.0 tons ash of malted malt.

1.0 tons ash of malted malt.

**ANALYSES, MADE BY THE GOVERNMENT,
in the Haver, and in the Provinces of the
Barley No. 2.**

Particulars of Sampling.			Actual Weights.				Per-centages.					
Dates of Sampling.	Number of Days growing.	Description.	As sampled.	Taken for drying at burning.	Dried at 212° F.	Burnt (Ash).	Dry in Fresh.		Ash in Fresh.		Ash in Dry.	
							Each Experiment.	Mean.	Each Experiment.	Mean.	Each Experiment.	Mean.
Nov. 3	—	Barley	25 ¹	99.46 80.53	80.53 2.052	80.37	81.04	2.063	2.083	2.548	2.571	
„ 6	—	As thrown from the couch . . .	25 ²	99.72 99.72	54.58 1.420	54.73	54.68	1.424	1.415	2.602	2.588	
„ 10	4	On the floor	25 ³	99.79 99.79	54.78 1.444	54.90	54.86	1.447	1.398	2.636	2.547	
„ 14	8	Do.	25 ⁴	99.81 99.81	55.96 1.537	56.07	56.19	1.540	1.539	2.747	2.739	
„ 17	11	Do. (1st third to kiln) . . .	25 ⁵	99.79 99.79	55.83 1.580	55.95	56.11	1.583	1.601	2.890	2.854	
„ 19	13	Do. (2nd third to kiln) . . .	25 ⁶	99.78 99.78	58.06 1.675	58.18	58.26	1.679	1.656	2.885	2.841	
„ 21	15	Do. (3rd third to kiln) . . .	25 ⁷	99.74 99.74	59.35 1.653	59.51	59.59	1.657	1.676	2.785	2.813	
		Screened Malt	25 ⁸	99.57 99.57	94.81 2.325	95.22	95.33	2.335	2.366	2.452	2.482	
		Malt-dust	12.5 ⁹	49.69 49.69	42.88 3.830	86.31	86.23	7.708	7.821	8.931	9.070	

1 2.158 ozs. of stones, &c., picked out.	3 0.415 ozs. of stones, &c., picked out.
4 0.771 "	6 0.469 "
7 1.043 "	9 1.266 "

TABLE VI.—SHOWING the ACTUAL WEIGHTS of the BARLEY, MALT (and intermediate Products), and also the QUANTITIES of each, and some of their CONSTITUENTS, as corrected by Calculation for the Amounts removed as Samples during the MALTING PROCESS.

Particulars of the Sampling.			Actual Weights.			Quantities as corrected for Samples taken.					
Dates of sampling. (1863.)	Number of Days growing.	Description.	Number of Bushels.	Weight per Bushel.	Total Weight.	Taken for Samples.	In Condition of Moisture. as sampled.	Total Dry Substance.	Mineral Matter (Ash).	Dry Organic Matter.	Nitrogen.
Barley "No. 1."											
Oct. 30		Barley	280	lbs. oza.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Nov. 2		As thrown from the couch	401	54 10	15,295	—	15,295	12,597	314½	12,282½	196.5
" 6	4½	On the floor	508	54 9-74	21,898	25	21,898	12,547½	284½	12,263½	195.7
" 10	8	Do.	547	42 3	21,434	25	21,458½	12,473½	283½	12,190½	195.8
" 12	10½	Do. (1st third to kiln)	—	37 12	20,657½	25	20,705	12,081½	281½	11,799½	194.5
" 14	12½	Do. (2nd third to kiln)	—	—	—	25	—	—	—	—	—
" 16	14½	Do. (3rd third to kiln)	—	—	—	25	—	—	—	—	—
		Screened Malt	290	41 10	12,002½	—	12,072½	11,268½	244½	11,024	178.0
		Malt-dust	—	—	334	—	336	296½	25	271½	12.7
							12,406½	11,565	269½	11,295½	190.7
Barley "No. 2."											
Nov. 3		Barley	280	50 10	14,175	—	14,175	11,487½	295½	11,192½	222.9
" 6		As thrown from the couch	397	52 9-1	20,869½	25	20,869½	11,411½	295½	11,116½	219.1
" 10	4	On the floor	—	—	—	25	—	—	—	—	—
" 14	8	Do.	594½	32 15-4	19,606	25	19,633½	11,043½	302½	10,741	216.5
" 17	11	Do. (1st third to kiln)	—	—	—	25	—	—	—	—	—
" 19	13	Do. (2nd third to kiln)	—	—	—	25	—	—	—	—	—
" 21	15	Do. (3rd third to kiln)	—	—	—	25	—	—	—	—	—
		Screened Malt	284	37 4	10,499	—	10,578	10,084	250½	9,833½	190.6
		Malt-dust	—	—	451	—	454½	392½	35½	356½	15.9
							11,032½	10,476	285½	10,190½	206.5

TABLE VII.—RESULTS of the partial ANALYSIS of WATER before and after being used for “STEEPING” BARLEY.

	Constituents.	Grains per Gallon.	
--	---------------	--------------------	--

Pump Water before being used for Steeping.

	Organic matter	2·79	
	Mineral matter	26·80	
	Total solid matter . . .	29·09	

Water from the Cistern after Steeping.

	Organic matter	265·05 ²	
	Mineral matter	155·70	
	Total solid matter . . .	420·75	

² Containing Nitrogen 8·62 = Nitrogenous substance 54·31.

TABLE VIII.—RESULTS of the PARTIAL ANALYSIS of the BARLEY, MALT, and MALT-DUST (and intermediate Products) used in EXPERIMENTS with SHEEP, at ROTHAMSTED, in 1849.

	Barley before steeping.	Growing.					Screened Malt.	Malt and Kiln-dust.
		Feb. 18; as thrown from the Couch.	Feb. 22; 4 Days on the Floor.	Feb. 26; 8 Days on the Floor.	March 3; 13 Days on the Floor.	Mar. 6; 16 Days on the Floor.		
1.—In the Condition of Moisture as sampled.								
Starch, sugar, woody-fibre, &c.	70·37						82·57	59·23
Albuminous (or "flesh-forming") matters'	9·14						10·21	25·83
Mineral matter (ash)	2·33						2·61	8·70
Total solid matter	81·84	57·74	57·86	58·22	58·83	59·76	96·39	98·76
Moisture	18·16	42·26	42·14	41·78	41·17	40·24	4·61	6·24
Total	100·00	100·00	100·00	100·00	100·00	100·00	100·00	100·00
(1) Containing nitrogen	1·45						1·62	4·10
2.—Exclusive of Moisture.								
Starch, sugar, woody-fibre, &c.	85·95						86·56	63·12
Albuminous (or "flesh-forming") matters'	11·21						10·71	27·59
Mineral matter (ash).	2·84						2·73	9·29
Total solid matter	100·00						100·00	100·00
(2) Containing nitrogen.	1·78						1·70	4·38

	Barley before Steeping.	Feb. 22; 4 Days on the Floor.	Feb. 26; 8 Days on the Floor.	Mar. 3; 13 Days on the Floor.	Final Products.			Loss.
					Screened Malt.	Malt and Klin- dust.	TOTAL.	

Actual Weights, Lbs.

As sampled	7,632	10,504	10,415	10,010	5,907½	266	6,173½	1,458½
Total dry or solid matter	6,246	6,078	6,063	5,889	5,635½	249½	5,885	361
Non-nitrogenous organic matter	5,371				4,878½	157½	5,036½	334½
Nitrogenous matter	697.2				602.9	68.7	671.6	25.6
Mineral matter	177.5				153.9	123.1	1,177.0	10.5

Proportion to 100 before steeping.

As sampled	100	137.6	136.4	131.2	77.41	3.48	80.89	19.11
Total dry or solid matter	100	97.3	97.1	94.3	90.23	3.99	94.22	5.78
Non-nitrogenous organic matter	100				90.83	2.94	93.77	6.23
Nitrogenous matter	100				86.47	9.85	96.32	3.68
Mineral matter	100				86.70	13.01	199.71	10.29

¹ The ash from the klin-dust included some incombustible matter from the furnace, and hence the amounts of mineral matter set down for the malt and klin-dust and for the total products, are somewhat too high, and the recorded loss of mineral matter is in a corresponding degree too low.

TABLE X.—EXPERIMENTS made with COWS, at New
1st Week.—Decem

—	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Dec. 21).	MONDAY.		Tue
						A.M.	P.M.	A.M.

Lot 1.—Special Food.—Unmalted

Yield of Milk, &c.	Nos.					lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May 20		1,086	15 4	9 12	13 7
	2	Cross Shorthorn .	6	Oct. 26		1,044	19 2	14 13	18 14
	3	Cross Shorthorn .	Aged	April 26		1,065	17 0	8 2	16 3
	4	Cross Shorthorn .	7	April 21		1,082	13 10	8 13	13 3
	5	Cross Shorthorn .	8	Sept. 26		1,020	15 7	13 4	16 0
	6	Cross Yorkshire .	7	April 14		1,164	14 9	8 3	14 6
	7	Cross Yorkshire .	10	April 9		1,260	14 2	8 7	14 0
	8	Cross Ayrshire .	5	June 15		944	16 8	9 6	16 9
	9	Shorthorn . .	4	June 15		1,156	14 10	7 7	13 7
	10	Shorthorn . .	7	Oct. 29		1,236	25 12	14 3	23 7
				Totals .		11,057	166 0	102 6	159 8
				Averages		1,106	16 10	10 4	15 15

Lot 2.—Special Food.—Malt and M

Yield of Milk, &c.	Nos.					lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Yorkshire .	8	March 10		1,184	12 0	7 8	12 0
	2	Cross Shorthorn .	7	Oct. 27		992	18 12	10 12	18 9
	3	Shorthorn . .	8	April 29		1,260	13 6	8 4	11 7
	4	Cross Shorthorn .	7	June 19		1,020	14 12	8 12	11 15
	5	Cross Shorthorn .	6	Oct. 29		914	15 8	10 4	13 10
	6	Cross Welch . .	8	June 18		1,316	16 10	8 12	14 12
	7	Cross Shorthorn .	7	June 24		1,008	16 12	9 8	16 7
	8	Cross Yorkshire .	10	April 26		1,065	16 8	10 11	14 15
	9	Cross Shorthorn .	6	April 19		1,040	18 14	10 8	18 2
	10	Shorthorn . .	9	Nov. 20		1,206	25 4	11 13	21 14
				Totals .		11,005	168 6	96 12	183 11
				Averages		1,101	16 13	9 11	15 6

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff

UGBY. DETAILED RECORD of the MILK YIELDED, &c.
er 27, 1863.

	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

lbs. per head per day.*

oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.
4	14 3	8 13	12 14	9 6	12 15	10 0	12 4	8 13	158 12	22 11	
9	17 7	15 4	17 2	14 1	17 5	15 12	17 11	13 11	229 2	32 12	
8	16 6	9 14	16 0	9 2	17 1	9 1	16 3	9 4	180 7	25 12	
6	14 0	10 3	14 7	10 1	14 8	8 6	14 13	10 8	165 14	23 11	
4	13 7	11 15	14 8	11 3	16 12	11 7	16 0	11 2	187 4	26 12	
10	13 11	8 0	14 0	8 4	14 11	8 5	14 6	7 13	157 6	22 8	
11	13 7	9 6	12 14	8 1	14 4	8 5	13 14	8 13	156 4	22 5	
12	16 1	9 6	17 1	8 9	16 12	9 3	16 10	8 12	177 13	25 6	
0	13 6	8 11	13 5	7 8	14 3	7 1	14 4	7 5	150 12	21 9	
10	24 7	14 4	24 6	12 6	25 8	13 4	24 2	12 6	265 12	37 15	
10	156 7	105 12	156 9	98 9	163 15	100 12	160 3	98 7	1,829 6	—	
4	15 10	10 9	15 11	9 14	16 6	10 1	16 0	9 14	182 15	26 2	

a. Barley (No. 1.), per head per day.*

a. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.
7 0	11 10	6 15	11 15	5 6	11 4	6 9	10 0	3 7	125 2	17 14	
4	18 14	11 1	17 15	9 2	18 3	11 2	18 8	9 4	202 8	28 15	
10	12 7	7 3	11 12	6 15	11 15	7 0	11 6	6 6	137 4	19 10	
12	12 5	10 6	14 9	7 6	14 14	7 15	14 0	8 8	157 15	22 9	
7	14 9	9 12	15 5	9 7	15 3	9 11	15 14	9 8	173 13	24 13	
14	15 5	9 4	14 15	8 0	16 1	8 6	15 12	7 10	169 7	24 3	
12	16 14	10 8	17 5	9 1	17 8	10 10	18 0	8 6	187 1	26 12	
14	16 7	9 11	17 0	8 7	16 8	9 9	16 5	9 0	180 5	25 12	
10	18 8	10 13	18 12	8 5	19 0	11 0	19 7	10 2	202 14	29 0	
0	24 15	13 14	23 6	12 4	24 15	13 4	25 0	12 1	260 15	37 4	
3	161 14	99 7	162 14	84 5	165 7	95 2	164 4	84 4	1,797 4	—	
10	16 3	9 15	16 5	8 7	16 9	9 8	16 7	8 7	179 12	25 11	

7 lbs.; and pulped swedes, 50·4 lbs., per head per day.

TABLE XI.—EXPERIMENTS made with COWS, at NEWLANDS
2nd Week.—December 28, 1863.

—	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Dec. 21.)	MONDAY.		TUESDAY.	
						A.M.	P.M.	A.M.	P.M.

Lot 1.—Special Food.—Unmalted Barley

Yield of Milk, &c.	Nos.					lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May	20	1,086	11 8	8 10	12 10	8 1
	2	Cross Shorthorn .	6	Oct.	26	1,044	18 16	14 13	19 5	14 2
	3	Cross Shorthorn .	Aged	April	26	1,065	14 14	8 8	16 0	8 13
	4	Cross Shorthorn .	7	April	21	1,082	13 5	8 14	13 8	8 9
	5	Cross Shorthorn .	8	Sept.	26	1,020	16 2	11 6	17 0	10 9
	6	Cross Yorkshire .	7	April	14	1,164	13 9	8 3	14 2	8 4
	7	Cross Yorkshire .	10	April	9	1,260	13 0	7 4	13 10	8 9
	8	Cross Ayrshire .	5	June	15	944	15 8	8 14	16 1	8 10
	9	Shorthorn . .	4	June	15	1,156	13 1	6 13	13 5	7 3
	10	Shorthorn . .	7	Oct.	29	1,236	23 0	13 1	23 4	13 6
				Totals .		11,067	152 15	96 6	158 13	85 4
				Averages		1,106	15 5	9 10	15 14	9 4

Lot 2.—Special Food.—Malt and Malt-dust

Yield of Milk, &c.	Nos.					lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Yorkshire .	8	March	10	1,184	9 0	4 9	9 0	4 13
	2	Cross Shorthorn .	7	Oct.	27	992	17 2	10 14	19 2	10 0
	3	Shorthorn . .	8	April	29	1,260	10 7	6 15	10 2	6 10
	4	Cross Shorthorn .	7	June	19	1,020	13 1	9 0	13 12	8 8
	5	Cross Shorthorn .	6	Oct.	29	914	15 6	8 2	14 15	9 11
	6	Cross Welsh . .	8	June	18	1,316	15 13	8 5	15 7	8 11
	7	Cross Shorthorn .	7	June	24	1,008	16 10	9 12	17 12	9 14
	8	Cross Yorkshire .	10	April	26	1,065	15 4	7 9	16 0	8 9
	9	Cross Shorthorn .	6	April	19	1,040	17 12	11 4	18 0	9 2
	10	Shorthorn . .	9	Nov.	20	1,206	23 10	12 8	24 3	13 4
				Totals .		11,005	154 1	88 14	158 5	89 8
				Averages		1,101	15 7	8 14	15 13	8 13

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.;

RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
 ry 3, 1864.

AY.	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head Per Day.
	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

3 lbs. per head per day.*

a. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.
3 7	10 9	8 8	10 6	8 0	10 3	7 0	11 8	6 2	134 8	19 3	
5 6	18 2	14 2	19 5	14 8	17 2	15 0	17 4	13 1	229 2	32 12	
3 15	16 6	9 3	16 7	8 14	16 0	8 11	16 4	7 9	172 12	24 11	
3 6	12 8	9 0	13 2	9 13	13 10	8 12	13 8	8 3	154 2	22 0	
1 6	16 13	10 6	16 11	10 0	15 1	11 0	14 9	10 0	186 13	26 11	
3 3	13 13	8 8	13 11	8 0	13 0	7 7	13 2	7 0	151 4	21 10	
3 5	13 12	8 9	13 10	8 7	13 14	7 12	12 9	7 1	150 4	21 7	
9 0	16 5	8 11	16 12	8 3	15 2	8 0	13 12	5 3	166 3	23 12	
7 1	13 1	7 5	12 7	7 7	13 4	6 3	13 12	6 12	140 12	20 2	
2 8	21 7	10 6	20 0	11 0	20 13	11 2	21 2	9 6	231 11	33 2	
7 7	152 12	94 10	152 7	94 4	148 1	90 15	147 6	80 5	1,717 7	—	
9 12	15 4	9 7	15 4	9 7	14 13	9 2	14 12	8 1	171 12	24 9	

ss. Barley (No. 1.), per head per day.*

a. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.	lbs. oza.
5 12	9 10	4 14	9 1	5 3	9 0	6 0	10 0	5 0	101 3	14 7	
0 12	18 6	9 9	18 4	9 5	17 0	7 2	17 0	8 8	190 3	27 3	
6 10	10 6	8 2	9 15	5 10	8 10	5 12	8 0	4 10	112 4	16 1	
8 8	14 8	9 10	13 11	9 10	14 13	7 0	14 0	7 6	159 2	22 12	
9 9	15 12	8 14	16 6	7 13	16 2	8 9	15 0	8 0	169 10	24 4	
8 11	16 0	8 12	15 5	9 0	14 0	8 2	14 11	6 14	167 7	23 15	
0 12	17 1	9 11	17 8	9 14	17 0	9 10	17 0	8 2	188 4	26 14	
8 9	15 0	8 7	14 12	8 9	14 13	8 12	15 12	6 8	164 10	23 8	
9 9	18 4	10 6	18 5	9 2	17 4	10 8	17 0	8 14	191 7	27 6	
13 15	24 15	12 7	23 12	12 11	22 1	13 4	24 0	10 5	255 12	36 9	
12 11	159 14	90 12	156 15	86 13	150 11	84 11	152 7	74 3	1,699 14	—	
9 4	16 0	9 1	15 11	8 11	15 1	8 8	15 4	7 7	170 0	24 5	

off, 7·7 lbs.; and pulped swedes, 50·4 lbs., per head per day.

TABLE XII.—EXPERIMENTS made with COWS, at NEWLANDS
3rd Week.—January 4

—	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Dec. 21.)	MONDAY.		TUESDAY.	
						A.M.	P.M.	A.M.	P.M.

Lot 1.—Special Food.—Unmalted Barley

Yield of Milk, &c.	Nos.					lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May 20		1,086	10 4	6 3	11 6	7 8
	2	Cross Shorthorn .	6	Oct. 26		1,044	19 0	14 5	17 8	14 1
	3	Cross Shorthorn .	Aged	April 26		1,065	16 0	9 1	14 5	8 15
	4	Cross Shorthorn .	7	April 21		1,082	13 11	8 3	12 14	9 7
	5	Cross Shorthorn .	8	Sept. 26		1,020	19 9	9 0	15 12	9 7
	6	Cross Yorkshire .	7	April 14		1,164	12 12	6 14	12 7	7 15
	7	Cross Yorkshire .	10	April 9		1,260	13 0	8 14	12 9	8 0
	8	Cross Ayrshire .	5	June 15		944	12 11	7 0	12 7	7 13
	9	Shorthorn .	4	June 15		1,156	12 0	6 12	11 9	6 9
	10	Shorthorn .	7	Oct. 29		1,236	20 10	12 0	23 4	12 6
				Totals .		11,057	149 9	88 4	144 1	91 8
				Averages		1,106	14 15	8 13	14 7	9 2

Lot 2.—Special Food.—Malt and Malt-dust

Yield of Milk, &c.	Nos.					lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Yorkshire .	8	March 10		1,184	9 4	5 15	9 1	6 4
	2	Cross Shorthorn .	7	Oct. 27		992	16 11	11 10	16 12	9 14
	3	Shorthorn .	8	April 29		1,260	8 7	4 14	6 4	5 2
	4	Cross Shorthorn .	7	June 19		1,020	10 11	8 8	12 6	7 14
	5	Cross Shorthorn .	6	Oct. 29		914	14 14	8 12	14 15	8 8
	6	Cross Welch .	8	June 18		1,316	14 10	8 8	14 8	8 8
	7	Cross Shorthorn .	7	June 24		1,008	16 7	8 14	16 4	9 6
	8	Cross Yorkshire .	10	April 26		1,065	14 1	8 8	12 8	8 7
	9	Cross Shorthorn .	6	April 19		1,040	16 8	11 8	16 4	9 12
	10	Shorthorn .	9	Nov. 20		1,206	22 0	12 14	23 8	12 12
				Totals .		11,005	143 9	89 15	142 6	86 9
				Averages		1,101	14 6	9 0	14 4	8 11

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.;

RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
May 10, 1864.

DAY.	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head Per Day.
P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

); 3 lbs. per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
7 6	9 1	7 4	10 0	7 1	8 8	7 9	9 0	8 6	119 4	17 1	
14 6	17 12	12 10	18 1	13 5	17 14	13 12	17 8	12 5	219 7	31 6	
8 9	14 14	8 7	15 6	8 14	16 0	8 1	15 4	8 12	167 8	23 15	
9 3	12 0	9 12	14 2	7 15	12 15	8 12	12 0	9 0	153 15	22 0	
11 1	16 2	9 0	15 0	10 8	14 7	10 0	15 2	9 11	179 12	25 1	
6 5	12 0	7 8	11 10	6 8	12 2	6 8	11 12	6 11	133 9	19 1	
7 10	12 13	8 5	13 11	8 0	13 3	7 14	12 0	7 3	146 3	20 14	
6 14	12 7	6 10	12 12	7 11	14 1	6 15	12 8	6 12	139 4	19 14	
6 14	11 6	6 10	11 11	6 7	12 0	6 12	12 0	6 5	127 13	18 4	
12 4	22 0	12 3	22 12	12 1	22 8	11 2	21 10	9 8	236 7	33 12	
90 8	140 7	88 5	145 1	88 6	143 10	87 5	138 12	84 9	1,623 2	—	
9 1	14 1	8 13	14 8	8 13	14 6	8 12	13 14	8 7	162 5	23 3	

3 lbs. Barley (No. 1.), per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
5 10	9 8	5 8	9 12	6 0	9 15	5 2	10 1	4 0	105 15	15 2	
9 5	17 0	9 13	16 1	10 4	17 11	9 2	17 6	8 3	188 12	26 15	
4 5	6 0	5 5	5 4	4 0	5 10	3 10	5 1	3 4	73 13	10 9	
8 3	11 1	7 9	12 14	7 1	13 15	7 2	14 0	6 7	139 4	19 14	
7 13	15 2	8 3	14 15	8 6	14 7	8 9	14 7	7 10	162 6	23 3	
7 3	14 5	8 12	15 6	8 0	15 14	8 0	15 3	7 8	161 7	23 1	
8 15	15 4	9 0	16 9	8 14	16 6	8 9	16 14	8 4	176 4	25 3	
7 3	12 0	7 7	13 14	7 0	13 15	6 8	15 8	7 5	148 1	21 2	
9 14	17 7	10 2	18 12	9 12	18 0	10 15	16 12	9 6	191 10	27 6	
12 3	21 8	12 14	23 0	12 0	22 8	10 10	23 7	11 4	242 14	34 11	
90 10	139 13	84 10	147 4	81 5	148 5	78 3	148 11	73 3	1,590 6	—	
8 1	14 0	8 7	14 12	8 2	14 13	7 13	14 14	7 5	159 1	22 12	

and 7-7 lbs.; and pulped swedes, 50-4 lbs., per head per day.

TABLE XIII.—EXPERIMENTS made with COWS, at NEWLANDS
4th Week.—January 11

	Cows	Breed	Years old	Dates of Calving	Weights of the Animals Dec. 11.	MONDAY.		TUESDAY.	
						A.M.	P.M.	A.M.	P.M.
Lot 1.—Special Food.—Unmalted Barley									
Yield of Milk, &c.	Nos.				Lbs.	Lbs. ozs.	Lbs. ozs.	Lbs. ozs.	Lbs. ozs.
	1	Cross Shorthorn .	7	May 20	1,066	10 2	7 12	9 12	8 8
	2	Cross Shorthorn .	6	Oct. 28	1,444	16 8	14 3	16 10	12 8
	3	Cross Shorthorn .	Agd.	April 28	1,065	13 10	9 6	16 1	8 8
	4	Cross Shorthorn .	7	April 21	1,082	11 12	9 2	12 13	8 10
	5	Cross Shorthorn .	8	Sept. 28	1,029	13 6	10 6	14 15	8 11
	6	Cross Yorkshire .	7	April 14	1,164	11 12	7 8	21 10	7 2
	7	Cross Yorkshire .	10	April 9	1,360	13 2	8 4	13 0	7 12
	8	Cross Ayrshire .	5	June 15	944	13 12	7 6	14 6	7 10
	9	Shorthorn . .	4	June 15	1,156	13 5	7 8	13 1	8 10
	10	Shorthorn . .	7	Oct. 29	1,236	21 8	11 8	22 0	11 8
Totals .					11,057	142 13	92 4	144 4	86 8
Averages					1,106	14 5	9 4	14 7	8 10
Lot 2.—Special Food.—Malt and Malt-dust									
Yield of Milk, &c.	Nos.				Lbs.	Lbs. ozs.	Lbs. ozs.	Lbs. ozs.	Lbs. ozs.
	1	Cross Yorkshire .	8	March 10	1,184	10 1	5 2	9 8	8 14
	2	Cross Shorthorn .	7	Oct. 27	992	18 2	9 3	17 12	9 12
	3	Shorthorn . .	8	April 29	1,260	5 13	4 0	5 11	3 8
	4	Cross Shorthorn .	7	June 19	1,020	13 8	8 10	12 6	10 1
	5	Cross Shorthorn .	6	Oct. 29	914	15 5	8 4	14 8	7 12
	6	Cross Welch . .	8	June 18	1,316	15 0	8 0	14 7	8 10
	7	Cross Shorthorn .	7	June 24	1,008	17 0	9 13	17 10	9 0
	8	Cross Yorkshire .	10	April 26	1,065	14 10	9 4	15 4	8 12
	9	Cross Shorthorn .	6	April 19	1,040	17 10	10 0	17 12	10 8
	10	Shorthorn . .	9	Nov. 20	1,206	23 0	13 1	22 8	12 4
Totals .					11,005	150 1	85 5	147 0	83 10
Averages					1,101	15 0	8 9	14 11	8 9

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.;

, RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
uary 17, 1864.

WEDAY.	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

1; 3 lbs. per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
7 14	9 10	7 8	10 9	7 6	9 8	7 14	9 0	7 2	121 14	17 7	
13 14	17 8	13 6	17 8	13 4	17 6	14 12	17 4	13 4	213 15	30 9	
9 2	16 0	8 14	16 8	8 5	15 14	9 7	16 0	8 11	173 6	24 12	
9 0	13 6	8 6	13 0	8 8	12 6	10 3	12 4	9 2	150 14	21 9	
10 10	13 0	10 15	15 2	9 12	15 9	9 14	14 6	9 7	170 14	24 7	
6 10	13 15	7 8	13 6	7 2	12 10	7 15	13 12	7 7	139 15	20 0	
7 13	13 8	7 14	13 12	7 5	13 1	7 15	14 6	6 13	147 9	21 1	
7 8	14 10	8 1	15 1	8 1	13 7	7 15	14 10	7 0	154 4	22 1	
7 0	12 9	6 14	13 5	6 9	12 13	7 3	12 6	5 12	137 13	19 11	
11 10	21 10	11 10	23 0	11 14	21 14	11 6	22 6	12 6	237 0	33 14	
5 00 1	145 12	91 0	151 3	88 2	144 8	94 8	146 6	87 0	1,647 8	—	
5 9 0	14 9	9 2	15 2	8 13	14 7	9 7	14 10	8 11	164 12	23 9	

3 lbs. Barley (No. 1.), per head per day.*

ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
6 5 4	10 0	3 12	10 9	5 10	9 13	5 10	10 3	4 9	105 15	15 2	
12 9 5	17 4	9 10	17 8	9 3	17 4	9 14	18 4	8 4	189 7	27 1	
6 4 0	3 6	3 2	4 6	3 0	4 4	2 10	2 9	3 0	53 0	7 2	
1 7 0	14 5	7 2	12 6	8 14	11 12	8 1	12 0	7 10	145 6	20 12	
6 7 12	15 0	9 10	13 15	7 12	12 4	7 4	12 13	6 6	151 0	21 2	
13 8 4	14 6	9 11	14 7	8 4	14 6	8 1	16 0	7 14	161 3	23 0	
0 9 3	17 0	10 0	17 15	9 12	18 8	10 2	18 8	10 4	192 11	27 8	
4 7 14	14 2	7 14	15 4	8 4	14 10	8 0	14 7	8 0	161 9	23 1	
0 9 9	18 0	10 4	17 12	10 9	18 1	18 1	10 3	20 0	197 5	28 3	
6 12 9	21 11	11 7	21 0	11 5	20 8	12 1	22 0	12 6	238 2	34 0	
11 00 12	142 9	82 8	145 2	82 14	141 6	81 14	146 9	78 5	1,595 10	—	
9 8 1	14 4	8 4	14 8	8 5	14 2	8 3	14 11	7 13	159 9	22 13	

and 7-7 lbs. ; and pulped swedes, 50-4 lbs., per head per day.

TABLE XIV.—EXPERIMENTS made with COWS, at NEWLANDS
5th Week.—January 18

	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Jan. 18).	MONDAY.		TUESDAY.	
						A.M.	P.M.	A.M.	P.M.
Lot 1.—Special Food.—Unmalted Barley.									
Yield of Milk, &c.	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May 20	1,134	11 0	7 12	8 14	7 12
	2	Cross Shorthorn .	6	Oct. 25	1,112	17 8	13 13	16 19	14 15
	3	Cross Shorthorn .	Aged	April 26	1,127	15 6	8 9	16 6	8 2
	4	Cross Shorthorn .	7	April 21	1,125	11 4	8 7	12 12	8 8
	5	Cross Shorthorn .	8	Sept. 26	994	14 2	9 7	15 0	10 2
	6	Cross Yorkshire .	7	April 14	1,242	12 14	7 8	13 10	7 11
	7	Cross Yorkshire .	10	April 9	1,284	12 14	7 10	13 3	8 3
	8	Cross Ayrshire .	5	June 15	954	14 3	7 11	14 13	7 19
	9	Shorthorn . .	4	June 15	1,198	12 11	6 0	12 14	6 8
10	Shorthorn . .	7	Oct. 29	1,312	22 3	11 0	22 8	11 3	
				Totals .	11,482	147 1	87 13	146 8	90 13
				Averages	1,148	14 11	8 13	14 10	9 1

Lot 2.—Special Food.—Malt and Malt-dust

	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
Yield of Milk, &c.	1	Cross Yorkshire .	8	March 10	1,216	10 4	5 10	9 11	5 11
	2	Cross Shorthorn .	7	Oct. 27	1,024	17 13	8 6	17 13	9 4
	3	Shorthorn . .	8	April 29	1,336	3 7	2 8	3 11	3 6
	4	Cross Shorthorn .	7	June 19	1,047	13 5	9 14	13 0	7 19
	5	Cross Shorthorn .	6	Oct. 29	942	11 6	7 5	11 8	9 2
	6	Cross Welch' . .	8	June 18	1,400	14 6	8 1	14 8	7 19
	7	Cross Shorthorn .	7	June 24	1,076	17 2	9 0	18 0	10 6
	8	Cross Yorkshire .	10	April 26	1,132	14 2	7 8	14 3	6 19
	9	Cross Shorthorn .	6	April 19	1,098	17 2	10 4	17 3	11 2
	10	Shorthorn . .	9	Nov. 20	1,237	22 2	11 1	21 0	11 11
				Totals .	11,508	141 1	79 9	140 9	82 14
				Averages	1,151	14 2	7 15	14 1	8 2

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.

RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
ary 24, 1864.

DAY.	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

); 3 lbs. per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
7 3	8 12	8 3	10 6	7 2	11 4	7 3	11 0	7 1	123 8	17 10	
13 11	17 4	13 10	18 1	13 0	18 5	13 8	17 10	13 0	219 0	31 5	
8 13	16 6	9 13	16 10	9 5	16 4	9 3	17 2	8 14	177 6	25 5	
9 0	13 3	8 12	13 6	8 3	13 10	8 8	13 8	7 14	153 5	21 14	
8 0	16 2	9 5	15 0	10 3	13 6	10 10	15 4	9 10	170 10	24 6	
7 8	12 14	7 13	13 6	7 11	13 11	7 4	13 0	6 13	145 3	20 12	
7 12	12 14	8 11	11 5	8 12	13 10	7 3	13 0	7 11	146 1	20 14	
7 8	14 4	7 12	14 12	7 14	14 4	8 1	14 12	7 0	155 2	22 3	
6 8	11 6	7 6	11 4	6 12	11 1	6 11	11 10	5 12	128 3	18 5	
12 8	21 2	12 8	21 10	13 4	21 10	12 5	21 0	10 14	235 7	33 10	
88 7	144 3	93 3	145 12	92 2	147 1	90 8	147 14	84 9	1,653 13	—	
8 14	14 7	9 5	14 9	9 3	14 11	9 1	14 13	8 7	165 6	23 10	

3 lbs. Barley (No. 1.), per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
5 10	9 10	5 10	10 6	5 13	9 4	5 9	9 0	5 9	108 0	15 7	
9 14	17 9	9 6	17 6	9 13	16 9	10 6	17 6	9 2	188 7	26 15	
3 0	2 4	2 3	2 2	2 0	2 3	2 0	2 0	1 6	35 6	5 1	
7 5	14 12	7 2	13 11	8 14	12 6	8 1	14 4	6 8	150 0	21 7	
8 4	13 2	8 1	13 12	7 12	14 1	7 10	13 2	8 0	147 2	21 0	
7 4	14 14	8 4	14 1	7 13	13 14	8 4	13 14	8 10	157 3	22 7	
9 5	16 11	10 5	16 4	7 12	15 10	8 3	16 2	9 0	180 12	25 13	
7 6	14 0	8 0	13 11	9 1	13 10	8 14	12 13	6 10	151 6	21 10	
11 4	18 1	11 9	17 11	11 3	18 6	9 7	18 12	10 15	201 11	28 13	
11 0	19 0	11 0	19 14	10 10	18 1	10 7	19 8	11 0	216 10	30 15	
80 4	139 15	81 8	138 14	80 11	134 0	78 13	136 13	76 12	1,536 9	—	
8 0	14 0	8 2	13 14	8 1	13 6	7 14	13 11	7 11	153 10	21 15	

aff, 7·7 lbs.; and pulped swedes, 50·4 lbs., per head per day.

TABLE XV.—EXPERIMENTS made with COWS, at NEWLANDS
6th Week.—January 25

—	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Jan. 15.)	MONDAY.		TUESDAY.	
						A.M.	P.M.	A.M.	P.M.
Lot 1.—Special Food.—Unmalted Barley									
Yield of Milk, &c.	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May 20	1,134	11 0	7 0	10 0	8 8
	2	Cross Shorthorn .	6	Oct. 26	1,112	18 12	12 6	17 8	13 4
	3	Cross Shorthorn .	Agd	April 26	1,127	16 8	9 14	16 3	9 4
	4	Cross Shorthorn .	7	April 21	1,125	13 1	8 10	14 2	8 10
	5	Cross Shorthorn .	8	Sept. 26	994	14 14	9 10	13 10	11 1
	6	Cross Yorkshire .	7	April 14	1,242	12 2	7 10	12 12	7 13
	7	Cross Yorkshire .	10	April 9	1,284	13 12	7 2	13 0	7 10
	8	Cross Ayrshire .	5	June 15	954	14 11	6 12	14 0	8 4
	9	Shorthorn . . .	4	June 15	1,198	11 8	6 3	11 8	7 0
10	Shorthorn . . .	7	Oct. 29	1,312	22 14	12 0	22 0	13 8	
				Totals .	11,482	149 3	87 3	145 5	94 1
				Averages	1,148	14 15	8 11	14 8	9 7

Lot 2.—Special Food.—Malt and Malt-dust

	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
Yield of Milk, &c.	1	Cross Yorkshire .	8	March 10	1,216	10 1	5 0	9 11	6 8
	2	Cross Shorthorn .	7	Oct. 27	1,024	17 6	8 12	17 0	10 0
	3	Shorthorn .	8	April 29	1,336	2 6	2 0	2 1	1 2
	4	Cross Shorthorn .	7	June 19	1,047	13 0	9 0	12 3	8 10
	5	Cross Shorthorn .	6	Oct. 29	942	14 0	7 12	14 0	7 10
	6	Cross Welch .	8	June 18	1,400	15 2	7 6	14 6	8 11
	7	Cross Shorthorn .	7	June 24	1,076	16 2	8 4	16 6	9 12
	8	Cross Yorkshire .	10	April 26	1,132	12 8	6 15	11 12	8 12
	9	Cross Shorthorn .	6	April 19	1,098	18 10	9 10	18 2	11 8
	10	Shorthorn .	9	Nov. 20	1,237	17 0	9 3	17 10	11 4
				Totals .	11,508	136 3	73 14	133 0	83 5
				Averages	1,151	13 10	7 6	13 5	8 8

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.;

RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
January 31, 1864.

WEDNESDAY.	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

); 3 lbs. per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
8 10	10 0	8 15	11 0	8 1	10 0	7 12	9 11	6 10	128 9	18 6	
12 10	17 3	13 4	17 13	13 13	18 6	12 8	18 0	9 0	212 13	30 6	
9 0	16 11	8 12	16 7	10 1	16 6	8 12	16 4	7 11	178 4	25 7	
9 6	14 0	8 4	12 13	9 7	13 7	8 13	13 0	7 3	153 14	22 0	
9 6	14 6	9 12	14 2	9 1	13 6	9 5	14 12	8 2	166 7	23 12	
7 2	15 5	7 4	11 8	8 0	12 11	7 4	12 0	6 2	140 5	20 1	
7 10	13 8	6 10	12 0	7 12	12 5	7 2	11 10	6 12	140 1	20 0	
8 0	14 15	7 12	14 12	8 6	15 11	7 0	14 2	6 11	156 9	22 6	
8 0	12 6	6 3	11 12	6 8	12 1	5 10	12 8	5 9	127 14	18 4	
12 4	21 0	11 2	21 2	12 0	20 0	10 8	19 4	7 14	226 12	32 6	
92 0	149 6	87 14	143 5	93 1	144 5	84 10	141 3	71 10	1,631 8	—	
9 3	14 15	8 13	14 5	9 5	14 7	8 7	14 2	7 3	163 2	23 5	

3 lbs. Barley (No. 1.), per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
6 2	9 14	6 3	10 2	5 14	10 2	5 10	9 12	5 2	109 14	15 11	
9 10	16 9	10 2	17 9	9 12	18 0	9 6	16 10	8 4	187 1	26 12	
1 10	2 0	1 5	1 8	1 6	1 5	1 0	1 2	0 10	21 8	3 1	
7 6	12 5	7 7	13 0	7 3	12 6	8 11	12 4	7 4	142 14	20 7	
8 0	13 11	8 0	14 0	8 12	14 1	7 0	13 14	7 0	151 8	21 10	
8 10	15 4	7 12	14 10	8 6	15 2	8 3	14 0	8 0	161 8	23 1	
10 2	17 6	9 4	18 0	9 10	17 10	8 10	16 2	9 0	183 8	26 3	
6 4	13 0	6 9	12 6	8 4	12 10	9 0	14 0	7 0	141 3	20 3	
11 0	18 2	10 13	17 12	11 6	18 12	10 10	19 2	10 6	203 8	29 1	
10 7	21 9	10 4	19 0	9 6	17 9	10 5	18 2	9 6	201 1	28 12	
79 3	139 12	77 11	137 15	79 15	137 9	78 7	135 0	72 0	1,503 9	—	
7 15	14 0	7 12	13 13	8 0	13 12	7 13	13 8	7 3	150 6	21 8	

and 7.7 lbs. ; and pulped swedes, 50.4 lbs., per head per day.

TABLE XVI.—EXPERIMENTS made with COWS, at NEWLANDS
7th Week.—February 1

	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Jan. 18.)	MONDAY.		TUESDAY.	
						A.M.	P.M.	A.M.	P.M.
Lot 1.—Special Food.—Unmalted Barley									
Yield of Milk, &c.	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May 20	1,134	11 0	7 1	10 2	7 12
	2	Cross Shorthorn .	6	Oct. 26	1,112	18 0	12 8	18 0	12 12
	3	Cross Shorthorn .	Aged.	April 26	1,127	16 4	8 13	17 0	9 8
	4	Cross Shorthorn .	7	April 21	1,125	12 10	8 12	14 1	8 8
	5	Cross Shorthorn .	8	Sept. 28	994	13 12	9 4	13 0	10 1
	6	Cross Yorkshire .	7	April 14	1,242	13 0	7 7	11 4	7 6
	7	Cross Yorkshire .	10	April 9	1,284	12 6	7 6	13 7	8 8
	8	Cross Ayrshire .	5	June 15	954	15 2	7 4	15 8	8 0
	9	Shorthorn . .	4	June 15	1,198	10 7	6 13	12 0	7 2
10	Shorthorn . .	7	Oct. 29	1,312	17 4	9 10	20 2	11 12	
				Totals .	11,482	140 5	84 14	144 8	90 11
				Averages	1,148	14 0	8 8	14 7	9 1
Lot 2.—Special Food.—Malt and Malt-dst									
Yield of Milk, &c.	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Yorkshire .	8	March 10	1,216	10 2	5 12	10 0	6 2
	2	Cross Shorthorn .	7	Oct. 27	1,024	18 6	9 10	18 12	9 6
	3	Shorthorn . .	8	April 29	1,336	1 8	0 11	1 6	0 15
	4	Cross Shorthorn .	7	June 19	1,047	10 10	7 13	12 5	7 14
	5	Cross Shorthorn .	6	Oct. 29	942	14 10	8 10	14 2	8 1
	6	Cross Welch . .	8	June 18	1,400	15 7	8 0	14 10	9 0
	7	Cross Shorthorn .	7	June 24	1,076	17 10	9 1	17 5	10 4
	8	Cross Yorkshire .	10	April 26	1,132	14 4	7 12	13 14	7 12
	9	Cross Shorthorn .	6	April 19	1,098	18 4	11 2	18 1	10 2
10	Shorthorn . .	9	Nov. 20	1,237	18 0	11 13	19 4	11 6	
				Totals .	11,508	138 13	80 4	139 11	80 14
				Averages	1,151	13 14	8 0	13 15	8 1

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.;

RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
February 7, 1864.

TUESDAY.	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

1); 3 lbs. per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
9 3	10 0	7 0	9 14	8 4	11 3	7 4	12 1	6 9	126 5	18 1	
14 8	17 10	13 7	18 1	14 2	17 6	14 14	19 4	10 7	218 11	31 4	
9 3	16 2	8 13	16 12	10 0	16 5	9 1	17 12	7 13	179 9	25 10	
8 11	12 13	9 3	13 10	9 1	13 10	8 13	14 4	6 12	153 12	21 15	
10 7	14 6	10 3	15 11	9 14	15 6	9 4	17 0	8 6	169 4	24 3	
7 3	13 4	7 2	13 14	7 12	12 12	7 6	13 12	6 0	139 13	20 0	
7 4	12 10	7 7	13 1	8 5	13 6	8 0	13 0	7 2	145 4	20 12	
8 12	15 8	8 4	16 0	7 12	15 10	8 14	17 0	8 2	166 12	23 13	
6 7	11 15	6 2	11 6	7 0	11 12	6 3	12 4	5 10	126 15	18 2	
11 5	21 12	11 2	22 0	10 12	20 10	10 13	21 12	7 15	218 1	31 2	
92 15	146 0	88 11	150 5	92 14	148 0	90 8	158 1	74 12	1,644 6	—	
9 5	14 9	8 14	15 0	9 5	14 13	9 1	15 13	7 8	164 7	23 8	

3 lbs. Barley (No. 1.), per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
6 4	9 6	5 12	9 12	5 14	10 4	5 8	10 4	4 6	108 8	15 8	
10 2	16 0	10 10	17 0	9 8	16 12	9 0	16 14	8 4	187 0	26 11	
0 10	1 0	1 3	1 0	0 7	1 0	0 7	1 2	1 0	13 10	1 15	
7 11	12 15	6 8	13 8	7 7	12 4	7 6	14 12	7 0	140 5	20 1	
9 0	13 4	7 8	13 2	8 1	12 0	6 15	14 0	6 11	150 0	21 7	
8 8	14 10	10 2	12 12	8 2	15 0	7 6	16 0	7 1	161 1	23 0	
10 6	17 5	7 6	17 3	9 15	17 6	9 10	17 4	8 10	186 13	26 11	
8 2	13 7	7 3	13 6	8 6	11 12	7 12	12 13	7 0	147 13	21 2	
11 12	18 2	11 2	19 3	10 11	19 9	10 9	18 3	9 10	207 7	29 3	
10 10	20 4	11 12	20 1	10 11	19 0	10 5	17 5	9 6	210 1	30 0	
83 1	136 5	79 5	136 15	79 2	134 15	74 14	138 9	69 0	1,509 10	—	
8 5	13 10	7 15	13 11	7 15	13 8	7 8	13 14	6 14	150 15	21 9	

chaff, 7·7 lbs.; and pulped swedes, 50·4 lbs., per head per day.

TABLE VIII.—Experiment made with COWS, at NEWLANDS
5th Week.—February 8

	Cows	Breeds	Years old	Dates of milking	Weights of the animals Jan. 1.	MILK		FEED		
						AM.	P.M.	AM.	P.M.	
Lot 1.—Special Food.—Unmalted Barley										
Yield of Milk, &c.	No.				lbs.	lbs. am.	lbs. pm.	lbs. am.	lbs. pm.	
	1	Cross Shorthorn	7	May 20	1,115	11	6	6	7	5
	2	Cross Shorthorn	4	Oct. 26	1,011	17	1	14	3	7
	3	Cross Shorthorn	April	April 26	1,027	15	2	5	12	4
	4	Cross Shorthorn	7	April 27	1,025	14	4	7	12	10
	5	Cross Shorthorn	8	Sept. 26	994	15	1	10	6	6
	6	Cross Yorkshire	7	April 24	1,042	13	0	7	2	7
	7	Cross Yorkshire	10	April 9	1,024	12	0	5	0	13
	8	Cross Yorkshire	5	June 15	953	17	0	5	10	9
	9	Shorthorn	4	June 15	1,136	11	7	6	7	13
	10	Shorthorn	7	Oct. 29	1,012	20	13	12	0	11
Totals					11,482	146	12	99	5	89
Averages					1,148	14	11	9	15	8

Lot 2.—Special Food.—Malt and Malt-dust									
Yield of Milk, &c.	No.				lbs.	lbs. am.	lbs. pm.	lbs. am.	lbs. pm.
	1	Cross Yorkshire	9	March 10	1,216	11	1	5	6
	2	Cross Shorthorn	7	Oct. 27	1,024	17	10	10	0
	3	Shorthorn	8	April 29	1,336	1	0	0	15
	4	Cross Shorthorn	7	June 19	1,047	12	1	7	0
	5	Cross Shorthorn	6	Oct. 29	942	14	10	6	12
	6	Cross Welch	8	June 18	1,400	14	11	8	5
	7	Cross Shorthorn	7	June 24	1,076	17	0	7	13
	8	Cross Yorkshire	10	April 26	1,132	13	11	6	12
	9	Cross Shorthorn	6	April 19	1,098	18	6	10	8
	10	Shorthorn	9	Nov. 20	1,237	18	14	9	8
Totals					11,508	139	0	72	15
Averages					1,151	13	14	7	5

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.;

RM, RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
February 14, 1864.

WEDNESDAY.		THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
A.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

No. 1.; 3 lbs. per head per day.*

A.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
10	7 11	10 0	8 4	11 0	8 13	11 9	7 12	12 0	6 0	129 7	18 8	
13	14 4	17 15	13 2	17 2	14 4	17 5	14 2	19 0	11 15	219 5	31 5	
16	10 2	15 6	9 4	16 3	9 12	16 4	9 5	16 12	8 8	176 8	25 3	
23	8 12	12 3	9 6	12 10	8 7	12 8	9 8	14 1	8 2	153 12	21 16	
25	9 5	16 0	10 2	15 10	9 10	15 2	10 4	16 0	8 12	175 6	25 1	
28	7 5	11 14	7 11	11 3	7 0	12 5	7 5	14 4	6 12	138 2	19 12	
29	7 0	12 5	8 0	12 10	8 2	12 15	8 0	15 0	6 12	143 8	20 8	
29	7 3	15 0	8 11	14 5	8 2	14 7	8 4	16 2	7 10	166 0	23 11	
29	6 0	11 8	6 7	11 0	5 14	12 8	6 10	11 2	5 5	125 2	17 14	
30	7 11 4	19 12	12 4	20 0	11 3	20 7	11 7	20 8	9 4	221 0	31 9	
14	68 14	141 15	93 3	142 0	91 3	145 6	92 9	154 13	79 0	1,648 2	—	
9	8 14	14 3	9 5	14 3	9 2	14 9	9 4	15 8	7 14	164 17	23 9	

No. 3 lbs. Barley (No. 1.), per head per day.*

A.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
9	5 14	9 8	5 10	9 11	6 0	9 4	5 11	10 6	5 11	110 1	15 12	
9	10 0	17 6	9 6	17 10	10 8	17 1	10 8	17 6	8 2	191 7	27 6	
13	0 12	0 12	0 10	0 13	0 10	0 10	0 8	0 7	—	9 9	1 6	
9	8 2	13 0	6 12	11 9	9 10	12 7	8 8	12 7	9 10	143 6	20 8	
8	7 0	12 18	7 2	12 4	7 7	12 12	8 5	14 1	6 0	141 1	20 2	
4	8 3	13 12	7 12	14 7	8 0	14 4	8 6	14 4	7 10	158 13	22 11	
4	9 3	15 14	9 8	16 15	10 3	17 2	9 5	16 14	8 12	179 14	25 11	
2	8 11	11 10	8 2	12 1	8 4	13 4	8 4	12 0	8 2	143 5	20 8	
10	10 12	18 5	10 8	18 12	10 14	19 0	11 12	20 12	10 8	205 15	29 7	
8	9 0	18 0	11 8	18 0	11 10	18 8	10 4	19 6	8 14	201 15	28 14	
1	77 9	131 0	76 14	132 2	83 2	134 4	81 7	137 15	73 5	1,485 6	—	
8	7 12	13 2	7 11	13 4	8 5	13 7	8 2	13 13	7 5	148 9	21 3	

W-chaff, 7·7 lbs.; and pulped swedes, 50·4 lbs., per head per day.

TABLE XVIII.—EXPERIMENTS made with COWS, at NEWL
9th Week.—February

—	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Feb. 15.)	MONDAY.		TUESD
						A.M.	P.M.	A.M.

Lot 1.—Special Food.—Unmalted E

Yield of Milk, &c.	Nos.	Breed.	Years old.	Dates of Calving.	lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May 20	1,116	12 0	8 6	11 4
	2	Cross Shorthorn .	6	Oct. 26	1,095	18 0	14 0	17 6
	3	Cross Shorthorn .	Aged	April 26	1,132	16 10	9 2	16 10
	4	Cross Shorthorn .	7	April 21	1,144	14 0	9 3	12 5
	5	Cross Shorthorn .	8	Sept. 26	1,016	14 11	10 10	15 0
	6	Cross Yorkshire .	7	April 14	1,280	12 0	7 6	12 12
	7	Cross Yorkshire .	10	April 9	1,300	13 8	7 0	12 0
	8	Cross Ayrshire .	5	June 15	964	16 2	8 8	15 6
	9	Shorthorn . .	4	June 15	1,239	11 12	5 10	11 0
	10	Shorthorn . .	7	Oct. 29	1,306	21 0	10 11	20 0
	Totals .				11,592	149 11	90 8	143 11
	Averages				1,159	15 0	9 1	14 6

Lot 2.—Special Food.—Malt and Malt

Yield of Milk, &c.	Nos.	Breed.	Years old.	Dates of Calving.	lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Yorkshire .	8	March 10	1,288	9 2	6 7	9 0
	2	Cross Shorthorn .	7	Oct. 27	1,026	18 0	10 3	17 2
	3	Shorthorn . .	8	April 29	1,392	—	1 0	—
	4	Cross Shorthorn .	7	June 19	1,047	14 2	8 0	11 0
	5	Cross Shorthorn .	6	Oct. 29	934	14 11	8 3	12 1
	6	Cross Welch . .	8	June 18	1,392	15 8	8 15	13 6
	7	Cross Shorthorn .	7	June 24	1,076	18 4	8 5	16 4
	8	Cross Yorkshire .	10	April 26	1,144	13 11	6 8	11 11
	9	Cross Shorthorn .	6	April 19	1,106	19 11	10 14	18 8
	10	Shorthorn . .	9	Nov. 20	1,194	20 4	10 6	19 8
	Totals .				11,597	143 5	78 13	128 8
	Averages				1,160	14 5	7 14	12 14

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14

RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
January 21, 1864.

WEDSDAY.	THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

1.); 3 lbs. per head per day.*

W.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
4	8 10	11 0	8 8	11 2	7 10	11 4	8 4	10 12	8 11	136 8	19 8
4	12 12	17 14	13 0	17 10	12 10	16 3	14 8	16 0	11 6	212 13	30 6
8	9 3	15 15	9 1	15 4	9 10	16 0	9 0	15 2	9 10	175 10	25 1
10	10 2	13 2	9 12	12 1	9 12	12 2	7 14	13 8	7 13	152 4	21 12
3	9 1	15 10	10 1	14 11	10 11	13 11	10 6	15 0	9 13	175 9	25 1
8	7 9	12 8	8 0	12 6	8 3	12 1	8 1	11 8	7 6	139 14	20 0
0	7 4	12 4	8 8	11 12	7 12	12 0	7 13	12 8	6 5	138 5	19 12
3	8 11	16 7	9 6	15 10	9 5	15 8	9 7	15 10	8 12	173 11	24 13
14	6 6	11 3	6 6	10 8	6 6	10 9	5 11	11 4	5 11	119 8	17 1
4	11 0	20 0	11 14	19 14	11 6	19 10	11 2	20 10	9 8	217 11	31 2
4	90 10	145 14	94 8	140 14	93 5	139 0	92 2	141 14	84 15	1,641 13	—
7	9 1	14 9	9 7	14 2	9 5	13 14	9 3	14 3	8 8	164 3	23 7

n 3 lbs. Barley (No. 1.), per head per day.*

W.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
3	5 7	8 12	5 8	8 5	5 7	7 11	5 14	8 7	5 5	101 4	14 7
4	10 0	17 6	9 13	16 12	10 6	16 10	10 4	17 8	8 14	190 6	27 3
-	0 8	—	0 6	—	0 4	—	0 5	—	0 3	3 4	0 7
6	8 10	13 0	8 8	13 1	7 10	11 6	9 0	11 8	9 9	146 6	20 15
12	9 6	12 8	8 6	12 0	8 15	11 6	6 13	10 12	7 0	141 9	20 4
6	7 14	15 2	9 10	14 10	9 10	14 4	7 14	15 12	8 8	165 7	23 10
14	9 4	16 8	9 10	16 4	9 8	15 13	9 10	17 6	9 3	181 14	26 0
11	8 0	12 14	7 10	12 2	7 3	12 0	7 14	11 14	6 6	139 7	19 15
8	11 3	19 8	11 0	17 12	10 14	17 12	11 0	16 14	10 0	206 11	29 8
12	10 6	17 10	10 6	16 11	10 14	18 7	9 12	18 7	9 3	203 4	29 1
6	79 10	133 4	80 13	127 9	80 11	125 6	78 6	128 8	74 3	1,479 8	—
7	7 15	13 5	8 1	12 12	8 1	12 9	7 13	12 14	7 7	147 15	21 2

chaff, 7·7 lbs.; and pulped swedes, 50·4 lbs., per head per day.

TABLE XIX.—EXPERIMENTS made with COWS, at NEWLANDS
10th Week.—February 22

	Cows.	Breed.	Years old.	Dates of Calving.	Weights of the Animals (Feb. 29.)	MONDAY.		TUESDAY.	
						A.M.	P.M.	A.M.	P.M.
Lot 1.—Special Food.—Unmalted Barley									
Yield of Milk, &c.	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Shorthorn .	7	May 29	1,136	11 3	7 8	11 14	7 8
	2	Cross Shorthorn .	6	Oct. 26	1,114	18 4	14 12	16 12	15 8
	3	Cross Shorthorn .	Agd.	April 28	1,150	16 6	9 13	15 10	8 12
	4	Cross Shorthorn .	7	April 21	1,150	13 0	8 6	12 0	8 13
	5	Cross Shorthorn .	8	Sept. 26	1,018	13 2	11 2	14 4	8 10
	6	Cross Yorkshire .	7	April 14	1,312	12 5	8 2	12 6	7 7
	7	Cross Yorkshire .	10	April 9	1,308	13 3	8 15	11 2	7 9
	8	Cross Ayrshire .	5	June 15	974	16 11	9 3	15 3	9 8
	9	Shorthorn . . .	4	June 15	1,232	11 5	6 0	11 0	5 14
	10	Shorthorn . . .	7	Oct. 29	1,347	21 0	11 8	20 0	10 2
Totals .					11,761	146 7	95 5	140 8	86 11
Averages					1,176	14 10	9 8	14 0	8 11
Lot 2.—Special Food.—Malt and Malt-dust									
Yield of Milk, &c.	Nos.				lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
	1	Cross Yorkshire .	8	March 10	1,310	8 0	5 14	7 0	5 2
	2	Cross Shorthorn .	7	Oct. 27	1,022	18 1	10 3	16 6	9 9
	3	Shorthorn . . .	8	April 29	1,428	—	—	—	—
	4	Cross Shorthorn .	7	June 19	1,066	11 10	8 0	11 8	8 1
	5	Cross Shorthorn .	6	Oct. 29	936	11 12	6 14	10 0	7 9
	6	Cross Welch . .	8	June 18	1,396	14 7	8 3	13 2	7 1
	7	Cross Shorthorn .	7	June 24	1,076	17 12	9 6	16 3	10 9
	8	Cross Yorkshire .	10	April 26	1,179	11 8	7 7	11 5	7 3
	9	Cross Shorthorn .	6	April 19	1,092	16 9	9 1	16 8	9 9
	10	Shorthorn . . .	9	Nov. 20	1,214	17 8	10 5	17 12	10 6
Totals .					11,719	127 4	75 5	118 12	73 3
Averages					1,172	12 12	7 9	11 14	7 3

* Also an allowance, which averaged—rape-cake, 2 lbs.; bean-meal, 2 lbs.; clover-chaff, 14 lbs.;

FARM, RUGBY. DETAILED RECORD of the MILK YIELDED, &c.
February 28, 1864.

WEDNESDAY.		THURSDAY.		FRIDAY.		SATURDAY.		SUNDAY.		Total in Seven Days.	Per Head per Day.
A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.	A.M.	P.M.		

No. 1.) ; 3 lbs. per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
11 0	7 4	11 12	7 10	11 0	7 12	11 2	8 2	11 8	7 10	132 5	18 14	
17 6	13 14	17 2	15 5	18 1	12 0	17 10	12 10	18 4	11 9	217 3	31 0	
16 2	9 6	16 4	9 0	14 12	8 7	16 11	9 2	17 0	8 9	175 14	25 2	
13 0	8 7	13 0	8 9	12 3	8 2	13 13	8 8	13 4	8 2	149 1	21 5	
15 0	9 13	15 8	9 10	15 5	9 2	15 12	11 0	15 12	10 2	174 2	24 14	
12 12	7 12	12 6	7 6	11 12	7 14	12 2	7 4	13 3	7 0	139 11	19 15	
12 6	7 6	11 10	7 9	12 1	6 14	12 6	7 2	13 0	5 2	136 7	19 8	
15 8	9 2	16 0	9 2	16 3	9 0	15 12	7 13	14 4	7 9	170 8	24 6	
11 3	5 12	10 14	6 10	10 6	5 3	10 12	5 14	11 6	5 3	117 6	16 12	
20 3	11 8	19 8	11 7	20 0	11 0	20 3	10 12	19 12	10 12	217 11	31 2	
144 8	90 6	144 0	92 4	141 11	85 6	146 3	88 3	147 5	81 10	1,630 4	—	
14 7	9 1	14 6	9 3	14 3	8 9	14 10	8 13	14 12	8 3	163 0	23 5	

from 3 lbs Barley (No. 1.), per head per day.*

lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
8 4	4 12	8 4	4 11	7 5	4 10	8 4	5 6	8 6	4 0	89 14	12 13	
16 0	9 6	16 10	9 12	16 0	9 0	17 0	10 11	17 4	10 6	186 1	26 9	
—	—	—	—	—	—	—	—	—	—	—	—	
12 0	6 14	12 1	8 2	11 4	7 13	11 4	7 8	13 0	7 0	135 9	19 6	
11 12	7 11	12 6	8 0	11 0	7 1	11 6	7 2	13 8	6 11	132 5	18 14	
14 4	8 0	14 2	8 10	14 2	7 3	14 8	8 12	15 4	7 0	154 10	22 1	
16 12	9 2	16 11	10 3	16 3	8 8	17 6	9 12	18 0	8 0	183 14	26 4	
2 0	6 13	12 8	8 0	11 4	6 4	12 2	7 11	13 4	6 8	134 13	19 4	
13 13	9 11	17 3	10 13	16 12	10 2	17 8	11 6	18 10	10 14	190 6	27 3	
10 10	2	18 5	10 13	18 3	9 10	17 6	10 9	17 1	9 7	195 0	27 14	
7 72	7	128 2	79 0	122 1	70 3	126 12	78 13	134 5	69 14	1,402 8	—	
10 7	4	12 13	7 14	12 3	7 0	12 11	7 14	13 7	7 0	140 4	20 1	

aw-chaff, 7·7 lbs. ; and pulped swedes, 50·4 lbs., per head per day.

TABLE XX.—EXPERIMENTS made with COWS, at NEWLANDS FARM, RUGBY.

Per-centage of Cream as shown by the Lactometer, in the mixed Milk of the 10 Cows in each Case.

Dates.	Cows.	Special Food.	PER CENT. CREAM.		
			Morning.	Afternoon.	Mean.
1863: Dec. 20.	Lot 1. .	(Before giving Special Food) Do. Do.	9½	12½	11½
	Lot 2. .		9½	12	10½
.. Dec. 29.	Lot 1. .	Unmalted Barley .	12	13	12½
	Lot 2. .	Malted Barley .	10	12	11
1864: Jan. 5.	Lot 1. .	Unmalted Barley .	12	14½	13½
	Lot 2. .	Malted Barley .	10½	13	11½
.. Jan. 12.	Lot 1. .	Unmalted Barley .	11½	14½	12½
	Lot 2. .	Malted Barley .	10	13½	11½
.. Jan. 19.	Lot 1. .	Unmalted Barley .	10½	13½	12½
	Lot 2. .	Malted Barley .	10	13	11½
.. Jan. 26.	Lot 1. .	Unmalted Barley .	12	14	13
	Lot 2. .	Malted Barley .	10	11½	10½
.. Feb. 2.	Lot 1. .	Unmalted Barley .	11	12½	11½
	Lot 2. .	Malted Barley .	9½	10½	10
.. Feb. 9.	Lot 1. .	Unmalted Barley .	11½	13½	12½
	Lot 2. .	Malted Barley .	10	12	11
.. Feb. 16.	Lot 1. .	Unmalted Barley .	12	14½	13½
	Lot 2. .	Malted Barley .	11	13	12
.. Feb. 23.	Lot 1. .	Unmalted Barley .	12	13½	12½
	Lot 2. .	Malted Barley .	11	12½	11½

SUMMARY.

Dec. 29, 1863, to Feb. 23, 1864	Lot 1. .	Unmalted Barley .	11½	13½	12½
	Lot 2. .	Malted Barley .	10½	12½	11½

and consumed.

Weights. Increase (or Loss) in Live-weight and Milk.

	Cows.				Weights.		Increase (or Loss) in Live-weight.		Milk yielded.		
	Total in 4 Weeks.	Per Head per Day.	Per 1000 lbs. Live-weight per Week.	To produce 100 lbs. Milk.	Dec. 21, 1883.	Jan. 18, 1884.	Total in 4 Weeks.	Per 1000 lbs. Live-weight per Week.	Total in 4 Weeks.	Per Head per Day.	Per 1000 lbs. Live-weight per Week.
Lot 1.—10 Cows; Special Food—Unmalted Barley "No. 1."											
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted					1	1,086	1,134	48	9.43	534½	19.1
Rape Cake	840	3.0	18.6	12.3	2	1,044	1,112	68		891½	31.8
Bean-meal	560	2.0	12.4	8.2	3	1,065	1,127	62		694	24.8
Clover-chaff	3,920	14.0	57.5	8.2	4	1,082	1,125	43		624½	22.3
Straw-chaff	2,156	7.7	47.8	31.6	5	1,020	994	-26		724½	25.9
Pulped Swedes	14,112	50.4	313.1	207.0	6	1,164	1,242	78	582½	20.8	
					7	1,260	1,284	24	600½	21.4	
					8	944	964	10	637½	22.8	
					9	1,156	1,198	42	557½	19.9	
					10	1,238	1,312	76	970½	34.7	
Totals						11,067	11,482	425	—	6,817½	—
Averages						1,106	1,148	42	9.43	681½	24.4
Lot 2.—10 Cows; Special Food—Malted Barley "No. 1." (with Malt-dust).											
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, malted (with its dust)	829½	3.06	18.4*	12.4*	1	1,184	1,216	32	11.17	438½	15.7
Rape Cake	560	2.0	12.4	8.4	2	1,992	1,024	32		771	27.5
Bean-meal	560	2.0	12.4	8.4	3	1,260	1,336	76		376½	13.4
Clover-chaff	3,920	14.0	57.1	8.4	4	1,020	1,047	27		601½	21.5
Straw-chaff	2,156	7.7	47.9	32.3	5	914	942	28		558½	23.5
Pulped Swedes	14,112	50.4	313.4	211.2	6	1,316	1,400	84	659½	23.6	
					7	1,008	1,076	68	744½	26.6	
					8	1,065	1,132	67	654½	23.4	
					9	1,040	1,098	58	783½	28.0	
					10	1,206	1,237	31	997½	35.6	
Totals						11,005	11,508	503	—	6,834½	—
Averages						1,101	1,151	50	11.17	684½	23.9

• These figures represent not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

TABLE XXII.—EXPERIMENTS MADE WITH COWS, AT NEWLANDS FARM, RUGBY.
Record of the Food consumed, of the Weights, of the Increase in Live-weight, and of the Milk yielded.
2nd Period, 4 Weeks January 18 to February 15, 1864.

Food consumed.				Weights. Increase (or Loss) In Live-weight and Milk.							
	Total in 4 Weeks.	Per Head per Day.	Per 1000 lbs. Live- weight per Week.	Cows.	Weights.		Increase (or Loss) in Live-weight.		Milk yielded.		
					Jan. 18.	Feb. 15.	Total in 4 Weeks.	Per 1000 lbs. Live- weight per Week.	Total in 4 Weeks.	Per Head per Day.	
Lot 1.—10 Cows; Special Food—Unmalted Barley "No. 1."											
	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted				1	1,134	1,116	—18		507½	18·1	142½
Rape Cake				2	1,112	1,095	—17		869½	31·1	
Bean-meal				3	1,127	1,132	5		711½	25·4	
Clover-chaff				4	1,125	1,144	19		614½	22·0	
Straw-chaff				5	994	1,016	22	2·38	681½	24·3	
Pulped Swedes	840	3·0	18·2	6	1,242	1,280	38		563½	20·1	
	560	2·0	12·1	7	1,284	1,300	16		574½	20·5	
	560	2·0	12·1	8	954	964	10		644½	23·0	
	3,920	14·0	84·9	9	1,198	1,239	41		508	18·1	
	2,156	7·7	46·7	10	1,312	1,306	—6		901½	32·2	
	14,112	50·4	305·8	Totals	11,482	11,592		—	6,577½	—	
				Averages	1,148	1,159	·38		657½	23·5	
											142½
Lot 2.—10 Cows; Special Food—Malted Barley No. 2" (with Malt-dust).											
	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley malted (with its dust)	829½	3·0*	17·9*	1	1,216	1,288	72		436½	15·6	130½
Rape Cake	560	2·0	12·1	2	1,024	1,024	—		754	26·9	
Bean-meal	560	2·0	12·1	3	1,356	1,352	—4		80	2·9	
Clover-chaff	3,920	14·0	84·9	4	1,047	1,047	—		576½	20·6	
Straw-chaff	2,156	7·7	46·7	5	912	934	—8		648½	21·1	
Pulped Swedes	14,112	50·4	305·4	6	1,400	1,392	—8	1·93	639½	22·8	
				7	1,076	1,076	—		731	28·1	
				8	1,132	1,114	—12		634½	20·8	
				9	1,009	1,106	97		815½	29·1	
				10	1,237	1,194	—43		829½	29·6	
				Totals	11,416	11,487	71		4,001½	—	

TABLE XXIV.—EXPERIMENTS made with COWS, at NEWLANDS FARM, I
SUMMARY of the WEIGHTS, INCREASE in LIVE-WEIGHT, and YIELD of MILK

Cows.	Weights at Com- mence- ment (Dec. 21, 1863).	Increase (or Loss) in Weight.			Weights at Con- clu- sion (Feb. 29, 1864).	Summary of the Increase (or Loss) in Weight.			Yield of Milk	
		In 4 weeks; Dec. 21, 1863, to Jan. 18, 1864.	In 4 weeks; Jan. 18, to Feb. 15.	In 2 weeks; Feb. 15, to Feb. 29.		Total in 10 weeks.	Per Head per week.	Per 1000 lbs. Live- weight per week.	First week.	Last week.
Lot 1.—10 Cows; Special Food—Unmalted Barley "No. 1."										
Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
1	1,086	48	—18	20	1,136	50	5 0	4 8	158 12	132 5
2	1,044	68	—17	19	1,114	70	7 0	6 7½	229 2	217 3
3	1,065	62	5	18	1,150	85	8 8	7 10½	180 7	175 14
4	1,082	43	19	6	1,150	68	6 12½	6 1½	165 14	149 1
5	1,020	—26	22	2	1,018	—2	—0 3½	0 3½	187 4	174 2
6	1,164	78	38	32	1,312	148	14 12½	11 15½	157 6	139 11
7	1,260	24	16	8	1,308	48	4 12½	3 11½	158 4	136 7
8	944	10	10	10	974	30	3 0	3 2	177 13	178 8
9	1,156	42	41	13	1,252	96	9 9½	7 15½	150 12	117 6
10	1,236	76	—6	41	1,347	111	11 1½	8 9½	265 12	217 11
Totals	11,057	425	110	169	11,761	704	—	—	1,829 6	1,630 4
Averages	1,106	42	11	17	1,176	70	7 0½	6 2½	182 15	163 0
Lot 2.—10 Cows; Special Food—Malted Barley "No. 1." (with Malt-dust).										
Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
1	1,184	32	72	22	1,310	126	12 9½	10 1½	125 2	89 14
2	992	32	—	—2	1,022	30	3 0	2 15½	202 8	186 1
3	1,260	76	56	36	1,428	168	16 12½	12 8	137 4	—
4	1,020	27	—	19	1,066	46	4 9½	4 6½	157 15	135 9
5	914	28	—8	2	936	22	2 3½	2 6	173 13	132 5
6	1,316	84	—8	2	1,396	80	8 0	5 14½	169 7	154 10
7	1,008	68	—	—	1,076	68	6 12½	6 8½	187 1	183 14
8	1,065	67	12	35	1,179	114	11 6½	10 2½	180 5	134 13
9	1,040	58	8	—14	1,092	52	5 3½	4 14	202 14	190 6
10	1,208	31	—43	20	1,214	8	0 12½	0 10½	260 15	195 0
Totals	11,005	503	89	122	11,719	714	—	—	1,797 4	1,402 8
Averages	1,101	50	9	12	1,172	71	7 2½	6 4½	179 12	140 4

TABLE XXV.—EXPERIMENTS made with OXEN, at ROTHAMSTED, HERTS.
led Record of the Food consumed, of the Weights, and of the Increase of Live-weight.
1st Period of 4 Weeks; December 1 to December 29, 1863.

Food consumed.					Oxen.	Weights, and Increase (or Loss) in Live-weight.				
-	Total in Four Weeks.	Per Head per Day.	Per 1000 lbs. Live-weight per Week.	To produce 100 lbs. Increase.		Weights.		Increase (or Loss) in Live-weight.		
						Dec. 1.	Dec. 29.	Total in Four Weeks.	Per Head per Week.	Per 1000 lbs. Live-weight per Week.

Lot 1.—10 Oxen; Special Food.—Unmalted Barley "No. 2."

					Nos.	lbs.	lbs.	lbs.	lbs.	lbs.
					1	1,151	1,156	5	1'3	
					2	1,079	1,170	91	22'8	
					3	1,183	1,275	92	23'0	
					4	1,097	1,163	66	16'5	
					5	1,095	1,166	71	17'7	
					6	1,039	1,107	68	17'0	
					7	1,152	1,215	63	15'8	
					8	1,015	1,131	116	29'0	
					9	1,102	1,174	72	18'0	
					10	1,008	1,082	74	18'5	
					Totals .	10,921	11,639	718	—	—
					Averages .	1,092	1,164	71½	18'0	16

Lot 2.—10 Oxen; Special Food.—Malted Barley "No. 2," (with Malt-dust).

					Nos.	lbs.	lbs.	lbs.	lbs.	lbs.
					1	1,204	1,226	22	5'5	
					2	1,057	1,148	91	22'8	
					3	1,148	1,208	60	15'0	
					4	1,088	1,142	54	13'5	
					5	1,176	1,184	8	2'0	
					6	1,092	1,079	—13	—3'2	
					7	1,078	1,090	12	3'0	
					8	1,029	1,085	36	9'0	
					9	1,078	1,661	88	22'0	
					10	1,008	1,156	148	37'0	
					Totals .	10,958	11,464	506	—	—
					Averages .	1,096	1,146	50½	12'7	11½

Figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which it be produced.

TABLE XXVI.—EXPERIMENTS made with OXEN, at ROTHAMSTED, HERTS.
 Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight.
 2nd Period of 4 Weeks: December 29, 1863, to January 26, 1864.

Food consumed.					Weights, and Increase in Live-weight.					
	Total Food per Week.	Per Head per Day.	Per 100 lbs. Live- weight per Week.	To produce 100 lbs. Increase.	Oxen.	Weights.		Increase in Live-weight.		
						Dec. 29, 1863.	Jan. 26, 1864.	Total in Four Weeks.	Per Head per Week.	Per 100 lbs. Live- weight per We
Lot 1.—10 Oxen; Special Food—Unmalted Barley "No. 2"										
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.
					1	1,156	1,228	72	18.0	
					2	1,170	1,222	52	13.0	
					3	1,275	1,368	93	23.3	
					4	1,163	1,219	56	14.0	
					5	1,166	1,222	56	14.0	
Barley, unmalted	1,120	4.0	23.3	147.0	6	1,107	1,185	78	19.5	15
Clover-chaff . .	2,800	10.0	58.2	367.5	7	1,215	1,323	108	27.0	
Swedes, ad lib.	25,511	91.1	530.6	3347.9	8	1,131	1,208	77	19.2	
					9	1,174	1,268	94	23.5	
					10	1,082	1,158	76	19.0	
					Totals . .	11,639	12,401	762	—	—
					Averages .	1,164	1,240	76.4	19.0	15
Lot 2.—10 Oxen; Special Food—Malted Barley "No. 2" (with Malt-dust).										
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.
					1	1,228	1,323	97	24.3	
					2	1,148	1,208	70	17.5	
					3	1,208	1,316	108	27.0	
					4	1,142	1,232	90	22.5	
					5	1,184	1,262	78	19.5	
Barley, malted, } with its dust }	1,120*	4.0*	23.6*	142.5*	6	1,079	1,132	53	13.2	10
Clover-chaff . .	2,800	10.0	59.0	356.2	7	1,090	1,164	74	18.5	
Swedes, ad lib.	26,067	93.1	549.4	3315.1	8	1,065	1,162	97	24.3	
					9	1,166	1,225	59	14.7	
					10	1,156	1,216	60	15.0	
					Totals . .	11,464	12,250	786	—	—
					Averages .	1,146	1,225	78.6	19.6	10

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they should be produced.

TABLE XXVII.—EXPERIMENTS made with OXEN, at ROTHAMSTED, HERTS.
ailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight.
3rd Period of 4 Weeks; January 26 to February 23, 1864.

Food consumed.					Oxen.	Weights, and Increase in Live-weight.				
	Total in Four Weeks.	Per Head per Day.	Per 1000 lbs. Live- weight per Week.	To produce 100 lbs. Increase.		Weights.		Increase in Live-weight.		
						Jan. 26.	Feb. 23.	Total in Four Weeks.	Per Head per Week.	Per 1000 lbs. Live- weight per Week.

Lot 1.—10 Oxen; Special Food—Unmalted Barley "No. 2."

	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.
Unmalted	1,120	4.0	21.9	156.3	1	1,228	1,299	71	17.8	14½
"	560	2.0	11.0	77.1	2	1,222	1,274	52	13.0	
chaff	3,360	12.0	65.8	462.8	3	1,368	1,465	97	24.2	
total	21,757	77.7	426.1	2996.8	4	1,219	1,302	83	20.8	
					5	1,222	1,291	69	17.2	
					6	1,185	1,260	75	18.8	
					7	1,323	1,393	70	17.5	
					8	1,208	1,288	80	20.0	
					9	1,268	1,351	83	20.7	
					10	1,158	1,204	46	11.5	
					Totals	12,401	13,127	726	—	—
					Averages	1,240	1,313	72½	18.2	14½

Lot 2.—10 Oxen; Special Food—Malted Barley "No. 2" (with Malt-dust).

	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.
Malted	1,120*	4.0*	22.3*	177.5*	1	1,323	1,400	77	19.3	12½
dust	560	2.0	11.1	88.7	2	1,218	1,295	77	19.3	
"	3,360	12.0	66.9	532.5	3	1,316	1,383	67	16.7	
total	20,930	74.8	416.4	3317.0	4	1,232	1,274	42	10.5	
					5	1,262	1,325	63	15.8	
					6	1,132	1,185	53	13.2	
					7	1,164	1,253	89	22.3	
					8	1,162	1,197	35	8.7	
					9	1,225	1,288	63	15.8	
					10	1,216	1,281	65	16.2	
					Totals	12,250	12,881	631	—	—
					Averages	1,225	1,288	63	15.8	12½

figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which
 d be produced.

TABLE XXVIII.—EXPERIMENTS made with OXEN, at ROTHAMSTED, Herts.
 Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight
 4th Period of 4 Weeks; February 23 to March 22, 1864.

Food consumed.					Weights, and Increase (or Loss) in Live-weight.					
—	Total in Four Weeks.	Per Head per Day.	Per 1000 lbs. Live- weight per Week.	To produce 100 lbs. increase.	Oxen.	Weights.		Increase (or Loss) in Live-weight.		
						Feb. 23.	Mar. 22.	Total in Four Weeks.	Per Head per Week.	Per 1000 lb. Live- weight per We
Lot 1.—10 Oxen; Special Food—Unmalted Barley "No. 2."										
					Nos.	lbs.	lbs.	lbs.	lbs.	
					1	1,299	1,316	17	4.3	
					2	1,274	1,328	54	13.5	
					3	1,465	1,470	5	1.2	
	lbs.	lbs.	lbs.	lbs.	4	1,302	1,314	12	3.0	
Barley, unmalted	1,120	4.0	21.3	1287.4	5	1,291	1,310	19	4.8	
Oilcake	560	2.0	10.6	643.7	6	1,260	1,238	-22	-5.5	
Clover-chaff	3,360	12.0	63.8	3862.1	7	1,393	1,392	-1	-0.3	
Swedes, <i>ad lib.</i>	18,175	64.9	345.0	20890.8	8	1,288	1,294	6	1.5	
					9	1,351	1,353	2	0.5	
					10	1,204	1,198	-5	-1.2	
					Totals	13,127	13,214	87	—	—
					Averages	1,313	1,321	8.7	2.2	
Lot 2.—10 Oxen; Special Food—Malted Barley "No. 2 (with Malt-dust)."										
					Nos.	lbs.	lbs.	lbs.	lbs.	
					1	1,400	1,437	37	9.3	
					2	1,295	1,300	5	1.2	
					3	1,383	1,430	47	11.8	
					4	1,274	1,311	37	9.2	
Barley, malted, } with its dust }	1,120*	4.0*	21.5*	388.9*	5	1,325	1,334	9	2.3	
Oilcake	560	2.0	10.7	194.4	6	1,185	1,210	25	6.2	
Clover-chaff	3,360	12.0	64.5	1166.7	7	1,253	1,276	23	5.8	
Swedes, <i>ad lib.</i>	19,332	69.0	371.1	6712.5	8	1,197	1,252	55	13.7	
					9	1,288	1,311	23	5.8	
					10	1,281	1,308	27	6.7	
					Totals	12,881	13,169	288	—	—
					Averages	1,288	1,317	28.8	7.2	

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

PLATE XXIX.—EXPERIMENTS made with OXEN, at ROTHAMPTON, HERTS.
led Record of the Food consumed, of the Weights, and of the Increase in Live-weight.
5th Period of 4 Weeks; March 22 to April 19, 1864.

Food consumed.					Oxen.	Weights, and Increase in Live-weight.				
	Total in Four Weeks.	Per Head per Day.	Per 1000 lbs. Live- weight per Week.	To produce 100 lbs. Increase.		Weights.		Increase in Live-weight.		
						Mar. 22.	Apr. 19.	Total in Four Weeks.	Per Head per Week.	Per 1000 lbs. Live- weight per Week.
Lot 1.—10 Oxen; Special Food—Unmalted Barley "No. 2."										
					Nos.	lbs.	lbs.	lbs.	lbs.	lbs.
unmalted " " " half " " Swedes " Mangolds	lbs. 1,120 1,120 3,360 3,856 13,980	lbs. 4·0 4·0 12·0 13·8 49·0	lbs. 20·3 20·3 61·0 70·0 263·9	lbs. 101·8 101·8 305·5 350·5 1,270·9	1	1,316	1,404	88	22·0	20
					2	1,328	1,386	58	14·5	
					3	1,470	1,591	121	30·3	
					4	1,314	1,404	90	22·5	
					5	1,310	1,422	112	28·0	
					6	1,238	1,344	106	26·5	
					7	1,392	1,566	174	43·5	
					8	1,294	1,424	130	32·5	
					9	1,353	1,482	129	32·2	
					10	1,199	1,291	92	23·0	
Totals .					13,214	14,314	1,100	—	—	
Averages .					1,321	1,431	110	27·5	20	
Lot 2.—10 Oxen; Special Food—Malted Barley "No. 2" (with Malt-dust).										
					Nos.	lbs.	lbs.	lbs.	lbs.	lbs.
malted, dust } " " " half " " Swedes " Mangolds	lbs. 1,120* 1,120 3,360 4,347 14,490	lbs. 4·0* 4·0 12·0 15·5 51·8	lbs. 20·7* 20·7 62·0 80·2 267·2	lbs. 144·7* 144·7 434·1 561·6 1,872·1	1	1,437	1,525	88	22·0	14½
					2	1,300	1,422	122	30·5	
					3	1,430	1,525	95	23·8	
					4	1,311	1,396	85	21·2	
					5	1,334	1,444	110	27·5	
					6	1,210	1,256	46	11·5	
					7	1,276	1,346	70	17·5	
					8	1,252	1,252	—	—	
					9	1,311	1,385	74	18·5	
					10	1,308	1,392	84	21·0	
Totals .					13,169	13,943	774	—	—	
Averages .					1,317	1,394	77½	19·4	14½	

Figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which
 to be produced.

TABLE XXX.—EXPERIMENTS made with OXEN, at ROTHAMPTON, HERT
 Summary of the Weights, and of the Increase in Live-weight. Total Period 20 Weeks
 December 1, 1863, to April 19, 1864.

OXEN.	Weights at Commencement, Dec. 1, 1863.	Increase (or Loss) in Weight.					Weights at Conclusion, April 19, 1864.	Summary of the Increase in Weight.		
		In Four Weeks; Dec. 1 to Dec. 29, 1863.	In Four Weeks; Dec. 29, 1863, to Jan. 26, 1864.	In Four Weeks; Jan. 26 to Feb. 23, 1864.	In Four Weeks; Feb. 23 to Mar. 22, 1864.	In Four Weeks; Mar. 22 to Apr. 19, 1864.		Total in Twenty Weeks.	Per Head per Week.	Per 1000 Live weight per Week.

Lot 1.—10 Oxen; Special Food—Unmalted Barley "No. 2."

Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. c.
1	1,151	5	72	71	17	88	1,404	253	12 10	9
2	1,079	91	52	52	54	58	1,386	307	15 6	12
3	1,183	92	93	97	5	121	1,591	408	20 6	14
4	1,097	66	56	83	12	90	1,404	307	15 6	12
5	1,095	71	56	69	19	112	1,422	327	16 6	13
6	1,039	68	78	75	—22	106	1,344	305	15 4	12
7	1,152	63	108	70	—1	174	1,566	414	20 11	15
8	1,015	116	77	80	6	130	1,424	409	20 7	16
9	1,102	72	94	83	2	129	1,482	380	19 0	14
10	1,008	74	76	46	—5	92	1,291	283	14 2	12
Totals	10,921	718	762	726	87	1,100	14,314	3,393	—	—
Averages	1,092	71½	76¼	72½	8½	110	1,431½	339¼	16 15	13

Lot 2.—10 Oxen; Special Food—Malted Barley "No. 2" (with Malt-dust).

Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. c.
1	1,204	22	97	77	37	88	1,525	321	16 1	11
2	1,067	91	70	77	5	122	1,422	355	18 4	14
3	1,148	60	108	67	47	95	1,525	377	18 14	14
4	1,088	54	90	42	37	85	1,396	308	15 6	12
5	1,176	8	78	63	9	110	1,444	268	13 6	10
6	1,092	—13	53	53	25	46	1,266	164	8 3	7
7	1,078	12	74	89	23	70	1,346	268	13 6	11
8	1,029	36	97	35	55	—	1,252	223	11 2	9
9	1,078	88	59	63	23	74	1,386	307	15 6	12
10	1,006	148	60	65	27	84	1,392	384	19 3	16
Totals	10,958	506	786	631	288	774	13,943	2,965	—	—
Averages	1,095½	50½	78½	63	28½	77½	1,394½	296½	14 15	13

Periods.		Food consumed.										Increase in Weight.	
Dates.	Number of Weeks.	Per Head per Week.				Per 1000 lbs. Live-weight per Week.						Per Head per Week.	Per 1000 lbs. Live-weight per Week.
		Barley (or Malt),	Oats,	Clover-chaff.	Roots.	Barley (or Malt),	Oats,	Clover-chaff.	Roots.	To produce 100 lbs. increase.	Roots.		
Lot 1.—10 Oxen; Special Food—Unmalted Barley "No. 2."													
1863-4.		lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Dec. 1 to Dec. 29	4	28-0	—	56-0	631½	24-8	—	49-6	531-1	156-0	—	312-0	3,464-4
Dec. 29 to Jan. 26	4	28-0	—	70-0	637½	23-3	—	58-2	530-6	147-0	—	367-5	3,347-9
Jan. 26 to Feb. 23	4	28-0	14-0	84-0	544	21-9	11-0	65-8	426-1	154-3	77-1	462-8	2,996-8
Feb. 23 to March 22	4	28-0	14-0	84-0	454½	21-3	10-6	63-8	345-0	1287-4	643-7	3902-1	20,890-8
March 22 to April 19	4	28-0	28-0	84-0	446	20-3	20-3	61-0	324-6	101-8	101-8	306-5	1,621-5
Dec. 1 to April 19	20	28-0	11-2	75-6	540½	22-3	8-9	59-9	428-6	165-1	66-0	445-6	3,187-3
													16 15½
													13 7½
Lot 2.—10 Oxen; Special Food—Malted Barley "No. 2" (with Malt-dust).													
1863-4.		lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Dec. 1 to Dec. 29	4	28-0	—	56-0	593½	25-0	—	50-0	529-6	221-3	—	442-7	4,693-5
Dec. 29 to Jan. 26	4	28-0	—	70-0	651½	23-6	—	59-0	549-4	214-5	—	356-2	3,316-1
Jan. 26 to Feb. 23	4	28-0	14-0	84-0	523½	22-3	11-1	66-9	416-4	177-5	88-7	532-5	3,317-0
Feb. 23 to March 22	4	28-0	14-0	84-0	457½	21-5	10-7	64-5	371-1	238-9	194-4	1166-7	6,712-5
March 22 to April 19	4	28-0	28-0	84-0	471	20-7	20-7	62-0	347-4	144-7	144-7	434-1	2,433-7
Dec. 1 to April 19	20	28-0	11-2	75-6	544½	22-5	9-0	60-7	437-4	187-6	75-0	506-5	3,648-4
													14 14½
													11 16½

* These figures represent not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

TABLE XXXIV.—EXPERIMENTS MADE WITH SHEEP, at ROTHAMPTON, HERTS.

Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight.
1st Period of 4 Weeks ; December 2 to December 30, 1863.

Food consumed.					Weights and Increase in Live-weight.				
—	Total in 4 Weeks.	Per Head per Day.	Per 100 lbs. Live-weight per Week.	To produce 100 lbs. Increase.	Weights.		Increase in Live-weight.		
					Dec. 2, 1863.	Dec. 30, 1863.	Total 4 Weeks.	Per 100 lbs. Live-weight per Week.	
Lot 1.—12 Sheep; Special Food—Unmalted Barley "No. 1."									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted	268	0.75	4.94	183.9	1	112	121	9	2.25
					2	101	115	14	3.50
Clover-chaff	335	1.00	6.59	945.3	3	101	114	17	3.50
					4	107	114	7	1.75
Sweden, old No.	3,698	10.98	75.29	2,698.0	5	100	109	8	3.00
					6	93	104	12	3.00
					7	100	110	10	3.50
					8	105	119	14	3.50
					9	88	108	12	3.50
					10	119	121	14	3.50
					11	88	100	12	3.00
					12	94	106	12	3.00
					Totals . .	1,207	1,244	187	—
					Averages .	101	112	114	2.69
Lot 2.—12 Sheep; Special Food—Malted Barley "No. 1" (with Malt-dust).									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
Barley, malted, with its dust	249°	0.74°	4.97°	128.7°	1	109	120	11	2.75
					2	94	104	10	2.50
					3	96	105	9	2.25
					4	100	108	8	2.00
					5	119	123	14	2.00
					6	112	126	14	3.50
					7	104	114	10	2.50
					8	94	114	19	3.00
					9	108	114	10	3.00
					10	104	114	10	3.00
					11	110	120	10	3.00
					12	105	114	10	3.00
					Totals . .	1,207	1,244	187	—
					Averages .	101	112	114	2.69

Barley, unmalted	lbs.	lbs.	lbs.	lbs.	114	159	6	2'00
Clover-chaff	252	0'75	4'96	108'4	100	116	11	2'75
Swedes, ad lib.	336	1'00	6'62	264'5	98	109	12	3'00
	3,668	10'95	72'47	2,597'6	8	91	7	1'75
					10	104	8	2'00
					11	104	12	2'50
					12	98	10	3'75
Totals . . .					1,206	1,333	127	—
Averages .					100	111	104	2'63
								2'50

Lot 4.—12 Sheep; Special Food—Malted Barley "No. 2" (with Malt-dust).

Barley, malted, with its dust . .	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Clover-chaff	252*	0'75*	4'92*	108'8*	119	126	12	1'75
Swedes, ad lib.	336	1'00	6'57	258'5	114	105	11	3'00
	3,668	10'92	71'75	2,521'5	94	105	11	2'75
					102	114	12	3'00
					91	103	12	3'00
					97	102	12	2'75
					102	114	12	3'00
					94	104	10	2'50
					100	110	10	2'50
					91	100	9	2'25
Totals . . .					1,213	1,343	120	—
Averages .					101	112	104	2'71
								2'54

Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley No. 2 (with Malt-dust).

Barley, unmalted	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, malted, with its dust . .	168	0'50	3'33	117'5	158	144	16	4'00
Clover-chaff	336	1'00	6'65	255'0	114	126	12	3'00
Swedes, ad lib.	3,668	10'92	71'63	2,565'0	104	114	10	2'50
					94	108	14	3'50
					96	112	16	4'00
					94	103	11	3'75
					202	114	12	3'00
					91	103	12	3'00
					88	108	10	2'50
					91	108	10	2'50
					91	101	10	2'50
Totals . . .					1,191	1,324	143	—
Averages .					99	111	12	2'98
								2'83

* These figures represent not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

	lbs.	lbs.	lbs.	1 st cut.	2 ^d cut.	3 ^d cut.
Barley, unmalted	552	4.53	225.0	129	134	37.00
				3	122	37.00
				4	116	37.50
				5	116	37.50
				6	106	37.50
				7	117	1.00
				8	91	1.75
				9	112	37.50
				10	122	37.50
				11	105	37.00
				12	93	37.00
				12	100	1.75
Totals	3,562.8	66.54	3,562.8	1,353	1,446	—
Averages				111	130	37.03

Lot 4.—12 Sheep; Special Food—Malted Barley "No. 2" (with Malt-dust).

[illegible]

Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley "No. 2 (with Malt-dust).

	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted	168	3'50	163'1	144	130	6	1'50	1'50	1'50
Barley, malted, with its dust . .	84*	0'25*	81'6*	136	120	10	2'30	2'30	2'30
Clover-chaff	336	1'00	328'2	114	100	8	1'50	1'50	1'50
Sweedes, <i>ad lib.</i>	3,698	63'46	3,592'3	108	118	10	2'50	2'50	2'50
				108	106	8	2'25	2'25	2'25
				111	108	10	2'00	2'00	2'00
				112	101	10	2'30	2'30	2'30
				Totals . .	1,334	103	—	—	—
				Averages .	111	84	2'15	2'15	2'15

* These figures represent not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

Unmalted Barley "No. 2."

	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted	252	0.75	4.19	2	134	134	144	9	2.95
				3	134	144	144	10	2.90
				4	134	133	142	8	2.90
				5	130	142	136	9	2.90
				6	117	136	136	9	2.95
				7	132	110	136	9	2.90
				8	132	133	133	11	2.95
				9	122	133	133	11	2.95
				10	115	134	134	10	2.90
				11	100	110	110	10	2.90
				12	100	110	110	10	2.90
Totals					1446	1358	1358		
Averages					121	120	120	9½	2.93
									1.86

Lot 4.—12 Sheep; Special Food—Malted Barley "No. 2 (with Malt-dust).

	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, malted, with its dust	231*	0.75*	4.27*	1	132	140	147	8	2.90
				2	136	147	147	11	2.75
				3	113	180	180	8	2.90
				4†	(105)	(95)	(95)	(-10)	—
				5	130	140	140	10	2.90
				6	109	110	110	7	2.75
				7	110	120	120	11	2.90
				8	110	120	120	12	3.00
				9	114	126	126	6	1.90
				10	112	118	118	6	1.90
				11	116	128	128	12	3.00
				12	112	118	118	6	1.90
Totals					1392	1400	1400	103	—
Averages					118	126	126	9½	2.94
									1.90

Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley "No. 2 (with Malt-dust).

	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted	166	0.50	2.83	1	150	163	163	13	3.25
				2	136	144	144	8	2.90
				3	110	118	118	8	2.90
				4	120	128	128	8	2.90
				5	118	126	126	8	2.90
				6	112	119	119	10	2.90
				7	112	123	123	10	2.90
				8	104	123	123	8	2.90
				9	112	118	118	6	1.90
				10	112	122	122	6	1.90
				11	108	114	114	6	1.90
				12	111	116	116	5	1.95
Totals					1457	1533	1533	96	—
Averages					120	128	128	8	2.90
									1.68

* These figures represent not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.
† This sheep had been observed to be unwell for about 8 weeks and was killed on 15th. Its consumption of food and weights are included in the calculated results given above.

TABLE XXXVII.—EXPERIMENTS MADE WITH SHEEP, AT ROTHEAMSTEAD, HERTS.

Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight.

4th Period of 4 Weeks; February 24 to March 23, 1864.

	Food consumed.			Sheep.	Weights, and Increase in Live-weight.			
	Total in 4 Weeks.	Per Head per Day.	Per 100 lbs. Live-weight per Week.		Weights.		Increase in Weight.	
					Feb. 24, 1864.	March 23, 1864.	Total in 4 Weeks.	Per Head Live-weight per Week.
Lot 1.—12 Sheep; Special Food—Unmalted Barley "No. 1."								
Barley, unmalted Clover-chaff, Sweden, ad lib.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
	268	0.75	3.96	1	133	137	4	1.00
				2	134	143	8	2.00
	236	1.00	5.28	3	136	144	8	2.00
				4	138	153	15	1.25
	4219	12.54	66.25	5	139	153	14	1.00
				6	131	137	6	1.50
				7	138	133	5	1.25
				8	140	130	10	2.50
				9	135	130	5	1.00
				10	130	134	4	1.00
				11	110	114	4	1.00
				12	122	127	5	1.25
			Totals . .	1556	1623	67	—	
			Averages .	130	135	54	1.40	
							1.05	
Lot 2.—12 Sheep; Special Food—Malted Barley "No. 1" (with Malt-dust).								
Barley, malted, with its dust Clover-chaff, Sweden, ad lib.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
	249*	0.74*	3.93*	1	134	144	10	2.50
				2	128	136	8	1.50
	236	1.00	5.31	3	123	134	11	1.50
				4	128	134	6	1.50
	4161	12.38	65.74	5	126	133	7	1.75
				6	144	151	7	1.75
				7	114	122	8	2.00
				8	126	130	4	1.00
				9	140	150	10	2.50
				10	119	125	6	1.50
				11	119	123	4	2.25
				12	114	116	2	0.50
							1.31	

Barley, unmalted	lbs.	0 75	lbs.	345*2	lbs.	144	150	6	1'20
						144	147	3	0'75
						142	142	10	2'90
						142	154	12	3'00
Clover-chaff						116	116	0	0'00
						116	114	4	1'00
						132	136	4	1'00
						133	143	9	3'25
Swedes, <i>ad lib.</i>		12'35	65'08	569*3		124	135	11	2'75
						110	117	7	1'75
Totals . . .						1556	1631	73	—
Averages . .						130	136	6	1'52

Lot 4.—11† Sheep; Special Food—Malted Barley "No. 2 with Malt-dust".

Barley, malted, with its dust . .	lbs.	0'75*	lbs.	399*1*	Nos.	lbs.	lbs.	lbs.	lbs.
					1	140	142	2	0'50
					2	147	154	7	1'75
					3	120	126	6	1'50
					4†	—	—	—	—
					5	140	147	7	1'75
					6	136	136	6	1'50
					7	128	130	6	1'50
					8	128	131	5	1'25
					9	126	131	5	1'25
					10	118	118	0	0'00
					11	128	136	8	2'00
					12	118	132	4	1'00
Totals . . .						1405	1464	59	—
Averages . .						126	133	54	1'34

Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley "No. 2 (with Malt-dust)".

Barley, unmalted	lbs.	0'50	lbs.	233*3	Nos.	lbs.	lbs.	lbs.	lbs.
					1	163	174	11	2'75
					2	144	144	0	0'00
					3	118	126	8	2'00
					4	128	133	5	1'25
					5	126	130	4	1'00
					6	123	127	7	2'00
					7	123	130	7	1'75
					8	132	136	4	1'00
					9	118	126	8	2'00
					10	122	126	4	1'00
					11	114	119	5	1'25
					12	116	124	8	2'00
Totals . . .						1333	1605	72	—
Averages . .						128	134	6	1'50

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

† No. 4 sheep of this lot (d) was killed Feb. 3. See Note to Table XXXV.

Lot 4.—11† Sheep; Special Food—Malted Barley No. 2 (with Malt-dust).

[illegible]

Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley “No. 2 (with Malt-dust)."

[illegible]

* These figures represent not the actual weights of malt and malt-ding, but the weight of the barley from which they would be produced. No. 4 sheep of this lot (4) was killed Feb. 3. See note to Table XXXVI.

TABLE XXXIX.—EXPERIMENTS MADE WITH SHEEP, at ROTHAMSTED, HERTS.

Summary of the Weights, and of the Increase in Live-weight.

Total Period, 20 Weeks; December 2, 1863, to April 20, 1864.

SHEEP.	Weights at Commencement, Dec. 2, 1863.	Increase (or Loss) in Weight.					Weights at Conclusion, April 20, 1864.	Summary of the Increase in Live-weight.		
		In 4 Weeks.						Total in 20 Weeks.	Per Head per Week.	Per 100 lbs. Live-weight per Week.
		In 4 Weeks; Dec. 2, to Dec. 20, 1863.	In 4 Weeks; Jan. 27, to Jan. 31, 1864.	In 4 Weeks; Feb. 24, to Mar. 23, 1864.	In 4 Weeks; Mar. 23, to April 20, 1864.					
Lot 1.—12 Sheep; Special Food—Unmalted Barley "No. 1."										
No. 1	lbs. 112	lbs. 9	lbs. 7	lbs. 4	lbs. 5	lbs. 142	lbs. 30	lbs. 0.75	lbs. 0.3	
2	101	12	10	8	8	130	49	2 7½	1 12½	
3	101	13	14	8	5	154	53	2 10½	2 1½	
4	107	7	8	5	4	137	30	1 8	1 3½	
5	100	8	10	4	8	140	40	2 0	1 10½	
6	92	12	11	6	6	134	42	2 1½	2 0	
7	100	10	10	5	7	140	40	2 0	1 13½	
8	105	14	10	10	12	143	45	2 10½	2 0½	
9	108	12	14	4	4	133	45	2 12½	2 0½	
10	119	9	10	4	7	161	49	2 1½	1 8	
11	98	5	5	4	5	119	31	1 8½	1 11½	
12	94	8	8	5	6	123	39	1 15½	1 11½	
Totals . . .	1307	94	116	67	82	1,705	498	—	—	
Averages . .	101	7½	9½	5½	6½	142	41½	2 1½	1 11½	
Lot 2.—12 Sheep; Special Food—Malted Barley "No. 1" (with Malt-dust).										
No. 1	lbs. 109	lbs. 11	lbs. 8	lbs. 10	lbs. 10	lbs. 154	lbs. 45	lbs. 0.38	lbs. 1.11½	
2	94	10	12	6	8	136	42	2 4	1 11½	
3	96	9	8	6	8	126	40	2 1½	1 13½	
4	100	12	8	6	6	140	40	2 0	1 11½	
5	119	14	11	7	8	171	52	2 0	1 10½	
6	112	16	8	8	8	161	49	2 9½	1 12½	
7	94	4	6	7	4	126	32	1 7½	1 7½	
8	102	12	8	6	4	137	33	1 12	2 1½	
9	103	13	12	10	10	140	37	1 19	2 1½	
10	94	7	5	6	7	124	34	1 11½	1 11½	
11	96	14	8	11	3	140	44	2 3½	1 13½	
12	98	14	4	3	8	124	36	1 13½	1 11½	

4	114	8	12	10	3	3	150	36	1 129	1 107
5	105	11	12	10	10	5	147	42	2 138	2 98
6	105	11	14	12	12	10	164	59	2 138	2 98
7	84	12	11	9	—	4	130	36	0 128	0 142
8	84	10	11	9	—	3	117	19	1 128	1 132
9	104	7	7	13	6	6	120	36	1 128	1 74
10	104	8	10	10	4	4	140	36	1 128	1 14
11	104	10	10	11	4	10	152	48	2 61	2 54
12	88	12	12	9	11	7	142	54	2 116	2 54
	86	7	7	10	7	9	125	40	2 0	1 144
Totals . . .	1206	127	113	112	73	78	1,709	508	—	—
Averages . .	100	104	94	94	6	64	143	42	2 14	1 114

Lot 4.—12* Sheep; Special Food—Malted Barley "No. 2" (with Malt-dust).

No.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.
1	112	12	10	10	7	13	129	31	1 94	1 124
2	114	11	11	11	6	8	164	50	2 0	1 12
3	94	12	—	(-10 ⁶)	—	—	—	40	—	—
4	102	12	—	—	—	—	—	49	2 74	1 134
5	104	12	—	—	—	—	—	57	1 11	1 11
6	91	12	—	—	—	—	—	57	2 134	2 74
7	107	12	—	—	—	—	—	59	2 144	2 8
8	104	12	—	—	—	—	—	53	1 144	1 74
9	102	12	—	—	—	—	—	53	1 94	1 74
10	94	10	—	—	—	—	—	46	2 44	1 14
11	100	10	—	—	—	—	—	39	1 134	1 124
12	91	9	12	6	4	8	130	—	—	—
Totals . . .	1213	130	64	103	59	95	1,539	448	—	—
Averages . .	101	104	54	94	54	84	143	404	2 04	1 104

Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley "No. 2" (with Malt-dust).

No.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.
1	128	16	6	13	11	12	166	56	2 144	1 134
2	124	12	10	8	8	—14	120	32	0 144	0 8
3	94	10	6	8	6	9	130	38	1 144	1 124
4	104	10	6	8	5	10	142	41	1 144	1 124
5	94	14	10	8	4	10	140	46	2 44	1 124
6	94	16	7	10	8	8	145	49	2 74	2 04
7	94	11	8	10	7	6	136	42	2 14	1 134
8	102	12	10	8	4	10	145	44	2 34	1 124
9	11	12	9	6	8	2	145	43	2 34	1 124
10	88	12	10	6	6	—	132	35	1 124	1 64
11	88	10	10	6	8	—	116	28	1 64	1 6
12	91	10	10	5	8	8	132	41	2 04	1 134
Totals . . .	1191	143	103	96	72	61	1,666	475	—	—
Averages . .	99	12	84	8	6	5	139	394	1 154	1 104

* No. 4 sheep of this lot was killed on February 3, and its weight was only taken into calculation in the total and average results given for the first and second months.

TABLE XL.—EXPERIMENTS MADE WITH SHEEP, at ROTHAMSTED, HERTS.

Summary of Food consumed, and Increase yielded, during each separate Month, and during the entire Period of Five Months.

Periods.		Food consumed.										Increase in Weight.	
Dates.	Number of Weeks.	Per Head per Week.			Per 100 lbs. Live-weight per Week.			To produce 100 lbs. increase.			Per Head per Week.	Per 100 lbs. Live-weight per Week.	
		Barley (or Malt.)	Clover-chaff.	Roots.	Barley (or Malt.)	Clover-chaff.	Roots.	Barley (or Malt.)	Clover-chaff.	Roots.			
Lot 1.—12 Sheep: Special Food—Unmalted Barley "No. 1."													
1883-4.													
Dec. 2 to Dec. 30.	4	5.3	7.0	76.8	4.94	6.59	72.29	183.9	215.3	2692.0	2	134	2 11
Dec. 30 to Jan. 27.	4	5.3	7.0	76.3	4.53	6.04	65.85	264.1	337.4	3497.9	1	104	1 11
Jan. 27 to Feb. 24.	4	5.3	7.0	74.5	4.21	5.61	59.69	213.6	244.7	3024.8	2	74	1 164
Feb. 24 to Mar. 23.	4	5.3	7.0	87.7	3.96	5.28	69.25	378.1	501.5	6248.6	1	64	1 04
Mar. 23 to April 20.	4	7.0	7.0	72.0	5.05	5.05	51.84	409.8	409.8	4215.8	1	114	1 34
Dec. 2 to April 20.	20	5.6	7.0	77.5	4.62	5.77	63.86	269.9	337.3	3733.9	2	14	1 114

Lot 2.—12 Sheep; Special Food—Malted Barley "No. 1" (with Malt-dust).

1883—4.													
Dec. 2 to Dec. 30.	4	5.2	7.0	76.9	4.87	6.84	72.30	189.7	217.1	2715.4	2	121	2 104
Dec. 30 to Jan. 27.	4	5.2	7.0	76.7	4.46	6.02	66.03	251.1	339.4	5720.2	2	1	1 124
Jan. 27 to Feb. 24.	4	5.2	7.0	76.4	4.16	5.63	60.63	226.3	346.4	3733.0	2	04	1 10
Feb. 24 to Mar. 23.	4	5.2	7.0	86.7	3.93	5.31	65.71	400.5	404.8	5013.3	1	114	1 5
Mar. 23 to April 20.	4	5.0	7.0	72.0	5.05	5.05	51.84	409.8	409.8	4215.8	1	114	1 34

[illegible]

Lot 4.—12† Sheep; Special Food—Malted Barley “No. 2” (with Malt-dust).

1933-4.		1934-5.		1935-6.		1936-7.		1937-8.		1938-9.		1939-40.		1940-1.		1941-2.		1942-3.		1943-4.		1944-5.		1945-6.		1946-7.		1947-8.		1948-9.		1949-50.		1950-1.		1951-2.		1952-3.		1953-4.		1954-5.		1955-6.		1956-7.		1957-8.		1958-9.		1959-60.		1960-1.		1961-2.		1962-3.		1963-4.		1964-5.		1965-6.		1966-7.		1967-8.		1968-9.		1969-70.		1970-1.		1971-2.		1972-3.		1973-4.		1974-5.		1975-6.		1976-7.		1977-8.		1978-9.		1979-80.		1980-1.		1981-2.		1982-3.		1983-4.		1984-5.		1985-6.		1986-7.		1987-8.		1988-9.		1989-90.		1990-1.		1991-2.		1992-3.		1993-4.		1994-5.		1995-6.		1996-7.		1997-8.		1998-9.		1999-00.		2000-1.		2001-2.		2002-3.		2003-4.		2004-5.		2005-6.		2006-7.		2007-8.		2008-9.		2009-10.		2010-1.		2011-2.		2012-3.		2013-4.		2014-5.		2015-6.		2016-7.		2017-8.		2018-9.		2019-20.		2020-1.		2021-2.		2022-3.		2023-4.		2024-5.		2025-6.		2026-7.		2027-8.		2028-9.		2029-30.		2030-1.		2031-2.		2032-3.		2033-4.		2034-5.		2035-6.		2036-7.		2037-8.		2038-9.		2039-40.		2040-1.		2041-2.		2042-3.		2043-4.		2044-5.		2045-6.		2046-7.		2047-8.		2048-9.		2049-50.		2050-1.		2051-2.		2052-3.		2053-4.		2054-5.		2055-6.		2056-7.		2057-8.		2058-9.		2059-60.		2060-1.		2061-2.		2062-3.		2063-4.		2064-5.		2065-6.		2066-7.		2067-8.		2068-9.		2069-70.		2070-1.		2071-2.		2072-3.		2073-4.		2074-5.		2075-6.		2076-7.		2077-8.		2078-9.		2079-80.		2080-1.		2081-2.		2082-3.		2083-4.		2084-5.		2085-6.		2086-7.		2087-8.		2088-9.		2089-90.		2090-1.		2091-2.		2092-3.		2093-4.		2094-5.		2095-6.		2096-7.		2097-8.		2098-9.		2099-00.		2100-1.		2101-2.		2102-3.		2103-4.		2104-5.		2105-6.		2106-7.		2107-8.		2108-9.		2109-10.		2110-1.		2111-2.		2112-3.		2113-4.		2114-5.		2115-6.		2116-7.		2117-8.		2118-9.		2119-20.		2120-1.		2121-2.		2122-3.		2123-4.		2124-5.		2125-6.		2126-7.		2127-8.		2128-9.		2129-30.		2130-1.		2131-2.		2132-3.		2133-4.		2134-5.		2135-6.		2136-7.		2137-8.		2138-9.		2139-40.		2140-1.		2141-2.		2142-3.		2143-4.		2144-5.		2145-6.		2146-7.		2147-8.		2148-9.		2149-50.		2150-1.		2151-2.		2152-3.		2153-4.		2154-5.		2155-6.		2156-7.		2157-8.		2158-9.		2159-60.		2160-1.		2161-2.		2162-3.		2163-4.		2164-5.		2165-6.		2166-7.		2167-8.		2168-9.		2169-70.		2170-1.		2171-2.		2172-3.		2173-4.		2174-5.		2175-6.		2176-7.		2177-8.		2178-9.		2179-80.		2180-1.		2181-2.		2182-3.		2183-4.		2184-5.		2185-6.		2186-7.		2187-8.		2188-9.		2189-90.		2190-1.		2191-2.		2192-3.		2193-4.		2194-5.		2195-6.		2196-7.		2197-8.		2198-9.		2199-00.		2200-1.		2201-2.		2202-3.		2203-4.		2204-5.		2205-6.		2206-7.		2207-8.		2208-9.		2209-10.		2210-1.		2211-2.		2212-3.		2213-4.		2214-5.		2215-6.		2216-7.		2217-8.		2218-9.		2219-20.		2220-1.		2221-2.		2222-3.		2223-4.		2224-5.		2225-6.		2226-7.		2227-8.		2228-9.		2229-30.		2230-1.		2231-2.		2232-3.		2233-4.		2234-5.		2235-6.		2236-7.		2237-8.		2238-9.		2239-40.		2240-1.		2241-2.		2242-3.		2243-4.		2244-5.		2245-6.		2246-7.		2247-8.		2248-9.		2249-50.		2250-1.		2251-2.		2252-3.		2253-4.		2254-5.		2255-6.		2256-7.		2257-8.		2258-9.		2259-60.		2260-1.		2261-2.		2262-3.		2263-4.		2264-5.		2265-6.		2266-7.		2267-8.		2268-9.		2269-70.		2270-1.		2271-2.		2272-3.		2273-4.		2274-5.		2275-6.		2276-7.		2277-8.		2278-9.		2279-80.		2280-1.		2281-2.		2282-3.		2283-4.		2284-5.		2285-6.		2286-7.		2287-8.		2288-9.		2289-90.		2290-1.		2291-2.		2292-3.		2293-4.		2294-5.		2295-6.		2296-7.		2297-8.		2298-9.		2299-00.		2300-1.		2301-2.		2302-3.		2303-4.		2304-5.		2305-6.		2306-7.		2307-8.		2308-9.		2309-10.		2310-1.		2311-2.		2312-3.		2313-4.		2314-5.		2315-6.		2316-7.		2317-8.		2318-9.		2319-20.		2320-1.		2321-2.		2322-3.		2323-4.		2324-5.		2325-6.		2326-7.		2327-8.		2328-9.		2329-30.		2330-1.		2331-2.		2332-3.		2333-4.		2334-5.		2335-6.		2336-7.		2337-8.		2338-9.		2339-40.		2340-1.		2341-2.		2342-3.		2343-4.		2344-5.		2345-6.		2346-7.		2347-8.		2348-9.		2349-50.		2350-1.		2351-2.		2352-3.		2353-4.		2354-5.		2355-6.		2356-7.		2357-8.		2358-9.		2359-60.		2360-1.		2361-2.		2362-3.		2363-4.		2364-5.		2365-6.		2366-7.		2367-8.		2368-9.		2369-70.		2370-1.		2371-2.		2372-3.		2373-4.		2374-5.		2375-6.		2376-7.		2377-8.		2378-9.		2379-80.		2380-1.		2381-2.		2382-3.		2383-4.		2384-5.		2385-6.		2386-7.		2387-8.		2388-9.		2389-90.		2390-1.		2391-2.		2392-3.		2393-4.		2394-5.		2395-6.		2396-7.		2397-8.		2398-9.		2399-00.		2400-1.		2401-2.		2402-3.		2403-4.		2404-5.		2405-6.		2406-7.		2407-8.		2408-9.		2409-10.		2410-1.		2411-2.		2412-3.		2413-4.		2414-5.		2415-6.		2416-7.		2417-8.		2418-9.		2419-20.		2420-1.		2421-2.		2422-3.		2423-4.		2424-5.		2425-6.		2426-7.		2427-8.		2428-9.		2429-30.		2430-1.		2431-2.		2432-3.		2433-4.		2434-5.		2435-6.		2436-7.		2437-8.		2438-9.		2439-40.		2440-1.		2441-2.		2442-3.		2443-4.		2444-5.		2445-6.		2446-7.		2447-8.		2448-9.		2449-50.		2450-1.		2451-2.		2452-3.		2453-4.		2454-5.		2455-6.		2456-7.		2457-8.		2458-9.		2459-60.		2460-1.		2461-2.		2462-3.		2463-4.		2464-5.		2465-6.		2466-7.		2467-8.		2468-9.		2469-70.		2470-1.		2471-2.		2472-3.		2473-4.		2474-5.		2475-6.		2476-7.		2477-8.		2478-9.		2479-80.		2480-1.		2481-2.		2482-3.		2483-4.		2484-5.		2485-6.		2486-7.		2487-8.		2488-9.		2489-90.		2490-1.		2491-2.		2492-3.		2493-4.		2494-5.		2495-6.		2496-7.		2497-8.		2498-9.		2499-00.		2500-1.		2501-2.		2502-3.		2503-4.		2504-5.		2505-6.		2506-7.		2507-8.		2508-9.		2509-10.		2510-1.		2511-2.		2512-3.		2513-4.		2514-5.		2515-6.		2516-7.		2517-8.		2518-9.		2519-20.		2520-1.		2521-2.		2522-3.		2523-4.		2524-5.		2525-6.		2526-7.		2527-8.		2528-9.		2529-30.		2530-1.		2531-2.		2532-3.		2533-4.		2534-5.		2535-6.		2536-7.		2537-8.		2538-9.		2539-40.		2540-1.		2541-2.		2542-3.		2543-4.		2544-5.		2545-6.		2546-7.		2547-8.		2548-9.		2549-50.		2550-1.		2551-2.		2552-3.		2553-4.		2554-5.		2555-6.		2556-7.		2557-8.		2558-9.		2559-60.		2560-1.		2561-2.		2562-3.		2563-4.		2564-5.		2565-6.		2566-7.		2567-8.		2568-9.		2569-70.		2570-1.		2571-2.		2572-3.		2573-4.		2574-5.		2575-6.		2576-7.		2577-8.		2578-9.		2579-80.		2580-1.		2581-2.		2582-3.		2583-4.		2584-5.		2585-6.		2586-7.		2587-8.		2588-9.		2589-90.		2590-1.		2591-2.		2592-3.		2593-4.		2594-5.		2595-6.		2596-7.		2597-8.		2598-9.		2599-00.		2600-1.		2601-2.		2602-3.		2603-4.		2604-5.		2605-6.		2606-7.		2607-8.		2608-9.		2609-10.		2610-1.		2611-2.		2612-3.		2613-4.		2614-5.		2615-6.		2616-7.		2617-8.		2618-9.		2619-20.		2620-1.		2621-2.		2622-3.		2623-4.		2624-5.		2625-6.		2626-7.		2627-8.		2628-9.		2629-30.		2630-1.		2631-2.		2632-3.		2633-4.		2634-5.		2635-6.		2636-7.		2637-8.		2638-9.		2639-40.		2640-1.		2641-2.		2642-3.		2643-4.		2644-5.		2645-6.		2646-7.		2647-8.		2648-9.		2649-50.		2650-1.		2651-2.		2652-3.		2653-4.		2654-5.		2655-6.		2656-7.		2657-8.		2658-9.		2659-60.		2660-1.		2661-2.		2662-3.		2663-4.		2664-5.		2665-6.		2666-7.		2667-8.		2668-9.		2669-70.		2670-1.		2671-2.		2672-3.		2673-4.		2674-5.		2675-6.		2676-7.		2677-8.		2678-9.		2679-80.		2680-1.		2681-2.		2682-3.		2683-4.		2684-5.		2685-6.		2686-7.		2687-8.		2688-9.		2689-90.		2690-1.		2691-2.		2692-3.		2693-4.		2694-5.		2695-6.		2696-7.		2697-8.		2698-9.		2699-00.		2700-1.		2701-2.		2702-3.		2703-4.		2704-5.		2705-6.		2706-7.		2707-8.		2708-9.		2709-10.		2710-1.		2711-2.		2712-3.		2713-4.		2714-5.		2715-6.		2716-7.		2717-8.		2718-9.		2719-20.		2720-1.		2721-2.		2722-3.		2723-4.		2724-5.		2725-6.		2726-7.		2727-8.		2728-9.		2729-30.		2730-1.		2731-2.		2732-3.		2733-4.		2734-5.		2735-6.		2736-7.		2737-8.		2738-9.		2739-40.		2740-1.		2741-2.		2742-3.		2743-4.		2744-5.		2745-6.		2746-7.		2747-8.		2748-9.		2749-50.		2750-1.		2751-2.		2752-3.		2753-4.		2754-5.		2755-6.		2756-7.		2757-8.		2758-9.		2759-60.		2760-1.		2761-2.		2762-3.		2763-4.		2764-5.		2765-6.		2766-7.		2767-8.		2768-9.		2769-70.		2770-1.		2771-2.		2772-3.		2773-4.		2774-5.		2775-6.		2776-7.		2777-8.		2778-9.		2779-80.		2780-1.		2781-2.		2782-3.		2783-4.		2784-5.		2785-6.		2786-7.		2787-8.		2788-9.		2789-90.		2790-1.		2791-2.		2792-3.		2793-4.		2794-5.		2795-6.		2796-7.		2797-8.		2798-9.		2799-00.		2800-1.		2801-2.		2802-3.		2803-4.		2804-5.		2805-6.		2806-7.		2807-8.		2808-9.		2809-10.		2810-1.	
---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	---------	--	----------	--	---------	--

Lot 5.—12 Sheep; Special Food—Unmalted and Malted Barley "No. 2" (with Malt-dust).

[illegible]

• These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

+ No. 4 sheep of this lot (4) was killed February 3, and its consumption of food, and weights, are only taken into account in the results given for the first two months.

‡ Two thirds of these amounts represent unmalted barley, and the remaining one-third the amount of barley from which the malt and malt-dust given would be produced.

TABLE XLI.—EXPERIMENTS MADE WITH SHIMP, at BETHAMPTON, ILLINOIS.

Weights of the Carcasses, and some other Separated Parts.

Designation of Parts.	MONTH.												Month.
	No. 1.	No. 2.	No. 3.	No. 4.	No. 5.	No. 6.	No. 7.	No. 8.	No. 9.	No. 10.	No. 11.	No. 12.	
Lot 1.—Special Food—Unmalted Barley "No. 1."													
Caul fat	lbs. oss. 0 8	lbs. oss. 0 12	lbs. oss. 0 0	lbs. oss. 0 12	lbs. oss. 0 12	lbs. oss. 0 12	lbs. oss. 0 4	lbs. oss. 0 4	lbs. oss. 0 4	lbs. oss. 0 12	lbs. oss. 0 0	lbs. oss. 0 0	lbs. oss. 0 0
Intestinal fat	4 8	0 0	0 0	4 0	4 12	4 0	0 4	0 4	0 4	0 12	0 0	0 0	0 0
Skin and wool	18 0	17 0	17 0	16 0	14 0	17 0	16 0	23 0	18 0	16 0	17 0	17 10	17 10
Total	26 0	27 12	26 0	26 12	26 0	27 12	26 12	29 4	26 0	26 0	26 0	27 0	27 0
Other offal parts, loss by } fasting, evaporation, &c. }	41 0	48 4	47 0	38 4	42 8	42 8	39 4	46 12	38 0	43 12	38 8	41 0	41 0
Carcass (cold)	73 0	74 0	79 0	73 0	69 0	66 0	74 0	84 0	73 0	82 0	60 0	74 0	74 0
Live-weight (unfasted)	142 0	150 0	164 0	137 0	140 0	134 0	140 0	162 0	138 0	161 0	119 0	149 10	149 10
Lot 2.—Special Food—Malted Barley "No. 1" (with Malt-dust).													
Caul fat	lbs. oss. 7 0	lbs. oss. 3 12	lbs. oss. 4 12	lbs. oss. 6 0	lbs. oss. 4 12	lbs. oss. 6 4	lbs. oss. 0 12	lbs. oss. 0 0	lbs. oss. 0 0	lbs. oss. 0 0	lbs. oss. 4 4	lbs. oss. 0 0	lbs. oss. 0 0
Intestinal fat	3 12	2 8	3 12	4 4	5 0	4 4	0 0	0 0	0 0	0 12	0 0	0 0	0 0
Skin and wool	16 0	20 0	17 0	16 0	21 0	22 0	16 0	19 0	14 0	16 0	16 0	17 10	17 10
Total	26 12	26 4	26 8	26 4	30 12	32 8	26 12	29 0	24 0	26 12	23 8	27 0	27 0
Other offal parts, loss by } fasting, evaporation, &c. }	41 0	48 4	47 0	38 4	42 8	42 8	39 4	46 12	38 0	43 12	38 8	41 0	41 0
Carcass (cold)	73 0	74 0	79 0	73 0	69 0	66 0	74 0	84 0	73 0	82 0	60 0	74 0	74 0
Live-weight (unfasted)	142 0	150 0	164 0	137 0	140 0	134 0	140 0	162 0	138 0	161 0	119 0	149 10	149 10

Total	28 12	25 4	18 0	18 0	3 12	21 0	3 4	4 0	3 8	15 0	21 0	20 0	16 0	16 0
Other offal parts, loss by fasting, evaporation, &c. }	47 4	51 12	37 0	39 4	50 0	36 4	26 8	23 12	24 4	33 8	27 4	23 4	26 0	26 0
Carcass (cold)	87 0	81 0	84 0	81 0	85 0	71 0	64 0	63 0	76 0	76 0	71 0	67 0	75 8	75 8
Live-weight (unfasted)	163 0	158 0	150 0	147 0	164 0	130 0	117 0	120 0	140 0	152 0	142 0	126 0	142 6	142 6

Lot 4.—Special Food—Malted Barley “No. 2” (with Malt-dust).

Caul fat	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
Intestinal fat	4 12	5 4	4 4	3 8	5 4	3 8	4 4	4 8	4 0	4 8	3 12	3 8	4 8	4 8
Skin and wool	21 0	20 0	16 0	•	20 0	15 0	20 0	19 0	18 0	18 0	19 0	19 0	19 0	18 10
Total	29 12	30 12	23 4		28 12	23 0	28 12	27 0	25 12	28 0	25 12	26 12	26 12	26 15
Other offal parts, loss by fasting, evaporation, &c. }	47 4	53 4	42 12		47 4	40 4	44 0	44 0	40 4	28 0	45 4	36 4	42 9	42 9
Carcass (cold)	73 0	80 0	68 0		11 0	66 0	71 0	69 0	74 0	70 0	75 0	67 0	72 2	72 2
Live-weight (unfasted)	150 0	164 0	134 0		157 0	128 0	144 0	140 0	140 0	126 0	146 0	130 0	141 1	141 1

Lot 5.—Special Food—Barley “No. 2,” } Unmalted and } Malted (with Malt-dust).

Caul fat	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.	lbs. ozs.
Intestinal fat	6 4	6 4	4 8	5 4	4 12	6 4	5 0	5 4	5 8	5 8	4 8	4 8	4 8	5 4
Skin and wool	22 0	15 0	16 0	21 0	15 0	20 0	20 0	20 0	17 0	21 0	18 0	18 0	18 0	18 9
Total	34 8	25 0	24 4	31 4	23 4	30 8	29 8	29 4	27 12	31 12	26 12	25 12	28 5	28 5
Other offal parts, loss by fasting, evaporation, &c. }	54 8	34 12	38 12	38 12	50 12	39 8	37 8	37 12	36 4	35 4	31 4	37 4	39 5	39 5
Carcass (cold)	97 0	66 0	67 0	73 0	66 0	75 0	69 0	79 0	70 0	66 0	51 0	69 0	71 2	71 2
Live-weight (unfasted)	166 0	126 0	130 0	142 0	140 0	145 0	136 0	146 0	134 0	133 0	116 0	132 0	138 13	138 13

* No. 4 sheep of Lot 4 was killed on February 3; it had been unwell about three weeks.

TABLE XLII.—EXPERIMENTS MADE WITH SHEEP, at ROTHAMSTED, HERTS.
Per-centages of the Carcasses, and some other separated Parts, in the Live-weights (unfasted).

[illegible]

Intestinal fat	3.81	3.64	3.83	2.38	2.58	2.69	3.40	3.22	3.29	14.04	12.70	12.67
Skin and wool	3.07	3.21	3.50	3.57	2.28	2.60	3.42	3.92	13.81	14.04	12.70	12.67
	11.66	10.13	12.00	12.25	12.80	12.31	15.38	14.17				
Total	17.64	16.98	19.33	18.20	17.68	17.60	22.65	17.33	21.38	19.19	18.46	18.76
Other offal parts, lost by fasting, evaporation, &c.	28.99	32.76	21.67	26.70	30.49	27.89	22.68	28.39	28.62	30.81	28.37	28.17
Carcass (cold)	53.37	51.27	56.00	55.10	51.83	54.61	54.70	52.50	50.00	50.00	53.17	53.07
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

Lot 4.—Special Food—Malted Barley "No. 2" (with Malt-dust).

Caul fat	3.17	3.21	3.17	*	3.24	2.74	2.95	3.21	2.86	3.57	2.57	3.04
Intestinal fat	2.66	3.35	2.24		2.23	2.73	3.13	2.50	2.69	2.08	3.27	2.85
Skin and wool	14.00	12.19	11.94		12.74	11.72	13.89	13.57	12.87	14.29	13.01	13.16
Total	19.83	18.75	17.35		18.31	17.19	19.97	19.28	18.42	22.22	17.64	19.05
Other offal parts, lost by fasting, evaporation, &c.	31.50	32.47	31.90		30.10	31.25	30.72	31.43	28.72	23.23	30.99	29.92
Carcass (cold)	48.67	48.78	50.75		51.59	51.56	49.31	49.29	52.86	55.56	51.37	51.03
	100.00	100.00	100.00		100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

Lot 5.—Special Food—Barley "No. 2" } Unmalted and } Malted (with Malt-dust).

Caul fat	3.36	4.96	3.46	3.70	3.39	4.31	3.67	3.59	4.10	4.14	3.88	3.83
Intestinal fat	3.36	3.18	2.88	3.52	2.50	2.93	3.31	2.74	3.92	3.95	3.66	3.20
Skin and wool	11.83	11.90	12.31	14.79	10.72	13.80	14.71	13.70	12.69	15.79	15.52	13.45
Total	18.56	20.04	18.65	22.01	16.61	21.04	21.69	20.03	20.71	23.88	23.06	20.48
Other offal parts, lost by fasting, evaporation, &c.	29.30	27.58	29.81	27.29	36.25	27.24	27.57	25.86	27.06	28.50	26.94	28.30
Carcass (cold)	52.15	52.38	51.54	50.70	47.14	51.72	50.74	54.11	52.24	49.62	50.00	51.22
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

* No. 4 sheep of Lot 4 was killed February 3; it had been unwell about 3 weeks.

TABLE XLIII.—EXPERIMENTS MADE WITH PIGS, AT ROTHAMSTED, HERTS.
Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight. 1st Period of 2 Weeks; Dec. 24, 1863, to Jan. 7, 1864.

	Food consumed.				Pigs.	Weights and Increase in Live-weight.			
	Total in 2 Weeks.	Per Head per Day.	Per 100 lbs. Live-weight per Week.	To produce 100 lbs. Increase.		Weights.		Increase in Live-weight.	
						Dec. 24, 1863.	Jan. 7, 1864.	Total in 2 Weeks.	Per Head per Week.
Lot 1.—8 Pigs; Food—Pea-meal, and Unmalted Barley "No. 2."									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
					1	136	182	24	12'0
					2	160	198	38	19'0
					3	133	165	33	16'5
					4	134	172	38	19'0
					5	135	161	26	13'0
					6	132	160	28	13'0
					7	122	160	38	19'0
					8	122	160	38	19'0
					Totals . .	1086	1360	274	—
					Averages .	136	170	34	17'1
									11'20
Lot 2.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 1" (with Malt-dust).									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
					1	172	189	17	8'5
					2	144	168	24	12'0
					3	149	164	22	11'0
					4	133	168	35	17'5
					5	130	161	31	15'5
					6	134	140	16	8'0
					7	119	143	23	11'5
					8	123	140	17	8'5
					Totals . .	1087	1272	185	—
					Averages .	136	159	23	11'6
									7'94
Lot 3.—8 Pigs; Food—Pea-meal, Unmalted and Malted Barley "No. 1" (with Malt-dust).									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
					1	147	174	27	13'5
					2	150	178	28	14'0
					3	150	168	18	9'0
					4	144	175	31	15'5
					5	149	154	25	13'5
					6	130	151	21	14'0
									8'43

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted, <i>ad lib.</i>	112	1'0	4'08	50'2	175	147	28	12'9	9'31
Barley, unmalted, <i>ad lib.</i>	800	7'1	33'40	338'7	170	171	31	14'0	14'0
					166	154	38	15'0	14'0
					133	161	28	14'0	14'0
					112	140	28	14'0	14'0
Totals . .					1309		173		—
Averages .					164		27½		9'31

Lot 5.—8 Pigs : Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, malted, with its dust, } <i>ad lib.</i>	112	1'0	4'67	64'7	170	14	30	15'0	7'21
	938*	8'4*	39'10*	543'3*	182	25	17	8'5	19'0
					164	38	17	8'5	19'0
					140	17	17	8'5	19'0
					147			8'5	19'0
Totals . .					1113		173		—
Averages .					139		21½		10'8

Lot 6.—8 Pigs ; Food—Pea-meal, and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2" (with Malt-dust *ad lib.*).

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted	112	1'0	4'62	48'1	182	192	35	17'5	9'61
Barley, unmalted	700	6'3	28'87	300'4	163	163	35	17'5	17'5
Barley, malted, with its dust . .	173*	1'6*	7'22*	75'1*	134	134	0	0'0	17'0
					170	34	33	16'5	14'0
					161	28	28	14'0	14'0
					112				—
Totals . .					1066		233		—
Averages .					137		29½		14'6

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

TABLE XLIV.—EXPERIMENTS MADE WITH PIGS, AT ROTHAMSTED, HERTS.
Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight. 2nd Period of 2 Weeks; Jan. 7 to Jan. 21, 1904.

Food consumed.				Weights and Increase in Live-weight.			
Total in 2 Weeks.	Per Head per Day.	Per 100 lbs. Live-weight per Week.	To produce 1 lb. live- weight increase.	Place.	Weights.	Increase in Live-weight	Per 100 lbs. Live-weight per Week.
					Jan. 7, 1904.	Jan. 21, 1904.	Total in 2 Weeks per Head
Lot 1.—8 Pigs: Food—Pea-meal, and Unmalted Barley "No. 1."							
Pea-meal	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.
Barley, unmalted, <i>ad lib.</i>	112	1'0	3'73	1	104	10'0	10'0
				2	106	10'0	10'0
				3	106	10'0	10'0
				4	104	10'0	10'0
				5	104	10'0	10'0
				6	104	10'0	10'0
				7	104	10'0	10'0
				8	104	10'0	10'0
				Total . . .	840	80'0	80'0
				Average . .	105	10'0	10'0
Lot 2.—8 Pigs: Food—Pea-meal, and Malted Barley "No. 1" (with Malt-dust).							
Pea-meal	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.
Barley, malted, with its dust, } <i>ad lib.</i>	112	1'0	4'08	1	104	10'0	10'0
				2	106	10'0	10'0
				3	106	10'0	10'0
				4	104	10'0	10'0
				5	104	10'0	10'0
				6	104	10'0	10'0
				7	104	10'0	10'0
				8	104	10'0	10'0
				Total . . .	840	80'0	80'0
				Average . .	105	10'0	10'0
Lot 3.—7† Pigs: Food—Pea-meal, Unmalted and Malted Barley "No. 1" (with Malt-dust).							
Pea-meal	lbs.	lbs.	lbs.	Nos.	lbs.	lbs.	lbs.
Barley, unmalted, <i>ad lib.</i>	96	1'0	3'91	1	104	10'0	10'0
				2	106	10'0	10'0
				3	106	10'0	10'0
				4	104	10'0	10'0
				5	104	10'0	10'0
				6	104	10'0	10'0
				7	104	10'0	10'0
				Total . . .	728	70'0	70'0
				Average . .	104	10'0	10'0

Pea-meal	lbs.	lbs.	lbs.	lbs.	1	172	208	30	19'0	9'47
Barley, unmalted, ad lib.	112	1'0	3'67	40'9	2	170	208	36	17'8	
	688	8'4	32'43	340'3	3	171	206	46	16'0	
					4	166	198	30	15'0	
					5	164	184	33	16'5	
					6	161	194	35	17'5	
					7	140	175	35	17'5	
Totals					8	1309	1583	274	—	
Averages						164	198	344	17'1	9'47

Lot 5.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	9'72
Barley, malted, with its dust, } ad lib.	112	1'0	3'93	40'4	1	170	210	40	20'0	
	1079*	9'6*	37'65*	387'1*	2	168	204	48	21'0	
					3	172	214	39	19'5	
					4	167	178	31	18'5	
					5	163	183	32	18'0	
					6	164	186	33	16'0	
					7	140	168	33	14'0	
					8	147	180	33	16'5	
Totals						1395	1568	277	—	—
Averages						161	196	34½	17'3	9'72

Lot 6.—7½ Pigs; Food—Pea-meal, and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2" (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	8'67
Barley, unmalted	98	1'0	3'74	43'2	1	168	200	44	20'0	
Barley, malted, with its dust	738½	7'5	28'21	325'2	2	162	198	59	14'5	
	183*	1'9	7'06*	81'3	3	163	194	31	15'5	
					4	163	—	—	—	
					5†	170	200	30	15'0	
					6	161	184	27	13'5	
					7	140	166	26	13'0	
Totals					8	1185	1428	227	—	—
Averages						171	203	32½	16'2	8'67

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

† No. 3 pig of this lot (3) being unwell, was killed Jan. 14, and its proportion of food and weight are excluded in the calculated results given above.

‡ No. 5 pig of this lot (6) being unwell, was killed Jan. 7, so that the results relate to 7 pigs only.

TABLE XLV.—EXPERIMENTS MADE WITH PIGS, AT ROTHAMSTED, HERTS.
Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight. 3rd Period of 2 Weeks; Jan. 21 to Feb. 4, 1884.

	Food consumed.			Pigs.	Weights and Increase in Live-weight.			
	Total in 2 Weeks.	Per Head per Day.	Per 100 lbs. Live-weight per Week.		Weights.		Increase in Live-weight.	
					Jan. 21, 1884.	Feb. 4, 1884.	Total in 2 Weeks.	Per Head per week.
Lot 1.—8 Pigs; Food—Pea-meal, and Unmalted Barley "No. 1."								
	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
				1	218	242	24	12.0
				2	226	269	33	16.5
		lbs.		3	196	226	28	14.0
				4	212	224	12	11.0
		51.1		5	189	218	29	14.5
				6	228	236	8	10.5
		461.2		7	189	210	21	10.5
				8	192	224	32	16.0
				Totals . .	1,642	1,861	219	—
				Averages .	205	233	27½	13.7
Pea-meal	112	1.0	3.20					6.25
Barley, unmalted, ad lib. . . .	1,010	9.0	28.83					—
Lot 2.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 1" (with Malt-dust).								
	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
				1	224	250	26	13.0
				2	198	224	26	13.0
				3	183	218	19	9.5
		65.9		4	206	231	25	12.5
				5	184	200	16	8.0
				6	174	196	22	12.0
		572.9*		7	169	186	16	9.0
				8	164	180	16	8.0
				Totals . .	1,511	1,681	170	—
				Averages .	189	210	21½	10.6
Pea-meal	112	1.0	3.51					5.33
Barley, malted, with its dust, ad lib.	974*	8.7*	30.51*					—
Lot 3.—7½ Pigs; Food—Pea-meal, Unmalted and Malted Barley "No. 1" (with Malt-dust).								
	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
				1	208	221	13	11.5
				2	210	238	28	14.0
				3½	—	—	—	—
				4	212	259	36	13.0
		3.35		5	184	210	26	13.0
				6	196	224	28	14.0
		26.64		7	186	224	38	14.0
Pea-meal	94	1.0	3.35					6.99
Barley, unmalted, ad lib. . . .	779	7.9	26.64					—

Pea-meal	lbs.	lbs.	lbs.	lbs.	1	210	231	21	11.5	6.00
Barley, unmalted, <i>ad lib.</i>	113	1.0	3.48	lbs.	2	208	231	25	11.5	
	890	7.9	26.41	lbs.	3	206	245	37	18.0	
					4	206	231	35	12.5	
					5	196	217	19	9.5	
					6	184	203	19	9.5	
					7	184	203	23	16.0	14.0
					8	175	203	23	14.0	
Totals . . .						1,563	1,767	204	—	—
Averages .						196	223	25.4	12.8	6.00

Lot 5.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	6.01
Barley, malted, with its dust, } <i>ad lib.</i>	113	1.0	3.37	lbs.	1	210	224	24	12.0	
	1,137*	10.1*	34.18*	lbs.	2	214	246	24	12.0	
					3	176	215	31	15.5	
					4	183	192	14	7.0	
					5	193	220	27	13.5	
					6	196	224	28	14.0	13.0
					7	168	194	26	13.0	
					8	180	206	26	13.0	—
Totals . . .						1,563	1,763	200	—	—
Averages .						196	220	25	13.5	6.01

Lot 6.—7½ Pigs; Food—Pea-meal and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2" (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.	6.30
Barley, unmalted	98	1.0	3.23	lbs.	1	200	224	24	17.0	
Barley, malted, with its dust	739.41	7.5	24.36	lbs.	2	200	212	30	10.0	
	169*	1.7*	6.11*	lbs.	3	194	224	30	15.0	
					4	194	224	30	15.0	
					5†	200	222	22	11.0	
					6	198	212	24	12.0	14.0
					7	166	194	26	14.0	
					8	166	194	26	14.0	—
Totals . . .						1,422	1,610	188	—	—
Averages .						203	220	26.5	13.4	6.30

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

† No. 3 pig of this lot (3) having been killed the results relate to 7 pigs only.

‡ No. 5 pig of this lot (6) having been killed the results relate to 7 pigs only.

TABLE XLVI.—EXPERIMENTS MADE WITH PIGS, at ROTHAMSTED, HERTS.
Detailed Record of the Food consumed, of the Weights, and of the Increase in Live-weight, 4th Period of 2 Weeks; Feb. 4 to Feb. 18, 1864.

	Food consumed.				Pigs.	Weights, and Increase in Live-weight.			
	Total in 2 Weeks.	Per Head per Day.	Per 100 lbs. Live-weight per Week.	To produce 100 lbs. Increase.		Weights.		Increase in Live-weight.	
						Feb. 4, 1864.	Feb. 18, 1864.	Total in 2 Weeks.	Per Head Live-weight per Week.

Lot 1.—8 Pigs; Food—Pea-meal, and Unmalted Barley "No. 2."									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
					1	242	267	25	12.5
					2	269	304	35	17.5
					3	226	242	16	8.0
					4	218	244	26	14.0
					5	236	270	32	13.0
					6	238	270	32	16.0
					7	210	228	18	9.0
					8	224	252	28	14.0
					Totals . .	1861	2009	208	—
					Averages .	233	250	26	13.0
Pea-meal	112	1.0	3.85	53.8					5.29
Barley, unmalted, <i>ad lib.</i>	966	8.6	24.58	464.4					—

Lot 2.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 1" (with Malt-dust).									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
					1	250	282	32	6.0
					2	224	252	28	14.0
					3	212	231	19	8.5
					4	201	247	46	9.5
					5	180	210	30	5.0
					6	186	216	30	8.0
					7	166	193	27	8.0
					8	160	188	28	4.5
					Totals . .	1661	1869	128	—
					Averages .	210	226	16	8.0
Pea-meal	112	1.0	3.21	87.5					3.67
Barley, malted, with its dust, <i>ad lib.</i>	838	7.4	22.86	631.0					—

Lot 3.—7 Pigs; Food—Pea-meal, Unmalted and Malted Barley "No. 1" (with Malt-dust).									
	lbs.	lbs.	lbs.	lbs.	No.	lbs.	lbs.	lbs.	lbs.
					1	231	250	19	14.0
					2	229	260	31	11.0
					3	226	259	33	10.5
					4	226	242	16	8.0
					5	210	243	33	12.0
					6	224	244	20	8.0
					7	224	244	20	8.0
					Averages .	224	244	20	8.0
Pea-meal	112	1.0	3.16	54.4					5.47
Barley, unmalted, <i>ad lib.</i>	800	8.2	24.33	444.4					—

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted, ad lib.	112	1'0	3'96	54'6	3	945	378	25	12'5
	929	8'2	34'40	499'8	8	231	256	25	11'5
					7	236	250	24	13'0
					8	233	251	28	14'0
Totals						1787	1992	205	—
Averages						223	249	25½	12'8
									5'42

Lot 5.—8 Pigs; Food—Pea-meal, and Malted Barley No. 2 (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, malted, with its dust, ad lib.	112	1'0	3'00	54'4	1	234	266	32	16'0
	1091*	2'7*	29'25*	589'8*	2	248	273	25	12'5
					3	245	272	25	14'0
					4	250	272	25	11'5
					5	254	269	26	13'0
					6	254	260	25	11'5
					7	194	217	23	12'5
					8	205	231	25	—
Totals						1763	1969	206	—
Averages						220	246	26	12'2
									5'42

Lot 6.—6† Pigs; Food—Pea-meal, and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2 (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted	84	1'0	2'96	57'9	1	282	315	31	15'5
Barley, malted, with its dust	565	6'7	19'93	390'4	2†	(269)	232	20	10'0
					3	212	245	21	10'5
					4	224	253	30	15'0
					5	222	253	25	10'0
					6	112	112	23	11'5
					8	194	217	25	—
Totals						1348	1493	145	—
Averages						225	249	24	12'1
									5'10

* These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

† No. 3 pig of this lot (3) having been killed the result relates to seven pigs only.

‡ No. 2 pig of this lot (6) having been killed on February 6th, and No. 5 pig on January 7th, the results relate to six pigs only.

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted, ad lib.	112	1'0	2'70	67'1	260	276	288	292	14'0
	9764	8'7	23'28	564'7	250	256	260	266	9'0
					251	251	250	250	9'5
					251	251	250	250	15'0
					251	251	250	250	10'5
Totals	1992				2159			157	—
Averages	240				270			20½	10'4
									4'08

Lot 5.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, malted, with its dust, } ad lib.	112	1'0	2'74	73'2	273	273	273	273	10'5
	1049*	9'4*	25'64*	688'7*	217	217	217	217	12'5
					231	231	231	231	6'5
					231	231	231	231	10'0
					231	231	231	231	10'0
					231	231	231	231	5'5
					231	231	231	231	10'5
Totals	1999				2129			153	—
Averages	246				265			19½	9'6
									3'74

Lot 6.—6½ Pigs; Food—Pea-meal, and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2" (with Malt-dust).

Pea-meal	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Barley, unmalted	84	1'0	2'68	56'8	232	232	232	232	17'5
Barley, malted, with its dust	572	6'8	18'25	368'5	232	232	232	232	9'0
	145*	1'7*	4'53*	90'9*	217	217	217	217	10'5
					217	217	217	217	10'5
					217	217	217	217	10'5
					217	217	217	217	10'5
					217	217	217	217	10'5
Totals	1483				1483			148	—
Averages	249				274			24½	12'3
									4'72

* These figures represent not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

† No. 2 pig of this lot (3) having been killed, the results relate to 7 pigs only.

‡ No. 2 pig of this lot (5) having been killed on Feb. 5 and No. 2 pig on Jan. 7, the results relate to 6 pigs only.

TABLE XLVIII.—EXPERIMENTS made with PIGS, at ROTHAMSTED, Herts.
Summary of the Weights, and of the Increase in Live-weight. Total Period; December 24, 1863, to March 3, 1864.

Pigs.	Weights at Commencement Dec. 24, 1863.		Increase (or Loss) in Weight.					Weights at Conclusion March 3, 1864.		Increase in Live-weight.	
	In 2 Weeks; Dec. 24, 1863, to Jan. 7, 1864.	In 2 Weeks; Jan. 7, 1864, to Jan. 21, 1864.	In 2 Weeks; Jan. 21, 1864, to Feb. 4, 1864.	In 2 Weeks; Feb. 4, 1864, to Feb. 18, 1864.	In 2 Weeks; Feb. 18, 1864, to March 3, 1864.				Total in 10 Weeks.	Per Head per Week.	Per 100 lbs. Live-weight per Week.

Lot 1.—8 Pigs: Food—Pea-meal and Unmalted Barley "No. 1."											
Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.
1	156	24	36	33	30	29	28	129	12	14*4	5 12½
2	160	38	33	35	32	30	29	137	13	17*6	7 14
3	133	33	32	16	22	22	22	131	13	17*6	7 14
4	134	38	40	23	28	25	25	153	15	4*8	7 4½
5	135	29	28	29	25	25	27	135	13	8*0	6 10½
6	122	39	47	32	32	27	25	175	17	8*0	8 5½
7	122	38	21	13	13	13	13	119	11	14*4	6 9
8	122	38	32	28	28	28	28	138	13	12*8	7 13½
Totals . . .	1056	274	292	219	208	193	193	2262	117½	—	—
Averages . . .	132½	34½	36½	27½	26	24½	24½	282½	147	14 11*2	7 0½

Lot 2.—8 Pigs Food—Pea-meal, and Malted Barley "No. 1" (with Malt-dust).											
Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.
1	172	35	35	32	38	38	38	290	118	11 12*8	5 1½
2	142	34	32	35	35	35	35	278	134	6*4	6 5½
3	142	32	32	19	16	16	16	245	100	0*0	5 3½
4	133	35	32	22	16	16	16	245	100	1*2	5 11½
5	130	31	23	16	14	14	14	224	94	6*4	5 16
6	124	16	24	18	12	12	12	228	104	6*4	5 14½
7	119	22	18	16	12	12	12	214	95	8*0	5 11½
8	122	17	16	9	7	7	7	196	73	4*8	4 9½
Totals . . .	1067	185	239	170	193	193	193	1912	825	—	—
Averages . . .	133½	23½	29½	21½	24	24	24	239	102½	10 5*0	5 8

Lot 3.—8* Pigs Food—Pea-meal, Unmalted and Malted Barley "No. 1" (with Malt-dust).											
Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.	lbs. ozs.
1	147	27	34	23	28	28	28	180	133	13 4*8	6 2½
2	150	28	28	28	22	22	22	207	137	13 11*2	6 4½
3*	144	21	27	26	21	21	21	183	136	13 12*8	6 7½
4	144	21	27	26	21	21	21	183	136	13 12*8	6 7½
5	144	21	27	26	21	21	21	183	136	13 12*8	6 7½
6	130	28	34	25	32	32	32	204	160	16 0*0	7 13½
7	134	24	34	28	32	32	32	204	160	16 0*0	7 13½
8	104	18	28	21	21	21	21	153	103	10 4*8	5 11½

[illegible]

Lot 5.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

[illegible]

Lot 6.—8† Pigs; Food—Pea-meal, and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2" (with Malt-dust) *ad lib.*

No.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. con.	lbs. con.
1	106	38	34	31	350	138	18	3.2	7	0.4
2	157	35	49	30	250	117	11	11.2	6	1.1
3	133	30	29	30	250	142	14	5.2	7	1.1
4	188	35	31	25	270	142	14	5.2	7	1.1
5	134	31	31	31	250	142	14	5.2	7	1.1
6	136	34	32	30	250	142	14	5.2	7	1.1
7	126	32	34	28	250	142	14	5.2	7	1.1
8	112	28	26	23	250	142	14	5.2	7	1.1
Totals	1095	253	227	145	1641	836	13	14.9	6	1.34
Averages	137	29.4	28.4	24.4	205.1	104.5	1.6	1.86	0.75	0.17

No. 3 slug of this lot (3) was killed Jan. 14, and its weights are only taken into calculation in the total and average results given for the first fortnightly period.

+ No. 5 pig of this lot (6) was killed Jan. 7, and No. 2 pig on Feb. 5, and the weights of the former are only taken into calculation during the first, and those of the latter during the first three of the fortnightly periods.

TABLE XLIX.—EXPERIMENTS MADE WITH PIGS, at ROTHEAMSTED, HERTS.
Summary of Food consumed, and Increase yielded, during each separate fortnight, and during the entire period of Ten Weeks.

Periods.		Food consumed.								Increase in Weight.	
Dates.	Number of Weeks.	Per Head per Week.				Per 100 lbs. Live-weight per Week.				Per Head per Week.	Per 100 lbs. Live-weight per Week.
		Pea Meal.	Unmalted Barley.	Malted Barley (with its dust).		Pea Meal.	Unmalted Barley.	Malted Barley (with its dust).			
Lot 1.—8 Pigs; Food—Pea-meal, and Unmalted Barley "No. 1."											
1893-4.											
Dec. 24 to Jan. 7	2	7.0	56.2	..	4.68	36.79	40.9	328.5	..
Jan. 7 to Jan. 21	2	7.0	62.4	..	3.73	33.31	39.7	354.6	..
Jan. 21 to Feb. 4	2	7.0	63.1	..	3.20	28.83	61.1	461.2	..
Feb. 4 to Feb. 18	2	7.0	60.3	..	2.85	24.58	53.8	484.4	..
Feb. 18 to Mar. 3	2	7.0	64.1	..	2.69	23.69	58.0	531.6	..
Dec. 24 to Mar. 3	10	7.0	61.3	..	3.35	29.28	47.6	416.8	..
Lot 2.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 1" (with Malt-dust).											
1893-4.											
Dec. 24 to Jan. 7	2	7.0	..	56.4	4.75	37.60	60.5	..	479.5
Jan. 7 to Jan. 21	2	7.0	..	56.3	4.02	37.52	48.9	..	438.9
Jan. 21 to Feb. 4	2	7.0	..	50.9	3.51	30.51	68.9	..	572.8
Feb. 4 to Feb. 18	2	7.0	..	52.1	3.21	23.83	87.5	..	551.0
Feb. 18 to Mar. 3	2	7.0	..	40.2	3.01	17.28	108.7	..	624.1
Dec. 24 to Mar. 3	10	7.0	..	54.8	3.70	29.22	67.9	..	531.0
Lot 3.—8 Pigs; Food—Pea-meal, Unmalted, and Malted Barley "No. 1" (with Malt-dust).											
1893-4.											
Dec. 24 to Jan. 7	2	7.0	34.2	21.0	4.65	22.76	..	13.95	65.2	269.7	165.4
Jan. 7 to Jan. 21	2	7.0	46.8	11.0	3.01	28.18	..	9.07	40.1	241.4	146.7

	Dec. 24 to Jan. 7	Jan. 7 to Jan. 21	Jan. 21 to Feb. 4	Feb. 4 to Feb. 18	Feb. 18 to Mar. 3	Dec. 24 to Mar. 3
lbs.	7-0	60-0	58-6	7-0	61-0	56-6
lbs.
lbs.	.	4-68	4-9	3-52	2-96	3-45
lbs.	33-40	32-43	26-41	24-40	23-52	27-90
lbs.	50-2	40-9	54-9	54-6	67-1	52-2
lbs.	389-7	342-3	436-3	449-8	534-7	431-9
tons.
tons.	13 15	17 2	12 12	12 13	10 7	13 6½
cows	9 5	9 7½	6 0¼	5 6¾	4 0¼	6 9¼

Lot 5.—8 Pigs; Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

[illegible]

Lot 6.—8½ Pigs; Food—Pea-meal, and Mixture of four-fifths Unmalted, and one-fifth Malted Barley “No. 2” (with Malt-dust).

1893-4.		lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs. ozs.
Dec. 24 to Jan. 7	2	7-0	43-8	21-9	28-57	27-02	30-4	275-1	14	9
Jan. 7 to Jan. 21	2	7-0	43-7	21-9	28-57	27-02	30-4	275-1	16	34
Jan. 21 to Feb. 4	2	7-0	43-8	21-9	28-57	27-02	30-4	275-1	16	34
Feb. 4 to Feb. 18	2	7-0	43-8	21-9	28-57	27-02	30-4	275-1	16	34
Feb. 18 to Mar. 3	2	7-0	43-8	21-9	28-57	27-02	30-4	275-1	16	34
Dec. 24 to Mar. 3	10	7-0	48-8	112-2	23-95	25-98	350-4	287-5	13	15

These figures represent, not the actual weights of malt and malt-dust, but the weight of the barley from which they would be produced.

These figures represent, not the actual weights of maize and manure, but the weight of the barley from which they were produced. † No. 3 pig of this lot (3) was killed Jan. 14, and its consumption of food and weights are only taken into account in the results given for the first fortnightly period. ‡ No. 5 pig of this lot (6) was killed Jan. 7, and No. 3 pig on Feb. 5, and of the former the consumption of food and weights are only taken into account in the results given for the first, and of the latter only in those given for the first three of the fortnightly periods.

TABLE L.—EXPERIMENTS MADE WITH PIGS, at ROTHAMSTED, HERTS.
Weights of the Carcasses, &c.

Designation of Parts.	PIGS.							
	No. 1.	No. 2.	No. 3.	No. 4.	No. 5.	No. 6.	No. 7.	Means.
Lot 1.—Food—Pea-meal, and Unmalted Barley "No. 1."								
Carcass (cold)	lbs. 227	lbs. 271½	lbs. 204	lbs. 224½	lbs. 208	lbs. 238	lbs. 199	lbs. 224½
Offal parts, loss by fasting, evaporation, &c.	60	64½	60	62½	62	59	42	58½
Live-weight (unfasted)	287	336	264	287	270	297	241	283½
Lot 2.—Food—Pea-Meal, and Malted Barley "No. 1" (with Malt-dust).								
Carcass (cold)	lbs. 227½	lbs. 217½	lbs. 188	lbs. 199	lbs. 173½	lbs. 180	lbs. 160½	lbs. 187½
Offal parts, loss by fasting, evaporation, &c.	62½	60½	54	41	50½	48	53½	51½
Live-weight (unfasted)	290	278	242	240	224	228	214	239
Lot 3.—Food—Pea-meal, and Unmalted and Malted Barley "No. 1" (with Malt-dust).								
Carcass (cold)	lbs. 223½	lbs. 227	lbs. ..	lbs. 231½	lbs. 232½	lbs. 231	lbs. 229½	lbs. 231½

Carcass (cold)	lbs.	213½	lbs.	255½	lbs.	236½	lbs.	209½	lbs.	204	lbs.	196	lbs.	220½	lbs.	192	lbs.	212½
Offal parts, loss by fasting, evaporation, &c.	lbs.	59½	lbs.	56½	lbs.	69½	lbs.	53½	lbs.	55	lbs.	49	lbs.	59½	lbs.	60	lbs.	57½
Live-weight (unfasted)	lbs.	273	lbs.	282	lbs.	306	lbs.	262	lbs.	259	lbs.	245	lbs.	280	lbs.	252	lbs.	269½

Lot 5.—Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

Carcass (cold)	lbs.	227½	lbs.	229½	lbs.	235	lbs.	180½	lbs.	203½	lbs.	210	lbs.	177½	lbs.	189	lbs.	206½
Offal parts, loss by fasting, evaporation, &c.	lbs.	56½	lbs.	64½	lbs.	63	lbs.	53½	lbs.	56½	lbs.	60	lbs.	50½	lbs.	63	lbs.	58½
Live-weight (unfasted)	lbs.	284	lbs.	294	lbs.	298	lbs.	234	lbs.	262	lbs.	270	lbs.	228	lbs.	252	lbs.	265½

Lot 6.—Food—Pea-meal, and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2" (with Malt-dust).

Carcass (cold)	lbs.	281	lbs.	..	lbs.	123	lbs.	213½	lbs.	..	lbs.	222	lbs.	206	lbs.	189½	lbs.	217½
Offal parts, loss by fasting, evaporation, &c.	lbs.	76	lbs.	..	lbs.	59	lbs.	60½	lbs.	..	lbs.	74	lbs.	50	lbs.	52½	lbs.	62
Live-weight (unfasted)	lbs.	357	lbs.	..	lbs.	252	lbs.	274	lbs.	..	lbs.	296	lbs.	256	lbs.	242	lbs.	279½

TABLE LI.—EXPERIMENTS MADE WITH PIGS, at ROTHAMSTED, HERTS.
Per-centages of the Carcasses, &c., in the Live-weights (unfasted).

Designation of Parts.	PIGS.									
	No. 1.	No. 2.	No. 3.	No. 4.	No. 5.	No. 6.	No. 7.	No. 8.	Means.	
Lot 1.—Food—Pea-meal, and Unmalted Barley "No. 1."										
Carcass (cold) Offal parts, loss by fasting, eva- poration, &c.	79.09	80.73	77.27	78.14	77.04	80.13	82.57	79.11	79.26	
	20.91	19.27	22.73	21.86	22.96	19.87	17.43	20.89	20.74	
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	
Lot 2.—Food—Pea-meal, and Malted Barley "No. 1" (with Malt-dust).										
Carcass (cold) Offal parts, loss by fasting, eva- poration, &c.	78.36	78.15	77.69	82.92	77.46	78.95	75.00	80.10	78.58	
	21.64	21.85	22.31	17.08	22.54	21.05	25.00	19.90	21.42	
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	
Lot 3.—Food—Pea-meal, and Unmalted and Malted Barley "No. 1" (with Malt-dust).										
Carcass (cold) Offal parts, loss by fasting, eva- poration, &c.	76.54	76.17	..	78.21	77.44	78.04	78.06	77.52	77.43	
	23.46	23.83	..	21.79	22.56	21.96	21.94	22.48	22.56	
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

Carcass (cold) Offal parts, loss by fasting, evaporation, &c.}	78.30	79.96	77.37	79.87	78.76	80.00	78.66	76.10	78.64
	21.70	20.04	22.63	20.13	21.24	20.00	21.34	23.81	21.36
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

Lot 5.—Food—Pea-meal, and Malted Barley "No. 2" (with Malt-dust).

Carcass (cold) Offal parts, loss by fasting, evaporation, &c.}	80.19	77.98	78.86	77.14	78.53	77.78	77.85	75.00	77.92
	19.81	22.32	21.14	22.86	21.47	22.22	22.15	25.00	22.08
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

Lot 6.—Food—Pea-meal, and Mixture of four-fifths Unmalted and one-fifth Malted Barley "No. 2" (with Malt-dust).

Carcass (cold) Offal parts, loss by fasting, evaporation, &c.}	78.71	..	76.59	77.92	..	75.00	80.47	78.30	77.83
	21.29	..	23.41	22.08	..	25.00	19.53	21.70	22.17
	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00



LONDON:
PRINTED BY WILLIAM CLOWES AND SONS, STAMFORD STREET,
AND CHANCING CROSS.

ON THE

COMPOSITION, VALUE,

AND

UTILIZATION OF TOWN SEWAGE:

BY

J. B. LAWES, Esq., F.R.S., F.C.S.,

AND

J. H. GILBERT, Ph.D., F.R.S., F.C.S.

LONDON :

PRINTED BY HARRISON & SONS, ST. MARTIN'S LANE.

1866.

RE-PRINTED BY DUNN & CHIDGEY, 155-57, KINGSLAND ROAD.

1888.



CONTENTS.

	PAGE
Position of the Sewage Question	3
Composition and Value of Town Sewage	10
Crops to which Sewage is most applicable.....	27
Results of direct Experiment on the Utilization of Sewage.....	30
Experience of Common Practice in the Utilization of Sewage	45
General Practical Conclusions.....	50

[Reprinted from the Journal of the Chemical Society, March and April, 1866.]

*On the Composition, Value, and Utilization of Town Sewage.**

By J. B. LAWES, Esq., F.R.S., F.C.S., and J. H. GILBERT,
Ph.D., F.R.S., F.C.S.

Position of the Sewage Question.

It is no less true than strange that, after so many centuries of advance in regard to almost every other requirement of civilized life, the lesson should not yet have been learnt of how to dispose of the excretal matters of large populations, in such a manner as to secure both their collection and removal without nuisance and injury to health, and their economical utilization for the reproduction of food. But it is undoubtedly the fact that, hitherto, where utilization has been the most complete, comfort and health have generally been in the greatest degree sacrificed, and where, on the other hand, the refuse matters of town populations have been the most rapidly and perfectly removed from their dwellings, there has been either no utilization at all, or it has been most imperfectly attained.

Sewage, the foul stream which flows through the underground veins and arteries of our great cities, carrying with it the excretal and other refuse matters of large populations, hitherto to little better purpose than to be wasted, and to be a source of pollution to our rivers—to destroy their fish, injure their channels, and

* The substance of this paper was given as a Discourse before the Chemical Society, February 1, 1866, by Dr. Gilbert.

brought us in this country, so far as utilization and the use of our rivers are concerned, that some authorities, especially on the Continent, incline to the reactionary conclusion that the cesspool system, or rather the adoption of some inferior barrel, tank, or cesspool system, of collection and removal of admixture with extraneous water is inevitable.

Before, therefore, entering upon the question of the comparative value, and modes and results of the utilization of diluted sewage, it will be well to call attention, though very briefly, to some of the results of experience hitherto attained, and to the systems of town purification, and other modes of utilizing the products, than the modern ones by means of water.

China and Japan are frequently cited as affording examples of very perfect utilization of human excretal matters, and, in consequence, of great productiveness of the soil and great concentration of population on a given area of land. The method of collecting, removing, and transporting human excretal matters in those countries is, however, such as to be quite inadmissible by our modern notions of cleanliness, decency, comfort, and health.

It is frequently stated that in Belgium human excretal matters are very perfectly utilized, and realise considerable money to the town populations. Indeed, in one of the applications made only last year to the Metropolitan Board of Works for the consideration of the Southern sewage of the Metropolis, and still under consideration of that body, it is stated that the excretal matter sold in Belgium for something over £1 per person per annum.

practices adopted are attended with, at any rate, so much of nuisance and discomfort as would not now be permitted in this country; whilst it would appear that a considerable proportion of the urine of the populations escapes collection and utilization. As the result of the same inquiries, it was concluded that, in no case, did the town population realise by the disposal of their excretal matters as much as averaged one franc per head per annum.

The conclusion that, as a rule but little, and frequently nothing, is realised by town populations when their excretal matters are collected under more or less modified cesspool or tank systems, as free as possible from extraneous water, and so removed for application to the land, is fully confirmed by the results of an enquiry conducted by a Commission sent out by the Prussian Government in 1864, to investigate the modes of collection, removal, and utilization, in various localities, with a view to the adoption of improved plans for the city of Berlin.

The Prussian Commissioners, Herren C. v. Salviati, O. Röder, and Dr. Eichhorn, visited and reported upon, not only the Belgium towns of Ghent, Ostend, and Antwerp, but likewise Hanover, Cologne, Metz, Carlsruhe, Strasburg, Basle, Lyons, Zurich, Munich, Nuremberg, Dresden, and Leipzig; and their report shows not only that the householders seldom realised anything like a franc per head per annum for their excretal matters, but that in the majority of cases it cost them something for the removal. Nevertheless, looking to the position and local circumstances of Berlin, and especially to the results of the water-system in this country hitherto, the Commissioners deprecate the adoption of that system for that city, and recommend more perfect arrangements and more stringent regulations for the emptying and removal of the contents of existing cesspools, and, where practicable, the adoption of a system under which the excretal matters of each house are to be collected in a barrel placed at the bottom of the shaft leading from the closets, which, when removed is covered with a closely fitting lid, and is of such dimensions that two men can carry it by means of handles attached for that purpose. They seem to anticipate little, if any, pecuniary profit to the town from these arrangements, but consider that they will be attended with scarcely any, or even no, nuisance or discomfort, and that by their means a large amount of valuable manure will be provided in a convenient form for transport and utilization. There can, however, we think, be little doubt that under such a

system the collection and removal must be attended with considerable nuisance, that the greater part of the urine will be lost and that the cost of the collection, removal and transport will be such as to render the utilization unprofitable beyond a comparatively limited distance from the city.

There is little probability that the difficulties of the water-system will lead us in this country to have recourse again in our large towns to any system of cess-pools, tanks, or barrels, however improved; but it may be well here to notice one or two attempts that have been made within the last few years to obviate the use of water, and thereby to avoid the pollution of rivers, and secure the collection of the manurial matters in a form more readily transportable by ordinary means, and therefore, more applicable for general agricultural use: for there cannot be a doubt that if any system could be devised by which human excretal matters could be collected and removed from dwellings, without either nuisance or injury to health, and obtained economically in a concentrated, dry, and portable condition, their utilization would be much more perfectly attained by such means than is at all likely or even possible, under the water-system.

Perhaps the most noticeable attempt of the kind in question is that which has been made at Hyde, in Lancashire, a manufacturing town of more than 20,000 inhabitants. Some few years ago a company contracted to carry out what they call the "Eurek system." They provided boxes to fit in at the back of the privy or closet of nearly every house, leaving scarcely a water-closet in the place. Some disinfecting or deodorising mixture is put in the box before it is placed in its position, and the box is exchanged for a fresh one after a certain number of days, according to the number of individuals frequenting the place; and it is stipulated that neither extraneous water, nor any other than human excremental matters, should be accumulated in these receptacles. The boxes, when removed, are covered with closely fitting lids, and so transported in closed vans to a manure manufactory close to the town. Here the matters are first well mixed, and then strained to remove rags, which are washed and sold for paper-making. More disinfectant is then added, and the matter concentrated by distillation, the distilled water being sold to dyers and bleachers. The residue thus thickened is then mixed with coal-ashes, which are collected in the houses in casks left for the purpose, and before being used are re-burnt in a reverberatory furnace, and finely ground.

On visiting Hyde in 1863, it certainly appeared that the mode of collection and preparation adopted was attended with, at any rate, very little unpleasant odour, and it was maintained by the advocates of the system, that its adoption had been successful in a sanitary point of view; though even at that time some difference of opinion existed, and a controversy on the subject was in progress. The system is still in operation; but we are informed that the feeling of the inhabitants is very strong against the maintenance of the works in the neighbourhood; indeed, that an injunction against them has been sought, though unsuccessfully, and that proceedings by indictment are now being taken. This opposition has reference not to the mode of collection, but to the conducting of the manufacture so near to the town. But, whether or not, the plan of collection and removal may have proved successful, so far as the avoidance of nuisance and injury to health are concerned, the process of manufacture seems, unfortunately, to offer but little prospect of successful utilization, so far at least as can be judged from the results of an analysis made at Rothamsted, of a sample of the manure obtained direct from the works. It was found to contain only between 1 and 2 per cent. of ammonia. Such a manure, although it might be useful enough when applied in quantities of many tons to the acre, would obviously be not worth more than its carriage beyond the distance of a very few miles. Besides the great dilution of the more valuable manurial matters by the admixture with ashes, a little consideration of the habits of the people is sufficient to account for the small quantity of ammonia found in the manure; for it is obvious that little of the urine beyond that passed once a day with the fæces would reach the boxes, and so find its way into a manure thus collected and prepared.

One more dry system, the offspring of the difficulties of the wet one, should be briefly noticed, namely, that of the Rev. Mr. Moule. Mr. Moule has invented and patented an arrangement for the use of dry sifted earth, instead of water. He states that by the use of about 4 lbs. per head per day of finely sifted clay, deposited by means of a mechanical arrangement upon the fæcal matters as soon as passed, they are at once entirely deodorised, and in a few weeks are so entirely disintegrated that neither excretal matters nor paper can be detected in the mass, which, he says, looks and smells like fresh earth, and may, after resifting, be re-used, until it has done

duty four times over, by which, of course, there is not only a great saving of material, but the value of the manure is considerably increased.

Very obvious objections to such a system are—the difficulties of the supply and preparation of the soil in the case of towns, or even in the country in wet seasons; the fact that but little of the urine containing as it does so large a proportion of the valuable manure constituents of human excretal matters, would reach the compost prepared; and that, in the manure produced, the more valuable matters would be diluted with so large a proportion of comparatively useless material, that beyond a very short distance the cost of carriage would be all that the manure was worth. On the other hand, that the adoption of such a system would be a great improvement in a sanitary point of view, in the cases of sick rooms, detached houses, or even villages, where the water-system is not available, and that it might be even economical where the expense for preparation and absorption, and the land for utilization, are in close proximity, may, perhaps, be readily granted. But we are certainly not so sanguine as the Rev. Mr. Moule, who seems to think that with the aid of Earth-closet Companies, his plan is as practicable for large towns as is the supply of water, gas, and coal, at present, and much more so than the removal and utilization of dilute town sewage.

Whilst it must be admitted that the agricultural utilization of human excretal matters has, hitherto, been much more completely attained under the system of collection without water than under our new one with it, it must not be forgotten that neither on the continent of Europe nor in this country has such utilization resulted in any substantial profit to the towns; and that it is, with the recorded results of China and Japan before us, and after many centuries of experience nearer home, of at least comparatively successful utilization, that the old systems have been abandoned as utterly inconsistent with advance in habits and notions of cleanliness, and with the maintenance of the comfort and health of large populations. Nor do the modifications of the dry system to which brief reference has been made, seem to hold out a hope of general and permanent applicability to large populations looking, as we must, to the combined requirements of convenience, comfort, health, and utilization. Our water-system of hot defecation and town cleansing is, on the other hand, scarcely more than a generation old. By its means excretal and other refuse

rs are more rapidly removed from dwellings than is possible by other; and, independently of the increased comfort and freedom from nuisance obvious to all, sanitary statistics have amply shown increased immunity from zymotic diseases, and increased longevity, as the result of the adoption of that system. It is that these advantages have, hitherto, been attained at the cost of the almost universal sacrifice of the manure, and of injury to our rivers.

It is, then, admittedly the existing dilemma of our modern cities. But public attention is now so thoroughly directed to the subject, that little fear need be entertained that either the systematic non-utilization of the sewage, or the pollution of our rivers by it, will long be permitted. Least of all is it reasonable to be discouraged in the fact, that the system which has done so much for some of our town populations in so short a time, and, not, at this early stage of its trial, have accomplished all that might be desired, or to conclude that the nuisances and difficulties incident to the old plans, which have remained unremedied for so many centuries, have much better chance now than formerly of being successfully obviated.

Assuming that there is more likelihood of the general application, success, and permanence of the water-closet system of urban defecation, it becomes important to consider the position, the value, and the modes and results of the utilization of the product of that system, namely, *dilute town sewage*.

Many plans have been proposed for the separation of the valuable constituents from sewer-water, and the manufacture of them into a dry and easily portable manure. But whilst several of these have been successful in separating the whole of the insoluble elementary matter, and even some small portion of the soluble constituents, leaving the fluid to a great extent, or at any rate sparingly, purified, and in a much less objectionable condition for running into rivers, none have succeeded in either adequate or permanent purification, or in the separation of the more valuable mineral matters, and the production of a concentrated solid fertilizer-manure, having a sufficient value to be remunerative, and to bear the cost of transport more than a very few miles;* nor

* Information in regard to some of the plans proposed for the partial purification of sewage-water, or for the separation of a solid manure from it, see—"On the Application of Sewage to Agriculture," by Dugald Campbell, Esq., F.C.S., Soc. Qu. J. vol. x., p. 272. "Report of Chemical Investigations relating to

when we consider the great solubility of some of the more active manurial constituents of sewage, and the great dilution of them in the sewage, can any hope be held out of so desirable a consummation;—desirable, indeed, for if human excretal matters, the residue of the constituents consumed as food, cannot be recovered in the form of a concentrated, dry, and easily transportable manure, little hope can be entertained of their re-distribution over anything like the area from which they came, or of their general use for the direct reproduction of the varied descriptions of food which were their source.

The questions arise: What is the amount, and what approximately the money value for the purposes of manure, of the constituents contributed to sewage by a given population? What their state of dilution in sewer-water? To what soils and crops is dilute sewage the most applicable? What is the money-value realisable in practice by sewage utilization? What are the conditions of profit or loss to towns of such utilization?

Composition and Value of Town Sewage.

It is one thing to determine the amount of constituents contained in sewage, or contributed to it by a given population, and to estimate their value accordingly, as if they existed in the dry and portable condition of the various concentrated manures of known value in the market; but it is obviously quite another to settle the really available or realisable value of the same constituents when they are distributed through an enormous volume of water, and if they must be transported and utilized in that condition. Let us first consider what may be called the theoretical value of the constituents of sewage, or their estimated value, taking as the measure the value of the same constituents in dry and portable manures.

Numerous authorities have undertaken the consideration of this question, and two chief methods have been adopted. One of these has been to take samples of sewage and determine its composition by analysis, to adopt such estimates as are at command relating to the amount of sewage available within a given time or

The Metropolitan Main Drainage Question," by A. W. Hofmann, LL.D., F.R.S., and Henry M. Witt, F.C.S., Report on Metropolitan Drainage, 1857. **Deodorization of Sewage, Second Report of the Royal Sewage Commission, 1862, Appendix No. 6, p. 64.**

from a given population, and so to reckon the amount and value of the constituents in a given quantity of sewage, or per head, or or a given number of persons, per annum. Another is to base the calculation upon the amounts of fæces and urine, or of the various constituents of these, which have been recorded as voided by individuals of different sexes and ages—sometimes making allowance, and sometimes not doing so, for other than human excretal matters reaching the sewers.

First, as to the results attained when the calculation is based upon the analysis of sewage, and estimates of the amount of it yielded by a given population.

In estimates of the value of the constituents of sewage, about three-fourths of the total value has generally been attributed to the ammonia (or nitrogen reckoned as ammonia); and it so happens that if a value of 8d. be put upon every lb. of ammonia shown by analysis to be contained in sewage, or if for each grain of ammonia per gallon, a value of one farthing be given to the total constituents in 1 ton of the sewage, the result will, in either case, agree almost exactly with that obtained by the elaborate method of giving the currently adopted market values to the several constituents, taking dry and portable manures as the standard.

One or two illustrations may be given of the applicability of the latter mode of reckoning. In the summer of 1863, Baron Liebig, adopting as the basis of his calculations an analysis of the Dorset Square sewage by Mr. Way, which showed nearly 18 grains of ammonia per gallon, estimated that (provided the quantity of phosphates which he considered requisite to render the whole of the ammonia available were employed with the sewage) the constituents in 1 ton of sewage of that composition would be worth about 4d. Now, according to our mode of estimate stated above, 18 grains of ammonia per gallon would indicate a value of 18 farthings, or 4½d., for the total constituents in 1 ton of the sewage. In January, 1865, Baron Liebig assumed the average sewage of the Metropolis to contain only 7·2 (instead of 18) grains of ammonia per gallon; and he estimated the value of the constituents in 1 ton of such sewage to be rather over 1½d. Our estimate would also give rather over 7 farthings, or 1½d. Lastly on his point, in 1857, Messrs. Hofmann and Witt concluded from their investigations that the average *dry weather* sewage of the Metropolis contained about 8·2 grains of ammonia per gallon;

and calculating the value of the sewage according to the amount of ammonia, organic matter, phosphoric acid, and potassa, they estimated that of the total constituents in 1 ton of such sewage to be about 2·11d. It is clear that giving a value of $\frac{1}{4}$ d. to the total constituents per ton of sewage, for each grain of ammonia per gallon, would yield almost identically the same result.

It is obvious, therefore, that in this part of the discussion we may, for all practical purposes, safely disregard everything but the amount of ammonia contained in the sewage, and that by so doing the consideration of the subject will be greatly simplified. It will be seen, too, that in adopting this course we do not in any way ignore, or undervalue, the importance of the associated constituents, but, on the contrary, accord to them the same value as Baron Liebig, Messrs. Hofmann and Witt, and others, have done by a much more elaborate process of calculation.

Numerous analyses have been made from time to time of samples of the Metropolitan and other sewage; and sometimes very important theoretical conclusions, and even propositions for the investment of enormous amounts of capital in utilization schemes, and anticipations of enormous profits from their adoption, have been based upon the results of a single analysis. Such, however, is the variation in the dilution of the sewage of any one locality at different times, that it is quite impossible to draw any safe conclusions from the results of analysis without carefully taking into consideration the circumstances affecting the dilution at the time of sampling. This is strikingly illustrated by the results given in Table I., in which are recorded the grains of ammonia per gallon, as determined by various experimenters, in samples of the Metropolitan sewage, taken at different times and places, and also the estimated value of the total constituents in one ton of the sewage, reckoned according to the number of grains of ammonia per gallon as above referred to.

TABLE I.

of Ammonia per gallon in different samples of Metropolitan Sewage, and estimated value of constituents in one ton.

Authority.	Name of Sewer.	Time of Sampling.	Ammonia per Gallon.	Estimated Value per ton.
			Grains.	d.
.....	Barrett's Court	Day	41·28	10½
	Dorset Square	Day	17·96	4½
	The Fleet.....	Noon	5·15	1½
		Midnight	8·50	2
	London Bridge	Noon	6·69	1½
		Midnight	8·10	2
	Dowgate Dock	Noon	10·03	2½
		Midnight	3·43	0½
	Iron Gate.....	Noon	8·13	2
		Midnight	6·20	1½
	Paul's Wharf	Noon	12·01	3
		Midnight	3·13	0½
	Whitefriar's Dock	Noon	5·35	1½
		Midnight	3·41	0½
.....	Custom House, West ...	Noon	6·25	1½
		Midnight	8·17	2
	Custom House, East ...	Noon	7·28	1½
		Midnight	15·01	3½
	Hambro' Wharf	Noon	7·69	2
		Midnight	5·69	1½
	Wool Quay	Noon	6·95	1½
		Midnight	5·00	1½
	Tower Dock.....	Noon	10·02	2½
		Midnight	7·15	1½
	Mean.....	7·24	1½
nn & Witt.....	Savoy Street	24 hours	8·21	2½

the results given at the head of the table, on the authority of Way, are those of probably the first analyses made of the Metropolitan sewage, and it is only fair to say that at the time he published them, he expressly stated that although they showed there was great manurial value in sewage, yet they could not be taken as in any way affording a measure of that value. It was, however, upon the analysis of the sample of the Dorset Square sewage, showing nearly 18 grains of ammonia per gallon, that Baron Pigou based his calculations as to the value of the Metropolitan sewage in 1863; and the advocates of particular sewage schemes, even members of Parliamentary Committees, have sought to build much upon the results of those analyses.

From the varying circumstances under which the sample analysed by Dr. Letheby were taken, as indicated in the table is obvious that the results, though very valuable in that respect must be considered rather as illustrations of the variation in composition of the Metropolitan sewage at different times and places, and as showing the danger of founding important practical conclusions upon the results of the analysis of an individual sample, than as affording direct evidence as to the average composition of the Metropolitan sewage.

The sample analysed by Messrs. Hoffmann and Witt was a mixture of equal portions taken every hour during twenty-four hours of dry weather, and there is no doubt that that sample has better claims to be taken as representing the average dry weather sewage of the Metropolis than any other that had up to that time been collected and examined. It was upon the analysis of this sample that Messrs. Hoffmann and Witt, calculating the value of the ammonia, organic matter, phosphoric acid, and potash which it contained, estimated that the constituents in one ton of such dry weather sewage would be worth rather over 2d., and according to the information supplied to them for the purpose of their calculations, the quantity of sewage, exclusive of rainfall, would be about 158,000,000 tons per annum, or scarcely three fifths as much as that assumed in the estimates of Baron Liel and Mr. Thomas Ellis, as the total sewage, namely, 266,000,000 tons. Yet, Messrs. Hoffmann and Witt's estimate of a little over 2d. for the value of the constituents in one ton of the normal dry weather sewage was taken by Mr. Ellis, in his application for a concession of the Metropolitan sewage, as applying to the whole amount of dilute sewage (inclusive of rainfall and subsoil water) which he estimated would be available for utilization (266,000,000 tons), and his calculations of profit to his Company and to the ratepayers were based upon this erroneous assumption.

To conclude in reference to the results recorded in Table I, attention may be called to the fact that the different samples show a variation of from about 8 to more than 41 grains of ammonia per gallon, representing approximately a difference of from about $\frac{1}{4}$ d. to about $10\frac{1}{4}$ d. for the estimated value of the total constituents in one ton of the sewage.

That the results of an analysis of a sample of sewage of a locality taken without careful reference to the circumstances of dilution, are not only entirely inadequate as the basis of gene-

conclusions, but may even be utterly misleading, is even more strikingly illustrated by the results next to be considered, which were obtained in the course of an investigation undertaken by the late Royal Sewage Commission.

Three members of the Commission, the late Mr. Henry Austin, C.E., Mr. Way, and one of the authors (J. B. Lawes) were appointed a sub-committee to undertake an investigation on the utilization of sewage. The agricultural experiments were conducted at Rugby, and their management, and the selection, collection, and preparation of samples for analysis, devolved upon the authors, the analyses being made in the laboratory of Mr. Way. The inquiry extended over a period of between three and four years, and involved the application of different quantities of sewage to meadow-grass and some other crops; the determination of the amounts of produce obtained; the feeding of fattening oxen and milking cows on the unsewaged and the sewaged grass; and the sampling, and more or less complete analysis, of the soil, of the sewage, of the drainage-water from the irrigated land, of the unsewaged and the sewaged grass, of the milk yielded by the cows fed upon it, &c., &c. It is proposed to embody in the sequel a brief abstract statement of some of the more important facts and conclusions brought out by the experimental inquiry above referred to, and the reader is referred for all fuller details to the Reports of the Commission.*

The mode of collecting samples of the Rugby sewage for analysis was, to take about a quart (from a gauge-tank holding between 3 and 4 tons, through which the sewage flowed before passing on to the land), at intervals of about two hours for several days together, well mix the quantity so accumulated, and take a sample of the mixture for analysis. 93 such mixed samples were collected and analysed, the period of collection extending over 31 months, from April, 1861, to October, 1863, inclusive. Table II. shows the highest, the lowest, and the average amounts of ammonia, and total solid matter, which the analyses of these numerous mixed samples indicated.

*Second Report of the Commission appointed to inquire into the best mode of Distributing the sewage of Towns, and applying it to beneficial and profitable uses; 1862. Third do. do., 1865.

TABLE II.

Showing the highest, lowest, and average amounts of Ammonia, and total Solid Matter, in mixed samples of Rugby Sewage at different times.

		Ammonia.		Total Solid Matter.	
		Grains per Gallon.	lbs. per 1000 Tons.	Grains per Gallon.	lbs. per 1000 Tons.
1861	Highest	15·64	500·5	216·5	6928
	Lowest	2·99	95·7	37·6	1213
	Mean of 24 analyses	6·39	204·5	75·1	2405
1861-2	Highest	11·38	364·2	129·3	4138
	Lowest	2·55	81·6	50·5	1616
	Mean of 34 analyses	5·95	190·4	80·3	2570
1862-3	Highest	12·81	409·9	269·9	8637
	Lowest	3·14	100·5	62·2	1989
	Mean of 35 analyses	7·08	226·5	103·2	3302

Thus, although each sample analysed was a mixture of samples taken over several days together, as above described, there was a variation among the 93 samples of from $2\frac{1}{2}$ to $15\frac{1}{2}$ grains of ammonia, and from $37\frac{1}{2}$ to 270 grains of total solid matter, per gallon; or, of from $81\frac{1}{2}$ to $500\frac{1}{2}$ lbs. of ammonia, and from 1203 to 8,637 lbs. of total solid matter, per 1,000 tons of sewage. Reckoned according to the number of grains of ammonia per gallon, the estimated value of the total constituents in 1 ton of sewage varied from about $\frac{1}{4}$ d. to nearly 4d.

Notwithstanding the very great differences in the composition of the Rugby sewage at different times, much greater, indeed, than could have been expected, considering the circumstances of the sampling, it is still believed that the mean of so many determinations may be taken as indicating, at any rate approximately, the average composition of the Rugby sewage over the period to which they refer. The probability of this will be seen on a consideration of the average results for each of the three seasons, and for the total period of 31 months of collection, given in Table III.

TABLE III.

Mean Composition of Rugby Sewage, in 1861, 1862, and 1863.

Constituents.		Means of			
		24 Samples April to Oct. 1861.	34 Samples Nov. 1861 to Oct. 1862.	35 Samples Nov. 1862 to Oct. 1863.	93 Samples April 1861 to Oct. 1863.
Grains per gallon.					
In suspension	Inorganic	14·36	20·86	34·45	24·30
	Organic	14·16	16·84	24·03	18·85
	Total	28·52	37·70	58·48	43·15
In solution	Inorganic	36·34	34·42	36·80	35·81
	Organic	10·28	8·20	7·92	8·63
	Total	46·62	42·62	44·72	44·44
Total inorganic ...		50·70	55·28	71·25	60·11
Total organic		24·44	25·04	31·95	27·43
Total solid matter .		75·14	80·32	103·20	87·59
Ammonia	In suspension.....	1·41	1·47	1·86	1·60
	In solution	4·98	4·48	5·22	4·89
	Total	6·39	5·95	7·08	6·49
lbs. per 1000 tons.					
In suspension	Inorganic	460	668	1102	778
	Organic	453	539	769	603
	Total	913	1207	1871	1381
In solution	Inorganic	1163	1101	1178	1146
	Organic	329	262	253	276
	Total	1492	1363	1431	1422
Total inorganic ...		1623	1769	2280	1924
Total organic		782	801	1022	879
Total solid matter .		2405	2570	3302	2803
Ammonia	In suspension	45	47	60	51
	In solution	159	143	167	157
	Total	204	190	227	208

It is seen that the mean result of the analyses of 24 samples collected from April to October, inclusive, 1861, indicates 6.3 grains of ammonia per gallon; that of 34 samples collected from November 1861, to October 1862, inclusive, 5.95 grains, and that of 35 samples, collected from November 1862, to October 1863, inclusive, 7.08 grains. This difference in the average concentration of the sewage of the different seasons is perfectly consistent with the difference in the character of the seasons themselves. Thus, the season of 1861-2 was much the wettest, and its sewage was, accordingly, the most dilute; the season of 1862-3 was much the driest, indeed extremely dry, and its sewage was the strongest; and the season of 1861 being intermediate in this respect, its sewage was of intermediate strength.

Looking to the average result of the 93 analyses, it will be observed that the sewage contained about $87\frac{1}{2}$ grains per gallon of total solid matter, of which about two-thirds was inorganic, and one-third organic. About half of the total solid matter was in suspension, and half in solution: of the half in suspension about four-sevenths was inorganic and three-sevenths organic, and of the half in solution, about four-fifths inorganic, and one-fifth organic. Lastly, of the nitrogen reckoned as ammonia, about one-fourth was in suspension, and three-fourths in solution.

The mean of the 93 analyses shows about $6\frac{1}{2}$ grains of ammonia per gallon, indicating a value of about $1\frac{3}{4}$ for the total constituents in 1 ton of the sewage. But taking into consideration the fact that the samples were not collected at exactly equal intervals throughout the total period, it is concluded that, by taking the mean result for each of the 31 months separately, and then the mean of the 31 means so obtained, the result will more nearly represent the real average composition of the sewage of the whole period, than will the direct mean of the 93 analyses; and the calculated average so obtained indicates about 7, instead of only $6\frac{1}{2}$, grains of ammonia per gallon.

From all the information at command as to the population contributing to the sewers, the water-supply, the rainfall, and the drainage area, it was concluded that, taking the average of seasons, there are about 60 tons of sewage per head of the population of Rugby, per annum; but that, as the period of the experiments was drier than usual, the amount probably then reached to only about 55 or 56 tons.

Now, if we reckon $6\frac{1}{2}$ grains of ammonia per gallon, and 60

tons of sewage per head per annum, it would result that $12\frac{1}{2}$ lbs. of ammonia were contributed annually for each average individual of the mixed population, of both sexes and all ages; or, if we reckon 7 grains of ammonia per gallon, and 56 tons of sewage per head per annum, we equally arrive at the amount of $12\frac{1}{2}$ lbs. of ammonia per head per annum; and from a careful consideration of the Rugby results, it was concluded, at the time the Report was issued, that this probably very nearly represented the actual fact.

Having, then, by means of the results of a great many analyses of sewage, and a consideration of the amount of sewage obtained from each average individual of the population, estimated that for each average individual there would be about $12\frac{1}{2}$ lbs. of ammonia annually contributed to the sewer-water, let us next see what

TABLE IV.

Amount of Nitrogen reckoned as Ammonia, and estimated value of total Constituents, in Human Voidings, per head per annum.

	Ammonia	Value of Total Constituents.
Adult Males; Hofmann and Witt.		
Urine.....	lbs. 15·8	s. d. 10 0½
Fæces.....	2·3	1 8½
Total.....	18·1	11 9¼
Adult Males; Thudichum.		
Urine.....	15·9	10 3½
Average, both sexes and all ages; Hofmann, Witt, and Thudichum.		
Urine.....	11·32	7 3
Fæces	1·64	1 2½
Total.....	12·96	8 5½
Average; both sexes and all ages; Lawes and Gilbert.		
According to	Food.....	12·2
	Voidings	12·6
	Voidings	12·7
	Mean	12·5
		} 8 4

Witt took the amount of urine estimated to be daily voided by an adult, and the amount of fæces recorded as voided on the average per head of the body-guard of the Grand Duke of Darmstadt (but allowing, as they said, a little more for Bull"), and applying the results of Berzelius' analysis and those of the analysis of Way, Liebig, and Wesarg, they calculated the amount of ammonia, and other constituents daily voided by such persons. According to their data, the amount of ammonia annually voided by an adult male was, in urine 18·1 lbs.; and the estimated money value of constituents was in urine 10s. 0½d., in fæces 1s. 8 11s. 9½d. The result so obtained for adult males is applicable to each individual of a mixed population, of both sexes and all ages, assuming that other matters reaching the excreta would probably make up the difference. There can be little doubt that this was making far too liberal an allowance for other human excretal matters contributing to the value of the excreta.

Some years later, in 1863, Dr. Thudichum, from much more comprehensive data, gave for the urine alone of an adult male 15·9 lbs. of ammonia, and 10s. 3½d. of value; amounts which will be seen, are almost identical with those of Messrs. Witt.

But Dr. Thudichum, instead of directly applying the result obtained for an adult male to each average individual of the population, considered that two adult males would appri-

sexes and all ages. By this process, as the Table shows, we have early 18 lbs. of ammonia, and nearly 8s. 6d. of value, to represent the mixed voidings of such an average individual.

In 1854, the authors, basing their estimates on very comprehensive data, relating both to the amounts of constituents consumed in the food, and voided in the urine and fæces, of persons of different ages and both sexes, concluded that probably about 18 lbs. of ammonia, and total constituents of the estimated annual value of about 6s. 8d., were annually contributed to the sewage per individual of a mixed town population. More recently, for the purposes of the Report of the Royal Sewage Commission, the estimates relating to the constituents voided were carefully revised, bringing into the calculations such further information as was then at command; * and the results so obtained are recorded in the Table (IV).

The amount of nitrogen estimated to be annually consumed in the food of an average individual was deduced from the calculation of 86 dietaries, arranged in 15 classes, according to sex, age, activity of mode of life, and other circumstances, and corresponded to about 12·2 lbs. of ammonia; from which, of course, a deduction has to be made for the nitrogen retained in the body, and for losses in various ways. When the calculation was based upon determinations or computations of the amounts of nitrogen or ammonia-yielding matters voided by persons of different sexes and ages, the result arrived at was 12·6 lbs. of ammonia; and when upon the recorded amounts of fresh urine and fæces voided, and the average composition of these, the amount indicated was 12·7 lbs. of ammonia per head per annum. A careful consideration, however, of the circumstances of the majority of the cases contributing to the averages among those divisions of the population in relation to which the evidence is the most plentiful, and of the relative character of the results where it is the most deficient, led to the conclusion that the estimate of 12·6, or 12·7 lbs. for the amount

* For nearly the whole, if not the whole, of the data upon which the new estimates are based, see "On the Sewage of London," by J. B. Lawes, F.R.S., *Journal of the Society of Arts*, March 9, 1855; "The Composition of the Urine in Health and Disease," by E. A. Parkes, M.D., 1860; "On an Improved Mode of collecting Excrementitious Matter, with a view to its Application to the benefit of Agriculture, &c.," by J. L. W. Thudichum, M.D., F.C.S., *Journal of the Society of Arts*, May 15, 1863; and "On the Elimination of Urea and Urinary Water, in relation to the period of the Day, Season, Exertion, Food, &c., &c.," by Edward Smith, M.D., LL.B., F.R.S., *Philosophical Transactions*, vol. cli, p. 747.

of ammonia voided annually by an average individual of a mixed population, was in all probability too high.

Reviewing the whole of the evidence, both that relating to the composition and the amount of the Rugby sewage, and that relating to the amount of constituents voided by an average individual, it was concluded that the amount of ammonia annually contributed to the sewer-water by an average person of a mixed population was pretty certainly more than 10 lbs., as formerly assumed, but probably less than 12 lbs.; and, making allowance for the fractional part of the excretal matters of horses, cows, dogs, and other animals, of the refuse of slaughter-houses, soot, and of other refuse matters that may reach the sewers, it was concluded that still not more than 12½ lbs. of ammonia would be contributed annually to the sewers from all sources, per head of mixed town population. This would indicate an estimated value of 8s. 4d. per annum for the total constituents in the sewage for each average individual.

It was admitted, however, to be a great desideratum, that when the Main Drainage of the Metropolis came to be completed, and the works to be in full operation, competent persons should be appointed to superintend the gauging, sampling, and analysis of the sewage, with a view to providing data which might serve to determine satisfactorily and conclusively the approximate amount, and average composition, of the Metropolitan sewage, as it will have to be dealt with in any plan of utilization, and also the relation of population to the composition of sewage generally.

Since the publication of the report of the commission, in March 1865, numerous gaugings and samplings of the sewage of the mid- and high-level sewers North of the Thames have been undertaken, and many samples have been analysed by Mr. Way and Dr. Odling. The results of this inquiry have not yet been published; but from information kindly communicated by Mr. Way, we are enabled to state their general bearing, so far, upon the point now under consideration.

From these new results it appears very probable that the amount of dry weather sewage averages only about two-thirds as much per head of the population as that generally supposed before, and assumed both in the inquiries of Messrs. Hofmann and Witt, and in the Report of the Sewage Commission; but the average amount of ammonia per gallon now found by Mr. Way in the dry weather sewage very nearly approaches that arrived at by

Messrs. Hofmann and Witt. Both Mr. Way and Mr. Cresy admit, however, in accordance with common experience, that further a subject is investigated, that there are still many questions, the settlement of which may materially affect the proper interpretation of the new gaugings.

Assuming them to indicate the result at present supposed, and as we have stated, it follows that the total amount of ammonia yielded by a given population will be only about two-thirds as much as was estimated by Messrs. Hofmann and Witt, on applying the results of their analysis to the higher estimated amount of the weather sewage. It further follows, from the same evidence, that the amount of ammonia annually contributed to the sewage, from all sources, per head of a mixed population, is more nearly 10 lbs., as formerly concluded by the authors, than $12\frac{1}{2}$ lbs., as recently estimated; and if this result should be confirmed, the former estimate of 6s. 8d. will more nearly represent the calculated annual value of the total constituents yielded per head of the population than the more recent one of 8s. 4d. It would have to be concluded, as indeed is not improbably the case, that, in the calculations based on the mean composition and the estimated total amount of the Rugby sewage, the latter had been put at too high a figure, too large a proportion of the rainfall having been assumed to reach the sewers; and that, in the estimates founded on the recorded amounts of constituents found, the incompleteness of the records, as already pointed out, as was supposed, led to too high an estimate.

We have, then, from 10 to $12\frac{1}{2}$ lbs. of ammonia, and an estimated value of from 6s. 8d. to 8s. 4d. for the total manurial constituents, contributed to sewage by each average individual of a mixed town population. Adopting these amounts, the questions arise—What will be the amount of ammonia, and what the estimated value of the constituents, in a given amount of sewage, at different dilutions? These points are illustrated in Table V.

TABLE V.

Ammonia per gallon, and estimated value of total Constituents in one ton of Sewage at different dilutions.

Dilutions supposed.		If 12½ lbs. Ammonia, per head per annum, from all sources.		If 10 lbs. Ammonia, per head per annum, from all sources.	
Per head per annum.	Per head per day.	Ammonia per gallon.	Estimated value per ton.	Ammonia per gallon.	Estimated value per ton.
Tons.	Gallons.	Grains.	Pence.	Grains.	Pence.
20	24½	9.77	2.44	7.81	2.00
30	36½	7.81	1.95	6.25	1.60
40	48½	6.50	1.67	5.21	1.33
50	60½	5.56	1.43	4.46	1.14
60	72½	4.88	1.25	3.91	1.00
80	96½	4.34	1.11	3.47	0.89
100	120½	3.91	1.00	3.13	0.80
200	240½	1.95	0.50	1.56	0.40

According to the information supplied to Messrs. Hofmann and Witt, the dry weather sewage of the Metropolis amounted to between 36 and 37 gallons per head per day = about 60 tons per head per annum. Their analysis showed 8.2 grains of ammonia per gallon, equivalent to about 15½ lbs. of ammonia per head per annum; and they reckoned the total constituents in 1 ton of such sewage to be worth 2.11d. But Table V shows that with a dilution of 60 tons, and with 12½ lbs. of ammonia per head per annum, there would be only 6.5 grains of ammonia per gallon, and total constituents in 1 ton of sewage worth only 1½d.; and that with only 10 lbs. of ammonia per head per annum, there would be only 5.2 grains per gallon, and constituents worth only 1¼d. in 1 ton of the sewage.

If, however, we take the dry weather sewage as indicated by the recent gaugings as more nearly 24 gallons per head per day = a rate of 40 tons per head per annum, we have then, with 12½ lbs. of ammonia per head per annum, 9.77 grains per gallon, and 2.44d. worth of constituents per ton; or, taking 10 lbs. of ammonia per head per annum, we have 7.8 grains per gallon, and constituents in 1 ton of an estimated value of nearly 2d. Now, Mr. Way's conclusion is, that the mid- and high-level dry weather sewage North of the Thames averages scarcely, but nearly, 8 grains of

ammonia per gallon, or almost exactly the amount last mentioned ; and as Messrs. Hofmann and Witt's analysis shows 8·2 grains, it will be seen that both estimates, taken in connection with the amended one as to the daily amount per head of the dry weather sewage, go to confirm the assumption that the amount of ammonia contributed to the sewage from all sources is much more nearly 12½ lbs. per head per annum.

Whatever may eventually prove to be the average dilution of the dry weather Metropolitan sewage, the actual amount of fluid varies immensely from time to time, according to rainfall and other circumstances. When it exceeds a certain amount, as in the case of continuous rains or storms, a portion will pass at once into the Thames ; and according to Mr. Bazalgette's figures it appears that this will happen when the volume is such as, if continuous, would represent something over 200 tons of fluid per head per annum. But, so far as information at present at command enables us to judge, it is probable that the amount, inclusive of rainfall and subsoil water, that will be available for utilization, will be somewhere about 80, and will pretty certainly not exceed 100 tons per head per annum ; that is, about twice, or not more than twice and a half, as much as the most recently estimated dry weather flow. Of course, to result in anything like such averages, the dilution would sometimes be at a rate very much greater than those amounts would indicate. But it may be observed, by way of illustration, that with 12½ lbs. of ammonia per head per annum, and an average of 80 tons of sewage, it would average less than 5 grains of ammonia per gallon, and only 25d. worth of constituents in 1 ton ; or, reckoning an average dilution of 100 tons, it would average less than 4 grains of ammonia per gallon, and only 1d. of value of constituents in 1 ton. In like manner, reckoning only 10 lbs. of ammonia per head per annum, a dilution of 80 tons would show less than 4 grains, and of 100 tons little over 3 grains of ammonia per gallon, and an amount of constituents in 1 ton worth only 1d. and 0·8d. respectively.

In comparison with the figures just given, it may be stated that both Baron Liebig, and Mr. Thomas Ellis (one of the applicants for the concession of the Metropolitan sewage) assume its total amount at 266,000,000 tons per annum, which, with 1,000,000 population, represents nearly 90 tons per head per annum ; and with this dilution, the former estimates the sewage

to contain an average of 7·2, and the latter 8·2 grains of ammonia per gallon; the latter, as already stated, applying the estimate Messrs. Hofmann and Witt for the dry weather sewage to the total estimated amount of available sewage, inclusive of rainfall.

It is sufficiently obvious that, however variable, the dilution of the constituents in town sewage is at any rate very great, and that in any scheme for the utilization of sewage large quantities will have to be dealt with. It will be useful, therefore, by way of illustration, and as a means of conveying a more definite idea of the extent of this dilution, to show the relation of a given amount—say 1,000 tons—of sewage of certain assumed dilutions, both to population, and to some well-known portable manure, such as Peruvian guano. This is done in Table VI, which shows the amount of guano which would supply as much nitrogen reckoned as ammonia as 1,000 tons of sewage of different dilutions, and the number of tons of sewage which would be equal in this respect to 1 ton of guano, and both on the alternative assumptions of 12½ lbs., and 10 lbs., of ammonia per head per annum. The assumed dilutions are 40, 50, and 60 tons per head per annum which may be taken to cover the minimum and maximum estimated rates of flow for the dry weather sewage of the Metropolis: 80 and 100 tons, which may be taken to represent the range for the average total available sewage, inclusive of rainfall and subsurface water, and 200 tons, the probable frequent dilution in wet weather.

TABLE VI.

Relation of Sewage to Peruvian Guano in amount of Nitrogen reckoned as Ammonia.

If Sewage per head per annum.	Contributing 1,000 tons Sewage.	If 12½ lbs. Ammonia, per head per annum, from all sources.		If 10 lbs. Ammonia, per head per annum, from all sources.	
		1,000 tons Sewage = Guano.	1 ton Guano = Sewage	1,000 tons Sewage = Guano.	1 ton Guano = Sewage
Tons.	Persons.	Cwts.	Tons.	Cwts.	Tons.
40	25	16½	1220	13	1525
50	20	13	1525	10½	1900
60	16½	11	1830	8½	2290
80	12½	8½	2440	6½	3050
100	10	6½	3050	5½	3810
200	5	3½	6100	2½	7620
1 Person = Guano.		½ cwt.		½ cwt.	

Thus, with $12\frac{1}{2}$ lbs. of ammonia, and the minimum estimated dilution of the dry weather sewage at a rate of 40 tons per head per annum, 1,000 tons of such sewage would only contain nitrogen, reckoned as ammonia, equal to that in about $16\frac{1}{2}$ cwt. of Peruvian guano, or to that in only 13 cwt. if the amount of ammonia per head per annum be reckoned at only 10 lbs. In other words, in the former case it would require 1,220 and in the latter 1,525 tons of sewage to supply the ammonia (or nitrogen reckoned as ammonia) of 1 ton of guano. In like manner, taking 80 tons of sewage per head per annum as a minimum estimate for the average sewage, inclusive of rainfall, with $12\frac{1}{2}$ lbs. of ammonia per head per annum, 1,000 tons would represent the nitrogen of $8\frac{1}{2}$ cwt., and with 10 lbs., $6\frac{1}{2}$ cwt., of Peruvian guano; or reckoning $12\frac{1}{2}$ lbs. of ammonia per head per annum, 1 ton of Peruvian guano would represent 2,440 tons, and reckoning 10 lbs., it would represent 3,050 tons.

The table also shows that reckoning $12\frac{1}{2}$ lbs. of ammonia per head per annum, the sewage of an average individual would annually represent in nitrogen $\frac{2}{3}$ cwt., or reckoning only 10 lbs. per head per annum only $\frac{1}{2}$ cwt., Peruvian guano, per head per annum.

Crops to which Sewage is most applicable.

Hitherto, on grounds shown to be fully justified, we have, for simplicity of illustration, confined attention to the amount of nitrogen or ammonia in sewage, as the measure or indication of its composition, and of the theoretical manurial value of its total solid constituents. It is, however, obviously of interest to consider whether or not the mineral or incombustible constituents of sewage exist in it in sufficient proportion to the ammonia or nitrogen, for the requirements of the crops to be grown; and, as the phosphoric acid and potassa (the one or the other, or both, according to circumstances) are, perhaps, the mineral constituents most likely to be deficient relatively to the nitrogen, their proportion to the latter in sewage, and in various crops, may appropriately be referred to in illustration of the point. Table VII shows the proportion of phosphoric acid and potassa to 100 of nitrogen in sewage, according to the mean of ten analyses of the Rugby sewage, in which the phosphoric acid and the potassa as well as the ammonia were determined. It also shows what

may be taken as approximately representing the average proportion of phosphoric acid and potassa to nitrogen in various crops.

TABLE VII.

Amount of Phosphoric Acid and Potassa to 100 Nitrogen, in Sewage and in various crops.

Rugby Sewage	Phosphoric Acid.			Potassa.		
	27			42		
	In Corn, Roots, &c.	In Straw, Leaves, &c.	In Total Produce.	In Corn, Roots, &c.	In Straw, Leaves, &c.	In Total Produce.
Meadow-Hay.....	27*	100
Clover-Hay	23	53
Wheat.....	48	42	46	28	108	57
Barley.....	40	34	38	34	126	60
Oats.....	28	37	30	25	155	65
Beans	25	46	30	32	123	50
Mangolds	17	100
Swedes	27	16	21	82	44	63
Common Turnips	28	18	26	160	71	117
Potatoes.....	42	123

It is obvious that since the phosphoric acid of sewage like the nitrogen, will be derived almost exclusively from excretal matter and food-refuse, its proportion to the nitrogen will, within comparatively narrow limits, be tolerably uniform; the amount of potassa on the other hand, will vary very much according to locality, and be considerably greater where the streets or roads are constructed of potassic minerals than elsewhere.

The table shows that, according to the analyses referred to, the Rugby sewage contained 27 parts of phosphoric acid and 42 parts of potassa, for 100 of nitrogen. It also shows that on the average, meadow hay contains almost exactly the same proportion of phosphoric acid to nitrogen as the sewage, but a much greater proportion of potassa than the latter.

In the cereal grains the proportion of phosphoric acid

* According to Baron Liebig's estimates, hay contains 51 parts of phosphoric acid to 100 of nitrogen; but having collated and averaged the results of numerous independent observers, we can see nothing to lead to the adoption of such a figure, whilst direct determinations in a number of samples of each, showed in the sewage-grass 25, and in the unsewaged 32 parts.

is, on the other hand, higher than in the sewage; in most of the other crops enumerated it is much about the same. Of potassa, the proportion is lower in the cereal (the only part of the crop which is, as a rule, sold off) than in the sewage, though in the other crops it is usually higher.

Under these various circumstances, the adequate discussion of sewage would occupy more space than it would be appropriate to their consideration here, which render it quite impossible to draw direct practical conclusions as to the application of sewage to different crops from what may appear, at first sight, the obvious indications of the figures in the table. Nevertheless, a careful consideration of the subject leads to the conclusion that, if sewage alone were applied constantly to meadow land, it would be more likely to become deficient than phosphoric acid; but that, if it were applied to the ordinary crops of rotation, phosphoric acid would be more likely to become deficient than nitrogen. Still, granting it to be clearly shown that with this or that description of soil or management, town-sewage was, in proportion to its nitrogen, deficient in this or that constituent for the cultivation of this or that crop, or crops generally, it would by no means follow that it was an inappropriate manure on that account; any defect in composition, whether in regard to phosphoric acid, potassa, or any other constituent, could be easily compensated for from other sources.

Indeed, independently of what we know of the sources of the constituents of sewage, and can judge therefrom of their appropriateness as manure for different crops, there is nothing in the results of the analysis of the solid matter of sewage, from which it could be justified in concluding that it is not applicable as manure to crops generally. On the contrary, a dry and portable solid matter, having the composition of the solid matter of town-sewage, would undoubtedly be generally applicable both to corn and other rotation crops, and to grass; and its constituents could fairly be valued by the same scale as other concentrated manures in the market.

Under the great dilution of town sewage, its large daily supply at the works, and its greater amount in wet weather when the land cannot bear, or least requires, more water, render it extremely appropriate for application on a comprehensive scale to arable land for the growth of corn and other ordinary rotation crops.

But, apart from these difficulties, if sewage can only be distributed in small quantities over large areas, at such a cost to the farmer as has yet been proposed, it is indeed vain to hope that any large proportion of the manurial constituents, derived from the consumption of human food in our towns, can be redistributed over the area from which they came; for such is the limit set by the climate to the amount of manure and of water applicable for crops that have to ripen their seed, that, for corn more especially, only comparatively small quantities per acre could be employed, and hence, were sewage systematically applied for their growth, the area of utilization must necessarily be very large. On this point it may be stated that Mr. Rawlinson, one of the members of the Royal Sewage Commission, has given it as his opinion that it would cost more to distribute 500 tons of sewage per acre, by means of pipes, hydrants, and hose and jet, as would be required in the case of application to arable land and crops generally, than to apply 5,000 tons per acre by means of open runs, as in the case of its application to grass.

From these considerations it will be obvious, that that which may be called the theoretical value of sewage, reckoned according to the constituents it contains, is not necessarily its practical available value when used in its highly diluted condition. It will be also obvious, that in that condition it is the most appropriate for grass, for which it can be employed at all seasons, and in comparatively large quantities on a limited area, and that it is least appropriate for crops which have to ripen. The question arises—what is the practical or realizable value of the constituents of sewage when they are utilized in the condition of dilution in which they exist in that fluid? This point will be illustrated by reference, both to the results of direct experiments, and to the experience of practical men who have utilized sewage with a view to profit.

Results of direct Experiment on the Utilization of Sewage.

At Rugby two fields of meadow land were experimented upon, in each one plot was left without sewage, one received sewage at the rate of 8,000 tons, one at the rate of 6,000 tons, and one at the rate of 9,000 tons, per acre per annum. The experiments were so conducted through three consecutive seasons, Table VIII summarizes the results obtained.

TABLE VIII.

ies of Sewage applied, and of Green Grass obtained, per acre per annum, in Experiments made at Rugby.

Seasons 1861, 1862, and 1863.

ns.	Plot 1. Unsewaged.	Plot 2. 3,000 Tons Sewage.	Plot 3. 6,000 Tons Sewage.	Plot 4. 9,000 Tons Sewage.
Grass obtained. Five-Acre Field.				
	Tons.cwts.qrs.lbs.	Tons.cwts.qrs.lbs.	Tons.cwts.qrs.lbs.	Tons.cwts.qrs.lbs.
1	9 5 3 5	14 16 3 8	27 1 0 10	32 16 3 8
2	8 3 1 10	27 18 0 18	34 10 0 19	32 9 2 22
3	4 18 3 13	22 5 0 11	34 18 1 27	37 0 2 5
.....	7 9 1 9	21 13 1 12	32 3 1 0	34 2 1 12
Ten-Acre Field.				
	8 18 0 15	15 16 3 2	22 15 2 12	26 13 3 12
2	16 10 0 25	27 11 0 20	32 2 1 14	31 12 1 20
3	8 0 3 19	25 5 1 8	30 11 2 12	34 19 1 21
.....	11 3 0 10	22 17 3 1	28 9 3 13	31 1 3 18
Averages :—the three years and both Fields.				
and 3	9 6 0 24	22 5 2 7	30 6 2 6	32 12 0 15

The five-acre field was much flatter than the other; its soil and subsoil were much more porous; the mechanical and chemical composition of samples, taken to the depth of 9 inches, showed the five-acre field to be much more stony, to retain much less water under the same external conditions, to contain much less organic matter, much less nitrogen, much less clay, and much more sand, than that of the ten-acre field. It was, in fact, considerably inferior in soil quality, and yielded, accordingly, considerably less produce per acre without manure. Notwithstanding this, it will be seen that it yielded upon the whole more total produce per acre under the influence of sewage than did the naturally better soil of the ten-acre field; and, it will be shown further on, that the sewage was in this case both more completely utilized and more completely converted into manure.

It would be inappropriate to discuss in detail here the influence of season and other circumstances upon the produce of the different years or the respective plots. It will be sufficient to call attention to the general character of the results, and to the practical conclusions to which they seem to lead. By the application of sewage a supply of green food was obtained much earlier and much later in the season, and the total quantity per acre was increased several fold. There was, generally, though not invariably, the more produce the greater the amount of sewage applied, the exceptions being in the wet and cold season of 1862. In the other seasons, and in both fields, there was an increase of produce with each increase in the amount of sewage applied; and the largest amounts of produce obtained at all were, in both fields, in the third season of application, and on the plots which had received the largest amounts of sewage. Still, it is important to remark, that the amounts of increase of produce for a given amount of sewage applied were the less where the larger quantities were employed. Experience abundantly shows, indeed, that if the only object were to get the largest possible amounts of produce per acre, as much as 30,000, 40,000, or even 50,000 tons of sewage might frequently be applied per acre with advantage; but under such conditions the sewage would be very inadequately both utilized and purified, and a minimum amount of increase would be obtained for a given amount of sewage applied.

Looking, however, both to urban and to rural interests, and to purification as well as utilization, much more moderate application than such as are required to yield the greatest amount of produce per acre, must be had recourse to. By way of practical suggestion on this point it may be stated that, on consideration of the circumstances under which the amounts of produce recorded in the Table were obtained, it is concluded that with an application of about 5,000 tons of average sewage per acre per annum, applied as it must be, pretty evenly throughout the year, there might be expected, taking the average of soils and seasons, an average of about 30 tons of grass. Assuming such a produce, and allowing £4 per acre for rent or natural yield, the grass would if sold at 10s. per ton, give a gross return of 0.53d. per ton of sewage employed, if for 12s. 6d. per ton 0.7d., and if for 15s. per ton 0.9d. From these amounts there would, of course, have to be deducted the cost of main distribution and application of the sewage, other expenses of the crop, and the farmer's profit, before

anything was available as payment to the town for the manurial matters.

In comparison with the result here assumed it may be observed that in the neighbourhood of Croydon, where about 250 acres are laid down for sewage irrigation, and where there are probably more than 6,000 tons of sewage annually available for each acre, from 15 to 30 tons of meadow grass, selling for from £20 to £25, are obtained per acre per annum; and after deducting as before £4 for rent, the gross return per ton of sewage employed is from 0·6d.

0·8d. With a somewhat similar application to Italian rye-grass, 10 to 35 tons, selling for from £25 to £30 are obtained, yielding, after deduction for rent or natural produce, from 0·8d. to 1d. per ton of sewage employed. It will be observed that in these cases the selling price of the grass is 16s. or 17s. per ton; but it is obvious that if sewage were extensively employed for the production of grass, its present price could not be maintained.

A marked effect of liberal sewage irrigation (indeed of active manures generally), on the mixed herbage of grass land, is greatly to develop the Gramineous plants, nearly to exclude the Leguminous, and to reduce the prevalence of miscellaneous or weedy plants, but much to encourage individual species. Among the grasses, according to locality or other circumstances, the rough meadow grass (*Poa trivialis*), couch grass (*Triticum repens*), rough cock's foot (*Dactylis glomerata*), woolly soft grass (*Holcus lanatus*), and perennial rye-grass (*Lolium perenne*), have been observed to become very prominent; two or three only remaining in any considerable proportion after some years of liberal sewage application. But sewaged produce being generally cut or grazed comparatively young, the tendency which the great luxuriance of a few very free-growing grasses has to give a coarse and stemmy later growth is not an objection, as in the case of meadows left for hay.

The chemical examination of the grass grown at Rugby showed that, at the stage of growth at which it was cut, the sewaged grass contained a less proportion of dry or solid substance than the unsewaged; that the grass cut during the later portions of the season (both unsewaged and sewaged) contained less solid matter than that cut during the more genial periods of growth; that the proportion of nitrogenous substance (and also of impure fatty or starchy matter) was much greater in the solid matter of the sewaged than in that of the unsewaged grass; that the proportion of nitrogenous substance was also much higher in the

solid matter of the grass grown towards the end than earlier in the season : that the proportion of indigestible woody-fibre was much about the same in the dry substance of the unsewaged and of the sewaged grass, but progressively diminished as the season advanced; and, lastly, that a given amount of the dry substance of grass grown in a cold and wet season, or during the cold and wet periods of the year, generally contained more nitrogenous substance than that of grass grown in more genial weather.

It will be seen presently, that, with these differences in botanical and chemical character between the unsewaged and the sewaged grass, when used as food, a given quantity of the fresh unsewaged grass was more productive of both meat and milk than an equal weight of the fresh sewaged grass; but that a given weight of the dry or solid substance of the sewaged grass was more productive than an equal weight of that of the unsewaged. Further, the less nitrogenous grass of the more genial periods of the season was more productive than the more highly nitrogenous produce of the less genial periods.

Experiments were made at Rugby with Italian rye-grass as well as meadow-grass, but the results were not sufficiently distinct in their character from those above described to render it of much interest to consider them in this place.

The next points to consider are—the comparative food-qualities of unsewaged and sewaged grass, and the best or most profitable mode of utilizing sewage-irrigated grass.

When in the experiments at Rugby the grass was cut green, and given to fattening oxen tied up under cover, more of the sewaged than of the unsewaged, reckoned in the fresh or green state, was both consumed by a given weight of animal within a given time, and required to produce a given weight of increase: but of real dry or solid substance, less of that of the sewaged than of the unsewaged grass was required to produce a given effect. When the grass was given alone the result was very unsatisfactory, but when oilcake was given in addition, the amount of increase upon a given weight of animal within a given time, and for a given amount of dry substance of food consumed, was not far short of the average result obtained when oxen are fed under cover on a good mixed diet. Still, the pecuniary result with the oxen, whether reckoned per acre or for a given amount of sewage, was by no means satisfactory.

It should here be mentioned that, at Croydon, although the land there was more liberally irrigated than at Rugby, much more satisfactory results have been obtained with fattening stock fed on the land. The practice there is, to irrigate for three or four days and nights together, to repeat the treatment two or three times for each crop, and, when the grass has got a sufficient head, to stop the application and turn the stock upon the land, where they remain until the grass is closely eaten down. They are then removed, the land is re-irrigated, and so on.

Very much better results were obtained at Rugby when the grass was given to milking cows. Referring to the Report of the Sewage Commission for all further details, the summary of the results with cows given in Table IX. will suffice for consideration here.

It may be stated generally, that when the cows were fed on grass alone, as much as they chose to eat, a given weight of the animal was more productive, both of milk and increase, but especially of milk, on the unsewaged than on the sewaged grass. More milk was also produced from a given weight of the unsewaged grass, reckoned in the fresh or green state, than from an equal weight of the fresh sewaged grass. Of dry or solid substance, however, a given weight of that of the sewaged grass produced, on the average, more milk than an equal weight of that of the unsewaged.

The milk from the cows fed on the sewaged grass was, upon the whole, slightly the less rich, containing generally somewhat less casein, butter, sugar, and total solid matter (though more mineral matter) than that from the unsewaged; but when oilcake was given with the grass, whether sewaged or unsewaged, the richness of the milk was notably increased.

The productive quality of the grass was very different in different seasons, and at different periods of the same season, being very inferior in the wet and cold season of 1862, and towards the close as compared with the earlier periods of the seasons.

Without commenting further on the difference of result obtained under different conditions of season, or under other varying circumstances, it will be sufficient briefly to call attention to the more general results which the records in the table bring prominently to view, and to the practical conclusion which, on a careful consideration of all the circumstances and details, may seem to be safely deducible from them.

TABLE IX.

Results obtained at Rugby, with Cows fed on Unsewaged and Sewaged Grass, 1861, 1862, and 1863.

	Plot 1. Unsewaged	Plot 2. 3,000 Tons Sewage.	Plot 3. 6,000 Tons Sewage.	Plot 4. 9,000 Tons Sewage.
Time each acre (with oilcake, if any) would keep 1 cow :—				
	Weeks.	Weeks.	Weeks.	Weeks.
1861—Grass (alone)	19	41	59	69
1862—Grass (with oilcake).....	42	63	73	72
1863— { Grass ($\frac{1}{4}$ without, $\frac{1}{4}$ with } { oilcake)	22	48	67	73
Means...	28	51	66	71
Milk from the Produce of each acre (exclusive of oilcake,* if any) :—				
	Gallons.	Gallons.	Gallons.	Gallons.
1861—Grass (alone)	321	571	820	961
1862—Grass (with oilcake).....	613	835	973	958
1863— { Grass ($\frac{1}{4}$ without, $\frac{1}{4}$ with } { oilcake)	414	876	1207	1327
Means...	449	761	1000	1082
Value of milk from the produce of each acre (ex. of oilcake,* if any), at 8d. per gall. ■				
	£ s. d.	£ s. d.	£ s. d.	£ s. ■
1861—Grass (alone)	10 14 3	19 0 6	27 6 11	32 0 1
1862—Grass (with oilcake).....	20 8 10	27 16 10	32 8 11	31 18 1
1863— { Grass ($\frac{1}{4}$ without, $\frac{1}{4}$ with } { oilcake)	13 16 0	29 3 9	40 4 7	44 4
Means...	14 19 8	25 7 0	33 6 10	36 1
Increased produce of milk per 1,000 tons sewage applied (ex. of oilcake,* if any) :—				
		Gallons.	Gallons.	Gallon. ■
1861—Grass (alone)		180	178	151
1862—Grass (with oilcake).....		74	60	38
1863— { Grass ($\frac{1}{4}$ without, $\frac{1}{4}$ with } { oilcake)		154	132	101
Means...		136	123	97
Increased value of milk (at 8d. per gall.) per 1,000 tons sewage applied (ex. of oilcake,* if any) :—				
		£ s. d.	£ s. d.	£ s. ■
1861—Grass (alone)		5 19 10	5 18 8	5 0 ■
1862—Grass (with oilcake).....		2 9 4	2 0 0	1 5
1863— { Grass ($\frac{1}{4}$ without, $\frac{1}{4}$ with } { oilcake)		5 2 7	4 8 1	3 7
Means...		4 10 7	4 2 3	3 4

* The value of the milk, "exclusive of oilcake," is reckoned by deducting cost of the cake consumed, less the estimated value of the manure it yields, from ■

It is seen that whether we reckon the total amount of food yielded per acre, or the amount, or the value, of the milk obtained from the consumption of the produce of each acre, there was a very great increase, varying from two to three-fold, according to season, by the use of sewage. The land upon which these experiments were made was good feeding pasture, of probably more than average quality, and the natural yield, without sewage, was, therefore, correspondingly high. Taking into consideration this fact, and other circumstances under which the results were obtained, it is concluded that, if not larger amounts of total produce per acre, at any rate larger amounts of increase for a given quantity of sewage may be expected when it is applied systematically over large tracts of land, with a view to the production of grass and milk.

It is estimated that with 5,000 tons of sewage per acre per annum, judiciously applied to Italian rye-grass or meadow-land properly laid down to receive it, an average *gross* produce of not less, and perhaps more, than 1,000 gallons of milk per acre per annum might be anticipated; and it may be observed that 1,000 gallons of milk at 8d. per gallon would represent a *gross* money return of £93 6s. 8d.

Putting the result in another way it may be stated that it required, according to circumstances, the consumption of between 5 and 6 tons of grass for the production of 1 ton of milk; and if we reckon 6 parts of grass for 1 of milk, and 30 tons of grass per acre, this would give a *gross* return in value of milk at 8d. per gallon of something over £37 per acre, or of about 25s. per ton, of grass consumed.

Still another illustration of the important bearing of the question of the utilization of the sewage of our town populations upon the re-production of food may be given. Supposing the whole of the sewage of a given population (which, however, would seldom be the case) were applied exclusively for the growth of grass for the production of milk, the result would be an increased yield of about $2\frac{1}{2}$ pints of milk per week, or about $\frac{1}{2}$ lb. per day, per head of such population. So far as the sewage were so applied, a portion of the milk produced would, of course, be represented, in con-

gross value inclusive of oilcake; and the amount of milk, "exclusive of oil-cake," by deducting from the gross amount of milk with oilcake at the rate of one gallon for every 8d. of deducted value. Such estimates are, however, obviously only approximations to the truth.

sumption, by its equivalent in butter and cheese. A portion of the grass would, however, be used directly for the production of meat ; and, in addition to the milk and meat produced by the consumption of the grass, a large amount of solid manure would be obtained, which would be applicable to arable land for the growth of corn and other rotation crops.

It would appear, then, that if town sewage were to a great extent utilized by the application of something like 5,000 tons per acre per annum to Italian rye-grass and meadow-land, a direct result would be a very greatly increased production of important articles of human food which are at present both scarce and dear. But the question remains—would the sewage, by such an application, be sufficiently purified to allow of the drainage from the irrigated land being turned into rivers which are to be used as a water-supply for other towns? Some light will be thrown on this subject by the results next to be considered.

In order to determine how far, in the experiments at Rugby, the sewage was deprived of its manurial or putrescible constituents in its passage over and through the land, samples of the drainage water were collected for analysis in each field, simultaneously with those of the sewage, commencing in May, 1862, and ending in October, 1863. In all 62 partial analyses of drainage-water, corresponding in detail with those of the sewage, were made. A few other analyses, in much more detail, were made of the sewage and drainage of the season of 1864. The results of the large number of partial analyses are summarized in Table X, which shows, in parallel columns, the average composition of corresponding samples of sewage and drainage.

TABLE X.

Mean Composition of the Rugby Sewage before application, and of the Drainage-water from the Irrigated Land, in the Seasons 1862 and 1863.

Grains per Gallon.

Constituents.		Five-Acre Field.		Ten-Acre Field.		The two Fields.		
		Sewage.	Drainage.	Sewage.	Drainage.	Sewage.	Drainage.	
Season 1862; May—October, both inclusive.								
In suspension	{	Inorganic	11 Samples. 25·67 14·69	8 Samples. 1·81 1·40	11 Samples. 24·69 17·14	11 Samples. 3·74 1·39	22 Samples. 25·25 15·92	19 Samples. 2·92 1·39
		Total ..	40·36	3·21	42·03	5·13	41·20	4·31
In solution..	{	Inorganic	34·49 7·83	34·50 7·18	32·38 7·60	37·10 7·83	33·44 7·71	36·01 7·56
		Total ..	42·32	41·68	39·98	44·93	41·15	43·57
		Total inorganic..	60·16	36·31	57·27	40·84	58·72	38·93
		Total organic ..	22·52	8·58	24·74	9·22	23·63	8·95
		Total solid matter ..	82·68	44·89	82·01	50·06	82·35	47·88
Ammonia	{	In suspension	1·37	0·24	1·52	0·33	1·44	0·39
		In solution ..	4·13	0·80	4·26	1·85	4·20	1·41
		Total ..	5·50	1·04	5·78	2·18	5·64	1·70
Season 1863; November, 1862—October, 1863, both inclusive.								
In suspension	{	Inorganic	23 Samples. 39·41 27·35	21 Samples. 2·14 1·41	22 Samples. 34·93 25·99	22 Samples. 3·93 3·29	45 Samples. 37·22 26·69	43 Samples. 3·06 2·37
		Total ..	66·76	3·55	60·92	7·22	63·91	5·43
In solution ..	{	Inorganic	39·57 8·35	33·55 7·46	38·77 8·30	41·35 7·98	39·18 8·32	39·98 7·73
		Total ..	47·92	46·01	47·07	49·33	47·50	47·71
		Total inorganic..	78·98	40·69	73·70	45·28	76·40	43·04
		Total organic ..	35·70	8·87	34·29	11·27	35·01	10·10
		Total solid matter ..	114·68	49·56	107·99	56·55	111·41	53·14
Ammonia	{	In suspension	2·08	0·15	1·98	0·31	2·03	0·23
		In solution ..	5·83	0·69	5·69	1·85	5·76	1·28
		Total ..	7·91	0·84	7·67	2·16	7·79	1·51

It is seen that of matter in suspension in the sewage, nearly the whole, both inorganic or organic, was retained by the soil; and probably a considerable part of the little which the drainage-water contained was derived from the soil itself.

Of matter in solution, on the other hand, a gallon of the

drainage-water contained, on the average, much about the same amount, both inorganic and organic, as a gallon of the sewage; though, doubtless, a considerable portion of the soluble matters in the drainage had their immediate source in the soil—the sewage giving up valuable manurial matters to the soil, and the fluid in its turn taking up substances from it.

It is important to remark that the drainage from the more porous and less naturally fertile soil of the five-acre field (which, however, gave the largest amount of increase for a given amount of sewage), contained less of almost every constituent, or class of constituents, enumerated, than did that from the more argillaceous and more naturally fertile soil of the more steeply sloping ten-acre field. The result is particularly marked in the case of the ammonia. The fact here indicated is of considerable practical, as well as scientific, interest: and it is perfectly consistent with the results of common experience, which tend to show that a soil which may contain a comparatively small proportion of clay, but which is thoroughly porous, is, as a rule, much better adapted for sewage irrigation, both as regards the utilization and the purification of the sewage, than one which, though richer in clay and of higher natural quality, is but imperfectly permeable by the fluid.

The results given in Table XI show in more detail the changes in the composition of the fluid in its passage through the soil. They relate to samples of sewage and drainage taken in another field at Rugby, during very dry weather, in the summer of 1864. The plan of collection was, to take of sewage about a gallon, and of drainage about half a gallon, eight or ten times during the ten or twelve working hours of the day; at the end of the day, after well shaking, to take a gallon from each mixture; and to repeat this for six consecutive days, until six gallons of each were obtained, when after well shaking, a two-gallon sample of each was bottled off for the purposes of analysis.

TABLE XI.

Detailed Composition of samples of the Rugby Sewage before application, and of the Drainage-water from the irrigated land, collected July, 1864.

Constituents.		Grains per Gallon.			
		Collected July 6—11.		Collected July 13—18.	
In suspension.	Inorganic matter :—	Sewage.	Drainage.	Sewage.	Drainage.
	Oxide of iron and alumina	4.57	...	6.30	...
	Lime	4.48	...	3.75	...
	Magnesia	0.65	...	0.25	...
	Carbonic acid	3.25	...	2.17	...
	Phosphoric acid	1.84	...	1.14	...
	Silica, sand, &c.	31.60	...	39.30	...
	Total	46.39	...	52.91	...
	Organic matter.....	40.40	...	32.40	...
	Total matter in suspension...	86.79	...	85.31	...
In solution.	Inorganic matter :—				
	Oxide of iron, &c.	Traces.	...	1.25	0.25
	Lime	8.45	10.25	8.23	10.08
	Magnesia	1.76	1.69	1.80	1.69
	Soda (1)	5.46	0.38	5.24	2.30
	Chloride of sodium (1)	6.82	9.73	8.53	9.21
	Chloride of Potassium (1)	6.08	1.50	6.17	2.34
	Sulphuric acid	4.39	6.55	4.01	6.75
	Phosphoric acid	1.28	0.44	1.66	0.32
	Carbonic acid.....	8.83	6.18	7.42	7.01
	Silica	1.80	0.80	1.00	0.80
	Total	44.87	37.52	45.31	40.75
	Organic matter.....	11.20	7.80	10.00	7.05
	Total matter in solution.....	56.07	45.32	55.31	47.80
Total inorganic matter		91.26	37.52	98.22	40.75
Total organic matter (2)		51.60	7.80	42.40	7.05
Total solid matter.....		142.86	45.32	140.62	47.80
(1) Containing	Potassa	3.84	0.94	3.90	1.48
	Soda	9.07	5.54	9.76	7.17
	Chlorine	7.03	6.61	8.10	6.70
(2) Containing	Ammonia	In suspension	2.42	...
		In solution	0.98	6.36	0.92
		Total.....	8.66	8.78	0.92
	Nitric acid in solution = Ammonia	 (3) 1.33	...	(4) 1.41

(3) 4.227 Nitric acid = 1.096 Nitrogen = 1.331 Ammonia.

(4) 4.483 „ = 1.162 „ = 1.411 „

The soil was light and gravelly, with a gravelly subsoil. An examination of the figures in Table XI shows, that, done the work of absorption, at any rate as well as, if not than, on the average, did the soils in the other fields. I intended to take samples for detailed analysis from this field under various conditions of the weather, but owing to the continuance of the drought, this could not be accomplished.

In judging of these results, as well as those already considered, it must, of course be borne in mind that, excepting when the land is already saturated with water, a gallon of drainage represents much more than a gallon of sewage; and that, the amount of any constituent of the sewage found in a gallon of the drainage must have been derived from more than a gallon of the former. The non-retention of valuable manurial matter by the soil was, therefore, not so great as would at first sight appear. On an inspection of the comparative composition of equal volumes of the sewage and of the drainage.

As in the larger number of cases, so in these, the quantity of matter in suspension in the drainage was very small, and obviously in great part derived from the soil, it was not suitable for quantitative analysis. A considerable proportion of the phosphoric acid of the sewage was in suspension, but there was very little of it in suspension in the drainage, the whole of the phosphoric acid existing in the sewage having been retained by the soil.

It is satisfactory to observe that among the inorganic constituents in solution in the sewage, by far the larger proportion of those which are, perhaps, the most likely to become relatively deficient, was retained by the soil. Thus, smaller proportions of boron, potassa and the phosphoric acid of the sewage passed off in the drainage than of any other constituents. Soda was also retained by the soil to a considerable extent, magnesia in a less degree, lime less still. Of lime, indeed, there was more in a gallon of drainage than in a gallon of sewage; of sulphuric acid also there was considerably more in the drainage than in an equal volume of the sewage. Lastly, of soluble silica a notable portion passed in the drainage.

Of organic matter in solution a very considerable quantity was found in the drainage-water. The character of the soluble organic matter in the drainage is, however, very different from that in the sewage. It contains very much less ammonia, or ammonifying matter; and, especially in periods of active vegetation

doubtless, frequently be derived from vegetable matter within the soil, rather than directly from the sewage.

A very important point to remark is, that, whilst the sewage scarcely contained an appreciable amount of nitric acid, the drainage contained more nitrogen in that form than as ammonia; the result being that the soil had retained a considerably less proportion of that important manurial constituent of the sewage than would have been supposed had only the more partial analyses been made.

The general result was, that, practically, the whole of the insoluble or suspended matter of the sewage was retained by the soil; and that, of the constituents of the sewage, whether in suspension or in solution, those which are of the most value, because the most liable to become relatively exhausted, were the most efficiently retained. Nevertheless, the drainage-water still retained so much of potassa, phosphoric acid, ammonia, and nitric acid, as clearly to show that the sewage had not been perfectly deprived of its valuable manurial matters, and also so much of total soluble matter, especially of soluble organic matter, as to show that it had not been by any means perfectly purified.

There is, indeed, a limit to the power which a soil possesses of removing substances from solution, or of preventing those already absorbed from being dissolved in water passing through it, the result being dependent on the physical and chemical characters of the soil itself, and on the amount and composition of the fluid passing through it. So far as the soluble organic matters of the drainage are derived from vegetable matter within the soil, it is a question whether there will not always be a considerable amount in that passing from land covered with luxuriant vegetation. So far, however, as the nitrogen of the drainage exists in the form of nitric acid, it is a pretty satisfactory indication that the organic matter has, to a great extent, already passed the stage of deleterious putrescence.

In the Rugby experiments the arrangements were not such as to allow of the water drained from one portion of the land being passed over another; but at Beddington, near Croydon, a great portion of the water does duty twice, and sometimes three times; and from results kindly communicated by Mr. Latham, the engineer to the Croydon Board of Health, and given in the following table, it would appear that there the water eventually passes from the land in a state of much greater purity than was the case in the Rugby experiments.

TABLE XII.

Partial Analyses of the Croydon Sewage before application, of the Drainage-water from the irrigated land, and of the River Wandle, above and below the Drainage Outfall from the irrigated land.

Constituents.	Croydon.		River Wandle.	
	Sewage.	Drainage.	Above Drainage Outfall.	Below Drainage Outfall.
	Grains per gallon.			
Inorganic matter.....	48.30	23.40	18.56	20.16
Organic matter	52.20	2.40	1.44	2.08
Total solid matter ...	100.50	25.80	20.00	22.24
Ammonia	6.70	0.21	0.18	0.18

The figures show much about the same amount of ammonia in the Sewage of Croydon, as was found on the average in that of Rugby; but the amount in the Croydon drainage was extremely small. It is unfortunate that the quantity of nitric acid was not also determined; but we are informed that it undoubtedly exists in some amount in the drainage from the Beddington meadows. Still, although formerly the Croydon Board had to meet numerous law-suits on account of the pollution of the river by the sewage, the fluid is now so far purified before being discharged, that those having the right of fishing in the river have found it worth while to fix gratings to prevent the fish going up the main outfall from the sewage-irrigated land.

The results obtained in regard to this part of the subject—that of purification—however interesting and important, must still be looked upon as little more than initiative; but there can be no doubt that, when large quantities of sewage are applied to grass-land, the arrangements should be such as to allow of the drainage-water being collected and re-used in such a manner as to insure as far as possible both complete utilization and complete purification. It must be admitted, however, that further experience, and further investigation, are still wanting, to determine what amount of sewage, provided the drainage water be properly re-distributed, can be

safely applied to a given area, under different conditions of soil and subsoil, and under different conditions of season, so as to insure its sufficient purification.

Experience of Common Practice in the Utilization of Sewage.

Leaving the results of experimental inquiry, it will be well briefly to notice those of practical experience hitherto, in regard to the value and utilization of town sewage. The instance most frequently quoted is that of the neighbourhood of Edinburgh, relating to which some particulars are given in the following Table:—

TABLE XIII.

Relating to the Sewage-irrigated Meadows near Edinburgh.

Names of Meadows.	Imperial Acres under Irrigation.	Approximate Population contributing to each Acre.	Approximate Quantity of Sewage available for each Acre.
Lochend, Spring Gardens, and Craighentenny.....	285	337	Tons. 20,500
Roseburn and Western Dalry	80	112	17,000
Quarry Holes	8	562	65,000
Broughton Burn.....	6	1,666	102,000
The Grange	16½	302	97,000

These tabular statements are chiefly based upon direct information, obtained in part from Mr. McPherson, the Edinburgh City Surveyor, and in part from the occupiers or managers of the respective meadows. To prevent misunderstanding, however, it must be explained with regard to them, that, as water-closets are not universal, and as the sewage is frequently allowed to pass unused, the records of the amount of population contributing to, and of sewage available for, each acre, do not show the amounts actually utilized, but only approximately the total amounts available, whether used or wasted.

Sewage has been applied to some portions of the land in the neighbourhood of Edinburgh for about 200 years, to a considerable portion for more than 60, and to most of that now under irrigation for more than 30 years. In two instances arrangements have been made for raising the sewage, by pumping, an inconsiderable

number of feet; but the cost has been found too great to allow a sufficient quantity being applied per acre, and hence the application in this way has been much limited, if not on some portions of the land entirely abandoned. The application is confined to meadow land and Italian rye-grass, and the distribution is entirely by means of open runs. When Italian rye-grass is grown, the land is periodically broken up, and one or two other crops taken without sewage before laying down again to grass. The application to ordinary rotation crops on arable land forms no part of the system adopted.

There is no doubt that at Edinburgh larger amounts of sewage are applied per acre than anywhere else, and that it is under the conditions that there are there obtained larger amounts of produce per acre than anywhere else. Nor is there any doubt, on the other hand, that there is, at Edinburgh, not only very great waste of manurial constituents, but very imperfect purification of the sewage. Hence the experience there, however interesting and important in some points of view, cannot be taken as the foundation either of estimates of the value realizable in practice by utilization of given amounts of sewage, or of the sewage of given population, or of safe conclusions as to the amount of sewage that can advantageously be applied per acre when the drainage has to be passed into a river, which may have to serve as the water-supply of other towns, instead of, as at Edinburgh, having an immediate outfall into the sea.

It may be mentioned that generally four or five crops of grass are obtained per acre annually, amounting, according to circumstances, to 30, 40, 50, 60, and even more tons per Imperial acre and selling for prices varying from £8 to over £40 per acre, averaging perhaps about £25. These results are, indeed, sufficiently striking, and well merit careful inquiry and consideration; but, for the reasons above stated, the exact practice of Edinburgh is not applicable to towns generally, and is especially inapplicable to inland towns.

Table XIV summarizes the results of the experience of the most important instances of sewage utilization in other localities.

TABLE XIV.

Relating to Sewage-irrigation in various localities.

Towns.	Population contributing.	Acres.		Crops, &c.	Annual Payment to Towns
		Original	Reduced.		
Alnwick	6,500	270	0	Arable and grass; abandoned	Nothing
Carlisle	22,000	70	...	Meadow grass; all grazed	?
Croydon	16,000	250	...	Meadow and rye-grass	£300
Exeter	4,000	50	...	Grass	Nothing
Leamington	6,700	{ 190 280	{ 20 100	Meadow and rye-grass Meadow; chiefly grazed	{ £50
London	6,000	95	...	Grass	Nothing
Northampton	4,000	210	{ 7 35	Rye-grass—Summer. Meadow grass—Winter	{ £10
Reading	7,000	42	...	Grass; not yet at work	Nothing

At Alnwick, the late Duke of Northumberland put down machinery and piping for the distribution of the sewage of the town over about 270 acres of mixed arable and grass land. After a very short time, the tenants, who had the free use of the sewage at the cost of its application, abandoned it altogether; and the Surveyor of the District, who reports the failure, expresses his opinion strongly against the general applicability of sewage to arable land.

At Carlisle, the sewage of only a portion of the town is utilized. It is deodorized by Mr. McDougall's disinfecting fluid, and raised by steam power some 10 or 12 feet into an open cut, from which it is diverted for application to the land by moveable iron troughs. It is estimated that from 8,000 to 9,000 tons of sewage are applied per acre per annum. It is understood that little or no benefit is realized by the town; but that the tenant makes a considerable profit by sub-letting the sewage-irrigated land for agricultural purposes.

In the Neighbourhood of Croydon, as already referred to, the sewage of nearly 20,000 persons is applied to about 250 acres of meadow and Italian rye-grass. It is calculated that more than 100 tons of sewage are available for each acre. A considerable quantity of the fluid is used two or three times over; and it finally

passes from the land pretty satisfactorily purified. It is estimated that, after making deduction of £4 for rental, the gross return per ton of sewage applied is, at the present prices of the produce, with Italian rye-grass from $\frac{3}{4}$ d. to 1d., and with meadow grass from $\frac{1}{2}$ d. to $\frac{3}{4}$ d. The sewage is not applied in any systematic manner to other crops, but it has been tried on a small scale on root-crops. An enlargement of the area of irrigation is contemplated, which will, if carried out, somewhat reduce the amount of fluid and excretal matters available per acre below the quantities above stated.

About 12 years ago, arrangements were made for collecting the sewage of Rugby in a tank, from which it is pumped, by a 12-horse power engine, through iron pipes laid down for the distribution over about 470 acres of mixed arable and grass land. Up to last year 190 acres were held by Mr. James Archibald Campbell but he has gradually limited the area of application, and during the last few years has abandoned the use of hose and jet, excepting occasionally on a small scale, and confined the application almost exclusively to from 12 to 20 acres of meadow and Italian rye grass. The remainder of the land, amounting to about 280 acres has passed through the hands of two tenants, both of whom are said to have sustained considerable loss. The last of the two has confined the application almost exclusively to about 100 acres of grass land, and applied the sewage almost entirely by open run. The whole is now in the hands of the landlord, Mr. G. H. Walker, who, it is understood, is contemplating the abandonment of the use of steam power, pipes, and hose and jet, and the application on a limited area by means of gravitation.

The general result at Rugby is, then, that after about a dozen years of practical experience, with arrangements adapted for the application of small quantities of sewage per acre, to arable as well as to grass land, and to all crops, the area has been greatly limited, the use to any other crops than meadow and Italian rye grass is quite exceptional, and the application by means of steam power, pipes, and hose and jet, will probably soon be entirely abandoned. It may be added that, at the time of the experiments of the Commission, the sewage, which was considerably stronger than that of the Metropolis, cost the tenants only about $\frac{3}{4}$ d. a ton at the hydrants in the fields; yet, rather than incur the loss of using it at that cost, both were glad to get rid of it to the Commission, at rates which, though three times as high during

six summer as during the six winter months, averaged the year round scarcely, but very nearly, 1d. per ton at the hydrants.

Some years ago, the Earl of Essex laid down pipes for the application of the sewage of Watford, by pumping and hose and jet, to about 210 acres of mixed arable and grass land. The results which his Lordship obtained on the application of only 134 tons of sewage per acre to wheat have frequently been held to be conclusive proof of its applicability in small quantities per acre over large areas, to arable land, and to all crops. But in the evidence given by his Lordship before the Sewage Committee of 1862, he stated, very emphatically, that his great error had been the piping of too much land; that he required 5,000 tons per acre for 10 acres of rye-grass; and that, applying the remainder to 35 acres of meadow, he had none to spare for wheat. In other words, although the abandonment of one acre of rye-grass would set free sewage enough for nearly 40 acres of wheat, if applied only at the rate which yielded the large gross return per ton of sewage so frequently quoted, yet his Lordship's practical experience had led him to prefer the application to the one acre of rye-grass rather than to the nearly 40 acres of wheat. Further, his Lordship gave it as his opinion that sewage would not be profitable to the farmer unless he could have it at from $\frac{1}{2}$ d. to $\frac{3}{4}$ d. per ton.

Referring to the question of the application of sewage to corn crops, it may be stated that, in an experiment made by the Commission at Rugby, with oats, a very high gross money return per ton of sewage was also obtained. The experiment was made in the unusually productive season of 1868, and with sewage of about double the average strength of that of the Metropolis, applied during a period of very dry weather. The results were, therefore, quite exceptional, and cannot be taken as affording any indication of what might be expected from the application of small quantities of sewage to corn crops generally, on different soils, and on the average of seasons. There cannot, indeed, be a doubt, that to obtain a maximum gross value of produce from a given amount of sewage, it should be applied in small quantities per acre, and in dry weather. But sewage is produced in large daily amount at all seasons, and must be disposed of as soon as it is produced. It must, therefore, be applied in winter, when of comparatively little value, as well as in summer, when of more, and it would frequently be quite inapplicable to arable land. Moreover, to obtain an increased gross money return per ton of sewage by using it on a

comprehensive scale for corn and other ordinary rotation crops, would involve the extra cost of main distribution over at least a ten-fold, if not frequently a twenty-fold area, and require the application to a great extent by the expensive means of pipes and hose and jet, instead of by the economical one of open runs.

At Malvern and Tavistock the application of sewage to grass land has now been carried on for some years, but at Worthing it has only very recently been commenced.

From this short review of the experience of practical men who have undertaken the utilization of sewage with a view to profit, it appears that, wherever arrangements have been made for the application of small quantities over large areas, to corn and other rotation crops on arable land, and by means of pipes and hose and jet, the undertaking has either been entirely abandoned, or the area greatly limited, and the application confined almost exclusively to meadow and Italian rye-grass. On the other hand, the undertakings which have been the most successful from the agricultural point of view are those in which the arrangements have been adapted for the almost exclusive application to grass, and the application to other crops is only exceptional.

General Practical Conclusions.

The practical conclusions deducible from the whole inquiry may be briefly stated as follows:—

1. It is only by a liberal use of water that the refuse matters of large populations can be removed from their dwellings without nuisance and injury to health.

2. That the discharge of town sewage into rivers renders them unfit as a water supply to other towns, is destructive of their fish, causes deposits which injure their channels, gives rise to emanations which are injurious to health, is a great waste of manurial matter, and should not be permitted.

3. That the proper mode of both utilizing and purifying sewage is to apply it to land.

4. That, considering the great dilution of town sewage, its constant daily supply at all seasons, its greater amount in wet weather when the land can least bear, or least requires, more water, and the cost of distribution, it is best fitted for application to grass, which alone can receive it the year round. It may, however, be

sionally applied with advantage to other crops within easy reach of the line or area laid down for the continuous application of grass.

i. That, having regard both to urban and rural interests, an application of about 5,000 tons of sewage per acre per annum, to meadow Italian rye-grass, would probably, in the majority of cases, prove to be the most profitable mode of utilization, though the quantity would have to be reduced, provided experience showed that the water was not sufficiently purified ; and it is pretty certain that a farmer would not pay $\frac{3}{4}$ d., and it is even very doubtful whether he could afford to pay $\frac{1}{2}$ d. per ton, the year round, for sewage of the average strength of that of the Metropolis (excluding storm-water) delivered on his land.

ii. That the direct result of the general application of town sewage to grass land would be an enormous increase in the production of milk, butter, cheese, and meat ; whilst, by the consumption of the grass, a large amount of solid manure, applicable to arable land and to crops generally, would be produced.

iii. That the cost or profit to a town of arrangements for the removal and utilization of its sewage must vary very greatly, according to its position, and to the character and levels of the land to be irrigated. Where the sewage can be conveyed by gravitation, and a sufficient tract of suitable land is available, the town may realize a profit ; but, under contrary conditions, it may have to submit to a pecuniary sacrifice to secure the necessary sanitary advantages.



FOOD IN ITS RELATIONS

TO

VARIOUS EXIGENCIES OF THE ANIMAL BODY.

THE appearance in the June (Supplementary) Number of the Philosophical Magazine of the interesting paper by Professors Fick and Wislicenus "On the Origin of Muscular Power," and the further interest excited in the subject by Professor Frankland's recent lecture at the Royal Institution, seem to render it opportune that the important question of the connexion between certain constituents of food and certain exigencies of the animal body should receive a little further consideration at the present time. Professor Frankland truly said that, since the appearance of Baron Liebig's masterly and highly suggestive work 'On Organic Chemistry in its applications to Physiology and Pathology' in 1842, his views of the relation of the nitrogenous and the non-nitrogenous constituents of food to certain requirements of the animal organism have been pretty generally adopted by text-book writers. It is also true, that authorities on the subject of the chemistry of food have, even so recently as last year and this, either directly maintained or taken for granted the correctness of Baron Liebig's views. It is, however, not the case, as was also assumed by Professor Frankland, that those views have remained unquestioned excepting in the one or two instances of criticism to which he referred.

This question, in various aspects, has occupied a great deal of our own attention for many years past; and so long ago as 1852, we advocated substantially the views now adopted by Professors Fick, Wislicenus, and Frankland; and we have on various occasions since that date expressed them with greater definiteness, and urged them the more emphatically, as new experimental evidence either of others or ourselves seemed to lend them support or confirmation. It may be well, therefore, to state very briefly

2 Messrs. J. B. Lawes and J. H. Gilbert on *Food in its*

the course of our own investigations bearing upon the subject and also the conclusions that we have based upon them.

In Baron Liebig's work above alluded to, and also in subsequent publications, he treated of the food requirements of the animal body generally—that is, under different conditions; starting from the fundamental assumptions, on the one hand, the direct connexion of the nitrogenous or, as he designated them, the “*plastic*” constituents of food, not only with the formation in the animal body of the compounds containing nitrogen but also with the development of muscular power, and, on the other, of the general relationship of the non-nitrogenous constituents of food with respiration, the development of heat, and the deposition of animal fat, he concluded that the relative value of different foods, as such, was to a great extent dependent on, and even measurable by, the proportion of nitrogenous constituents which they contained.

It was upon the assumption of the broad and fundamental classification of the constituents of food according to their various offices in the animal economy, as above stated, that numerous analyses of food were undertaken, and, founded upon the results obtained, Tables constructed professing to arrange current articles of food, both of man and other animals, according to their comparative values as such; and whether the object were the feeding of animals for the production of milk, the so-called fattening of them for the production of meat, or the support of the body by the exercise of muscular power, the proportion of nitrogenous constituents was generally taken as the measure of that value.

Omitting, for the sake of brevity, any special reference to the labours or views of others, it will suffice here to make a few such quotations from Baron Liebig's works as will best convey shortly in his own words a pretty clear indication of his own views and at the same time pretty fairly represent those of a large proportion both of systematic writers and experimenters, on the points in question. Speaking of the nitrogenous constituents of food, he said:—

“It is found that animals require for their support less of a vegetable food in proportion as it is richer in these peculiar matters, and cannot be nourished by vegetables in which these matters are absent.” (Chemical Letters, 3rd edition, p. 349.)

Again:—

“The admirable experiments of Boussingault prove, that the increase in the weight of the body in the fattening or feeding of stock (just as is the case with the supply of milk obtained from milch cows), is in proportion to the amount of plastic constituents in the daily supply of fodder.” (Chemical Letters, 3rd edit. p. 36)

In regard to the exercise of force, he said:—

"As an immediate effect of the manifestation of mechanical force, we see, that a part of the muscular substance loses its vital properties, its character of life; that this portion separates from the living part, and loses its capacity of growth and its power of resistance. We find that this change of properties is accompanied by the entrance of a foreign body (oxygen) into the composition of the muscular fibre (just as the acid lost its chemical character by combining with zinc); and all experience proves, that this conversion of living muscular fibre into compounds destitute of vitality is accelerated or retarded according to the amount of force employed to produce motion. Nay, it may safely be affirmed, that they are mutually proportional; that a rapid transformation of muscular fibre, or, as it may be called, a rapid change of matter, determines a greater amount of mechanical force; and conversely, that a greater amount of mechanical motion (of mechanical force expended in motion) determines a more rapid change of matter." (*Organic Chemistry in its application to Physiology and Pathology*, 1842, pp. 220 & 221.)

And again:—

"The amount of azotized food necessary to restore the equilibrium between waste and supply is directly proportional to the amount of tissues metamorphosed.

"The amount of living matter, which in the body loses the condition of life, is, in equal temperatures, directly proportional to the mechanical effects produced in a given time.

"The amount of tissue metamorphosed in a given time may be measured by the quantity of nitrogen in the urine.

"The sum of the mechanical effects produced in two individuals, in the same temperature, is proportional to the amount of nitrogen in their urine; whether the mechanical force has been employed in voluntary or involuntary motions, whether it has been consumed by the limbs or by the heart and other viscera." (*Ibid.* p. 245.)

Our own direct experiments have had reference chiefly to the feeding of fattening animals; but the characteristic food requirements of the body, when fed with the view to the exercise of muscular power, have also been made the subjects of enquiry.

Referring to the feeding of fattening animals, the question arises, whether in the use of the currently adopted food-stuffs the amount of food consumed by a given weight of animal within a given time, and the amount of increase produced, are more influenced by the amount of the nitrogenous or of the non-nitrogenous constituents which the food supplies; that is to say, whether the sum of the requirements of the animal system under these circumstances is such that, in the use of the ordinary articles of food, the amount taken or increase produced will

be more regulated, or measurable, by the supplies of the nitrogenous or "flesh-forming" constituents, or by those of the more specially respiratory and fat-forming non-nitrogenous constituents.

To acquire the data necessary for the satisfactory solution of this question, some hundreds of animals—oxen, sheep, and pigs—have been experimented upon. Comparative lots being selected the general plan of the feeding-experiments was to give to some a fixed and limited amount of food of known composition in regard to its contents of nitrogenous and non-nitrogenous constituents, to others a fixed and limited amount of food of different composition in this respect, and to allow all to take as much as they chose to eat of some other food, also of known composition, the quantity consumed being weighed. In some cases a single description of food only, or a mixture of several descriptions in known proportions, was given *ad libitum*, but weighed; and in others, several descriptions of food were allowed, each separately, *ad libitum*, but weighed. It will be seen that in this way great variation in the amount and proportion of the nitrogenous and non-nitrogenous constituents supplied was attained, whilst the animals, according to the nature of the food within their reach, fixed for themselves the limit of their consumption. All such comparative experiments were conducted for many weeks, or even for several months, consecutively, and the weights of the animals themselves were determined at the commencement, at stated periods during the progress, and at the conclusion of the experiment.*

To determine the character and composition of the gross increase in live-weight, the weights of the individual internal organs and of other separated parts of several hundred animals of different descriptions and in different conditions as to age, maturity, and fatness, were taken; whilst in some carefully-selected cases the total amounts of fat, nitrogenous substance, mineral matter, and water, were determined †.

* "On the Composition of Foods in relation to Respiration and the Feeding of Animals," Report of the British Association for the Advancement of Science for 1852. "Agricultural Chemistry: Sheep-Feeding Manure," part 1, Journ. Roy. Ag. Soc. Eng. vol. x. part 1, 1849. "Reports of Experiments on the Comparative Fattening Qualities of different Breeds of Sheep," *ibid.* vol. xii. part 2, 1851; vol. xiii. part 1, 1852; vol. xiv. part 1, 1855. "Agricultural Chemistry: Pig Feeding," *ibid.* vol. xiv. part 2, 1853. "On the equivalency of Starch and Sugar in Food," Rep. Brit. Assoc. for 1854.

† "Experimental Inquiry into the Composition of some of the Animals Fed and Slaughtered as Human Food," Phil. Trans. part 2, 1859; also Proceedings of the Royal Society, vol. ix. p. 348. "On the Composition of Oxen, Sheep, and Pigs. and of their Increase whilst Fattening," Journ. Roy. Ag. Soc. Eng. vol. xxi. part 2, 1860. "On the Chemistry of the Feeding of Animals for the production of Meat and Manure," Proc. Roy. Dub. Soc. March 31, 1864.

that, in the case of fattening animals, the amount *ed in relation to a given body-weight within a given* regulated, not only by the demands of the system of respiration, perspiration, &c., and for the repair of nitrogenous substance, but also by the addi-
 ents for growth and increase; whilst, on the other
 unt required to be consumed *for the production of*
t of increase will, in its turn, include that due to
 f the system for respirable and perspirable matter
 of the waste of nitrogenous substance. Whether,
 xperimental results were calculated so as to show
 nsumed per 100 lbs. live-weight per week, or to
 s. increase in live-weight, it was strikingly brought
 parable experiments, that it was in neither case the
 rogenous constituents, but in both the amount of
 available non-nitrogenous (or total organic) sub-
 od that had regulated the results obtained.

ie reader to our former papers for all experimental
 the fuller discussion of the results and statement
 ons, we will close this part of the subject in words
 paper given at the Meeting of the British Asso-
 ast in 1852*. The sentence as quoted had refer-
 ults obtained with sheep; but subsequently those
 pigs were summarized in almost the same words:—
 : consider that it is the results obtained under the
 of animal life that we are seeking to measure and
 es, and if we also bear in mind the various sources
 to which our actual figures must be submitted in
 their true indications, we think that it cannot be
 beyond a limit below which few, if any, of our
 ng food-stuffs are found to go, it is their available
 s constituents, rather than their richness in the
 es, that measure both the amount consumed to a
 of animal, within a given time, and the increase in
 d."

mind the nature of the respiratory process, and the
 which its demands must necessarily exercise over
 food consumed, it will scarcely appear surprising
 on at least should be chiefly regulated by the sup-
 od of non-nitrogenous constituents; but that *the*
ease obtained in feeding animals for the butcher
 ear a closer relationship to the supply of the
 is than to that of the nitrogenous constituents,

omposition of Foods in relation to Respiration and the
 als," Report of the British Association for the Advance-
 for 1852.

might perhaps well be looked upon as inconsistent with the currently adopted views as to the highly nitrogenous character of the increase of animals fed for human food, and, indeed, of the highly nitrogenous character of the animal portion of human food generally.

The investigation into the composition of the fattening animals, and their increase, above alluded to, showed, however, how small was the proportion of the nitrogenous substance of the food that was stored up in the increase of the animal, and also that the proportion of fat in the increase was much greater than had previously been supposed. The results further led to the remarkable conclusion, that, reckoning the fat of the estimated total consumed portions of animals admitted to be in only a proper condition of fatness, into its starch-equivalent, there was, on the average, a higher proportion of so-reckoned non-nitrogenous substance to one of nitrogenous substance in such animal food than in bread itself. It was concluded, indeed, that, on the large scale, the introduction of animal aliments into our otherwise chiefly farinaceous diet did not increase, but diminish the *relation* of the nitrogenous or so-called "flesh-forming," to the non-nitrogenous constituents (reckoned according to their respiratory and fat-forming capacity) in the collective food. The important bearing of these facts in forming an estimate of the characteristics of different human dietaries will be at once apparent.

So much, then, for the characteristic food requirements of animals exposed to as little exertion as possible, and fed with the express view of accumulating flesh and fat in their bodies. Concurrently with the earlier experiments to determine the relations of food and body-weight and increase above referred to, the question of the relation of the amount of the constituents voided (especially the nitrogen) in the liquid and solid excrements to that in the food consumed, was also investigated. Consistently with the results obtained in regard to the amount and character of the increase resulting from the consumption of very different amounts of nitrogenous substance, it was found that the amount of nitrogen voided by fattening animals fed under equal conditions as to the exercise of force, bore a very direct relation to that supplied in the food. So direct, indeed, is the connexion between the composition of the matters excreted and that of the food consumed, that we have constructed Tables showing the relative value of the manure produced by fattening animals from a given weight of different food-stuffs according to the composition of the latter.

But more to our present purpose—so striking were the results obtained in regard to the connexion between the compos-

relations to various exigencies of the Animal Body.

tion of the food on the one hand, and the amount consumed, the amount and character of the increase produced, and the composition of the excreted matters, on the other, and, on some important points, so contrary in their indications to the prevailing views, that we were led at once to turn our attention to human dietaries, and especially to a consideration of the management of the animal body undergoing somewhat excessive labour, as, for instance, the hunting horse, the racer, the cab-horse, and the foxhound, and also pugilists and runners. The conclusions to which we were led by this study were briefly summarized in 1852 as follows* :—

“.....that in the cases, at least of ordinary exercise of force, the exigencies of the respiratory system keep pace more nearly with the demand for nitrogenous constituents of food than is usually supposed ;”

And further :—

“A somewhat concentrated supply of nitrogen does, however, in some cases, seem to be required when the system is over-taxed; as for instance, when day by day, more labour is demanded of the animal body than it is competent without deterioration to keep up; and perhaps also, in the human body, when under excitement or excessive mental exercise. It must be remembered, however, that it is in butcher's meat, to which is attributed such high flesh-forming capacity, that we have also, in the fat which it contains, a large proportion of respiratory material of the most concentrated kind. It is found, too, that of the dry substance of the *egg*, 40 per cent. is pure fat.

“A consideration of the habits of those of the labouring classes who are under- rather than over-fed, will show, that they first have recourse to fat meat, such as pork, rather than to those which are leaner and more nitrogenous; thus perhaps indicating, that the first instinctive call is for an increase of the respiratory constituents of food. It cannot be doubted, however, that the higher classes do consume a larger proportion of the leaner meats; though it is probable, as we have said, that even with these as well as pork, more *fat*, possessing a higher respiratory capacity than any other constituent of food, is taken into the system than is generally imagined. Fat and butter, indeed, may be said to have about twice and a half the respiratory capacity of starch, sugar, &c. It should be remembered, too, that the classes which consume most of the leaner meats, are also those which consume the most butter, sugar, and in many cases, alcoholic drinks also.

“It is further worthy of remark, that wherever labour is expended in the manufacture of staple articles of food, it has
Report of the British Association for the Advancement of Science for 1852.

generally for its object the concentration of the non-nitrogenous, or more peculiarly respiratory constituents. Butter, and alcoholic drinks are notable instances of this. (Which at first sight might appear an exception, is in reality not so; for those cheeses which bring the highest price are those which contain the most butter; whilst butter is always dearer than cheese.)

"In conclusion, it must by no means be understood that we would in any way depreciate the value of even a somewhat liberal amount of nitrogen in food. We believe, however, that on the current views too high a relative importance is attached to it; and that it would conduce to further progress in this most important field of enquiry if the prevailing opinion on the subject were somewhat modified." *

It will be borne in mind that, at the time the statement here quoted was made, the opinions expressed were directly contrary to all recognized authority on the subject, and since that date that so much evidence has been accumulated in regard to the amounts of urea, and the amounts of carbonic acid and other products, given off under varied conditions of food and exercise. Still, from the facts even then at command, it was concluded that the increased demand for food resulting from the exercise of muscular power was specially characterized by the requirement for an enhanced amount of the non-nitrogenous constituents. Confirmatory evidence was, however, long wanting.

In 1854 we selected two pigs as nearly as possible of equal weight and character; to one was given, *ad libitum*, lentil-meal (containing about 4 per cent. of nitrogen), and to the other, *ad libitum*, barley-meal (containing less than 2 per cent.). After the animals had been kept for a certain time on their respective foods, one comparative experiment was conducted for a period of three days, and another for a period of ten days. At the end of each experiment, and, besides other particular details, the amounts of nitrogen consumed in food, and voided as urea, were determined.† The result was, that with exactly equal conditions as to exercise, both animals being in fact at rest, the amount of urea passed by the one feeding on the highly nitrogenous lentil-meal was more than twice as great as that

* It is worthy of remark, too, that neither are the most highly nitrogenous wheats the most valued by the baker for the purposes of bread-making, nor is the most highly nitrogenous bread the most valued by the chiefly sedentary working man. See "On some Points in the Composition of Grain, its Products in the Mill and Bread," Journ. Chem. Soc. vol. 1, p. 100.

† Phil. Trans. part 2, 1859, p. 554.

by the one fed on the barley-meal. We have since made other such experiments with similar results.

It was clear, therefore, that the rule laid down by Liebig, and assumed to be substantially correct by so many writers, did not hold good—namely, that “The sum of the mechanical effects produced in two individuals, in the same temperature, is proportional to the amount of nitrogen in their urine; whether the mechanical force has been employed in voluntary or involuntary motions, whether it has been consumed by the limbs or by the heart and other viscera”—unless, indeed, as has been assumed by some experimenters, there is, with increased nitrogen in the food, an increased amount of mechanical force employed in the “involuntary motions” sufficient to account for the increased amount of urea voided.

It was at any rate obvious that, if the amount of urea voided by one animal at rest could be from two to three times as great as that voided by a similar animal also at rest, and under otherwise equal conditions, provided only that the food of the one contained from two to three times as much nitrogen as that of the other, the amount of urea passed could not be any measure of the amount of muscular power exerted; and this evidence, considered in connection with that relating to the demands of the system not only of the fattening animal but of the animal body fed with a view to mechanical exertion, afforded further confirmation of the view we had already put forward as above quoted, and also led to the extension and more definite expression of it.

The results of Bischoff and Voit, conducted through a period of many months, with a dog, either submitted to hunger or fed from time to time on foods containing very different amounts of nitrogenous substance, showed a very variable amount of urea voided, although the animal were kept under equal conditions as to exercise. Still, on the publication of those results in 1860, the authors assumed that although there had been no greater exercise of force manifested in the form of external work, yet when the amount of nitrogenous substance in the food was greater, and the amount of urea voided correspondingly greater, there must have been a corresponding increase in the force exercised in the conduct of the actions proceeding within the body itself in connection with the disposition of the increased amount of nitrogenous substance consumed. When, however, they subsequently found that the amount of urea passed by the animal when subjected to somewhat severe labour was, other things being equal, no greater than when at rest, whilst the carbonic acid evolved was much increased by such exercise, their view was of necessity somewhat modified.

Again, the results of Dr. Edward Smith, which showed great

10 *On Food in its relations to various exigencies of the Animal Body.*

variation in the amount of urea passed when there was concurrent variation in the amount of nitrogenous substance in the food, and comparatively little variation in the amount of urea voided with great variation in the amount of labour performed, but, on the other hand, great increase in the carbonic acid evolved with increased exercise of force, obviously still further pointed to the correctness of the view that with muscular exertion there was a more marked increased demand for the non-nitrogenous than for the nitrogenous constituents of food.

That this was the necessary conclusion from the results of our own investigations, and also from those of the researches of Bidder and Schmidt, Bischoff, Voit, Pettenkofer, E. Smith, and others, we have frequently maintained. Indeed the view urged in public discussion has been, that all the evidence at command tended to show that by an increased exercise of muscular power there was, with increased requirement for respirable material, probably no increased production and voidance of urea, unless, owing to excess of nitrogenous matter in the food, or a deficiency of available non-nitrogenous substance, or diseased action, the nitrogenous constituent of the fluids or solids of the body were drawn upon in an abnormal degree for the supply of respirable material.

From the facts briefly summarized in the foregoing pages, it will be obvious that the generally accepted views in regard to the adaptation of food, according to its composition, to the various exigencies of the animal body, require modification in other respects than in so far as they relate to the source or developement of muscular power alone. At the same time we hail with much satisfaction the confirmation of the views we have so long maintained on the point in opposition to general authority, which has recently been afforded by the results of the interesting, though limited experiment of Professors Fick and Wislicenus, so ably discussed by them in their paper, and by Professor Frankland in his lecture.

(Read August 24, Section D, and August 27, Section B.)

ON THE SOURCES OF THE FAT OF THE ANIMAL BODY.

BY

LAWES, F.R.S., F.C.S., and J. H. GILBERT, Ph.D., F.R.S., F.C.S.*

Baron Liebig had concluded that the fat of Herbivora must be derived part from the carbo-hydrates of their food, but that it might also be from nitrogenous compounds. MM. Dumas and Boussingault at first called in this view; but subsequently the experiments of Dumas and Milne with bees, of Persoz with geese, of Boussingault with pigs and ducks, and authors with pigs, had been held to be quite confirmatory of Liebig's view; and so far as the formation of fat from the carbo-hydrates was concerned, however, at the Bath Meeting of the British Association, Dr. Hayden, of read a paper before the Physiological Section, in which, basing his conclusion on certain physiological considerations of a purely qualitative kind, he expressed doubt on the point. In August 1865, again, at a meeting of the Congress of Natural Chemists, held at Munich, Professor Voit, from the results of experiments with dogs fed on flesh, maintained that fat must have been produced from nitrogenous constituents of the food, and that these were probably the chief, if not the only source, of the fat even of Herbivora. In the course of the discussion allowed, Baron Liebig disputed this conclusion; and his son, Hermann von Liebig, has since written a paper on the subject, in which, illustrating his views by experiments with cows, he admits the probability that fat may be derived from nitrogenous substance, but nevertheless concludes that this is neither the chief nor even the chief source of fat, in the ordinary feeding of Herbivora.

The authors agreed with the conclusions of these latter authorities. The data afforded by Hermann von Liebig did not, however, afford conclusive evidence on the point, and they considered that the results of experiments with cows were, in many respects, less appropriate for the purposes of the inquiry than those with some other animals. They showed, illustrating the various points by reference to experiments of their own, that, compared with either cows, oxen, or sheep, the pig had a less proportion of alimentary organs and contents, consumed food of a much less character, produced a much larger amount of fat both in relation to a given weight of animal within a given time and to the amount of food consumed, voided a less proportion of the solid matter of its food in its solid and liquid excreta, and, finally, its increase contained a larger proportion of fat. For these results obtained with pigs must be much more conclusive as to the sources in which the fat which they yield than those with either cows, oxen, or sheep. The authors were exhibited showing the results which had been obtained by the authors' various experiments with pigs; and from these the following main conclusions were drawn:—

1. That certainly a large proportion of the fat of the Herbivora fattened for food must be derived from other substances than fatty matter in the food.
2. That when fattening animals are fed upon their most appropriate food, much of the stored-up fat must be produced from the carbo-hydrates it supplies.
3. That nitrogenous substance may also serve as a source of fat, more especially when it is in excess, and the supply of available non-nitrogenous constituents is defective.

* For fuller report, see the Philosophical Magazine for December 1866.



ON THE SOURCES OF THE FAT OF THE ANIMAL BODY.

IN 1842, Baron Liebig* maintained that the fat of Herbivora must be derived in great part from the carbo-hydrates of their food, but considered that it might also be produced from nitrogenous compounds. MM. Dumas and Boussingault † at first called in question this view: but subsequently the experiments of Dumas and Milne-Edwards ‡ with bees, of Persoz § with geese, of Boussingault || with pigs and ducks, and of ourselves with pigs ¶, were held to be quite confirmatory of Liebig's view, at any rate so far as the formation of fat in the animal body from carbo-hydrates was concerned.

In 1864, however, at the Bath Meeting of the British Association for the advancement of Science, Dr. Hayden, of Dublin, read a paper before the Physiological Section, in which, basing his conclusions upon certain physiological considerations of a purely qualitative kind, he argued that fat was not producible in the body from sugar and allied substances, but that both eventually served for the production of carbonic acid and water; and sugar being the most readily oxidized, so saved the combustion, and favoured the storing of fat.

Again, in August 1865, at a Meeting of the Congress of Agricultural Chemists, held in Munich (at which one of the authors was present), Professor Voit**, from the results of experiments with dogs fed on flesh, maintained that fat must have been produced from the nitrogenous constituents of the food, and that these were probably the chief if not the only source of the fat, even of Herbivora. In favour of the probability of this view, Professor Voit refers to the formation of adipocere from nitrogenous substance; but he mainly relies upon the fact that, in experiments by Pettenkofer and himself, in which large quantities of flesh were given to a dog, the whole of the nitrogen reappeared in the form of urea and in the fæces, whilst only a portion of the carbon was recovered in the urine, fæces, and the products of respiration and perspiration, from which it was concluded that some had been retained in the body, and had con-

* Organic Chemistry of Physiology and Pathology, p. 81 *et seq.*

† Balance of Organic Nature, 1844, p. 116 *et seq.*

‡ Comptes Rendus de l'Académie des Sciences, vol. xvii. p. 531.

§ Ann. de Chim. et de Phys. vol. xiv. p. 408 *et seq.*

|| Ann. de Chim. et de Phys. vol. xiv. p. 419 *et seq.*; xviii. p. 444 *et seq.*

¶ "On the Composition of Foods in relation to Respiration and the Feeding of Animals," Report of the British Association for the Advancement of Science for 1862.

** Versuchs-Stationen Organ. vol. viii. No. 1, 1866, p. 23.

tributed to the formation of fat. That animals nevertheless not become fat when fed upon very highly nitrogenous food is sufficiently explained by the greater number of blood-corpuscles, the result of such diet, and the greatly increased activity of oxidation of nitrogenous substance under such conditions; whilst, on the other hand, the accumulation of fat when fat and carbo-hydrates are supplemented to a liberal nitrogenous diet, he considers to be connected with the much less activity of oxidation of the nitrogenous substance and fatty matter then takes place, rather than attributable to the direct production of fat from the carbo-hydrates.

In the discussion which followed the reading of P. Voit's paper, Baron Liebig forcibly called in question P. Voit's conclusions, maintaining not only that it was impossible to form conclusions on such a point in regard to Herbivora from the results of experiments made with Carnivora, but that direct quantitative results obtained with herbivorous animals had afforded apparently conclusive evidence in favour of the opposite view.

Since the Munich Meeting, Hermann von Liebig, Baron Liebig, has written a paper on the subject*, in admitting the probability that fat may be formed from nitrogenous substance, he nevertheless concludes that this is not its only, nor even its chief source, in the ordinary feed of Herbivora.

After referring to the leanness of the South Russian shepherds who consume very large quantities of dried meat, and the rotundity of the peasantry, especially the women, in Prussia where bread and fruits constitute the chief articles of food, von Liebig proceeds to illustrate the formation of fat from the nitrogenous constituents of food by our domestic Herbivora. In the calculation of the results of numerous experiments with cows in 1857, by Knop, Arendt, and Behr, in which details as to food, live-weight, and quantity and composition of milk, were accurately recorded. According to the method of calculation adopted, it appeared that, after deducting from the amount of nitrogenous substance taken in the food, that essential to be required by the system for other purposes, there was generally little or none remaining for the production of fat. In his calculations, however, H. von Liebig, besides taking account the probable amount of nitrogenous substance stored in the body in increase with gain of weight, or set at liberty when there is a loss of weight, as the case might be, deducted from the total amount of nitrogenous substance given in the food, not only that required for the production of the caseine of the milk, but

* *Versuchs-Stationen Organ.* vol. viii. No. 8, 1866.

the whole of that estimated to be required for the mere sustenance of the animal (according to its weight), independently of gain or loss, or milk produced.

It is obvious, however, as pointed out by Voit, and as afterwards admitted by H. von Liebig, that if nitrogenous substance may break up into urea and fat (with other products), the amount estimated to be required for the mere sustenance of the body should not be considered inadmissible for the formation of fat as one of its products, and therefore should not be deducted (with that appropriated for the production of increase and of the caseine of the milk) from the amount supplied in the food in estimating whether or not it provided sufficient for the formation of the fat known or calculated to be produced.

H. von Liebig states that he selected experiments with cows as the basis of his illustrations, considering that, when in a normal state, the change in the solid substance of the body of the animal was comparatively small, if not indeed immaterial, and that the fixed products of the food, beyond what might be required for the mere maintenance of the body, were accumulated and easily estimated in the milk collected; whilst he considered, on the other hand, that the point in question could not be settled by reference to results relating to fattening animals, without the aid of an apparatus for the determination of the products of respiration and perspiration. We believe, however, that with a proper selection of fattening animals it may be satisfactorily illustrated without the aid of any such apparatus; and it is the object of this paper briefly to discuss the question of the sources of the fat of the animal body by reference to the results of experiments with such animals.

As already intimated, the objections of Dr. Hayden to the supposition that fat is formed from the carbo-hydrates of the food, were based upon physiological considerations of a qualitative, but not at all of a quantitative kind. Voit's argument was, on the other hand, founded upon strictly quantitative results, obtained, however, under conditions as to choice of animal and of food, in which the formation of fat, if it took place at all, must of necessity be attributed to the nitrogenous constituents consumed. H. von Liebig also relied upon quantitative results as the basis of his illustrations; but those selected, when properly considered, afforded, to say the most, only negative evidence on the point.

The question arises—What description of animal is likely to yield the most direct and conclusive evidence as to the source of the fat stored up in its body? Obviously the one which is fed more especially with a view to the production of fat, which consumes in its most appropriate fattening food a relatively large proportion of carbo-hydrates, and which yields a

large proportion of fat, both in relation to the weight of animal within a given time, and to the amount of food consumed. The following Table (I.), which summarizes the results of a great many direct experiments of our own*, will show that of the ox, the sheep, and the pig—the most important of the animals fed and slaughtered as human food—the last pre-eminently supplies the required conditions.

TABLE I.—Comparative fattening-qualities of different animals.

	Oxen.	Sheep.	Pigs.
Relation of parts in 100 live-weight.			
Average of.....	16	249	*59
Stomach and contents	11·6	7·5	13
Intestines and contents	2·7	3·6	6·2
Heart, aorta, lungs, windpipe, liver, gall-bladder } and contents, pancreas, spleen, and blood	14·3 7·0	11·1 7·3	7·5 6·6
Per 100 live-weight.			
Dry substance consumed in food per week.....	12·5	16·0	27·0
Increase yielded per week	1·13	1·76	6·43
Per 100 dry substance of food.			
Total dry substance in increase	6·2	8·0	17·6
Fat in increase	5·2	7·0	15·7
Total dry substance in urine and faeces	36·5	31·9	16·7
Average fat per cent.			
In lean condition	16·0	18·0	22·0
In fat condition.....	30·0	33·0	44·0
In increase whilst fattening	60·0	65·0	70·0

Looking first to the comparative structure of the animals, as far as it may be considered characteristic or indicative of the description of the food, it is seen that, of stomach and contents, the ruminant ox has a much larger proportion than the ruminant sheep, and the ruminant sheep in its turn much more than the non-ruminating pig. Consistently with these facts we find that the ox consumes in its food a much larger proportion of

* For the data upon which most of the average results given in the Table are founded, see "Experimental Inquiry into the Composition of some of the Animals fed and slaughtered as Human Food," Phil. Trans. Part II. 1859. In the estimates given "per 100 live-weight" and "per 100 dry substance of food," it is assumed that the oxen and sheep are liberally fed on oil-cake, clover-chaff, and roots, and the pigs on barley-meal alone; with different foods the results will, of course, be different.

only slowly digestible, or indigestible, cellulose than the sheep, and the sheep again very much more than the pig. The usual food of oxen and sheep, consisting as it does in large proportion of unripened or imperfectly ripened vegetable matter, is, in fact, essentially crude, containing not only a considerable amount of defectively elaborated and probably unassimilable nitrogenous substance, but also a large proportion of comparatively indigestible non-nitrogenous matter. Accordingly, complexity and great capacity of stomach, and slow progress of the food through the organ, are characteristics of the structure and digestive process of the animals.

Of intestines and contents, on the other hand, the ox has a less proportion than the sheep, and the sheep considerably less than the pig.

In fact, the relatively very small proportion of stomach and contents, and relatively very large proportion of intestines and contents in the pig are very striking. But when we consider that his most appropriate fattening food consists of ripened seeds and highly starchy roots, containing little indigestible woody fibre, and their non-nitrogenous constituents almost wholly in the form of starch, the primary change of which is known to take place almost throughout the length of the intestinal canal, the reason of the relatively small proportion of stomach, and large proportion of intestines, seems to be at once apparent.

Passing from a consideration of the receptacles and, so to speak, first laboratories of the food, we will only remark, in reference to the remaining results given in the upper portion of the Table, that, of what may be called the further elaborating organs of the body, and their fluids—the heart, liver, lungs, blood, &c.—the proportion, taken in the aggregate, is strikingly similar in the three descriptions of animal.

The second division of the Table shows that, notwithstanding its much larger proportion of stomach and contents, the ox consumes, for a given live-weight within a given time, only about three-fourths as much dry substance of the food as the sheep, and less than half as much as the pig with its very small proportion of stomach and contents. The ox gives, too, in proportion to a given live-weight within a given time, much less increase than the sheep, and only from one-fifth to one-sixth as much as the pig.

Reckoned in proportion to a given amount of dry substance of food consumed, the ox gives less, both of total dry substance in increase and of fat in increase, than the sheep, and only about one-third as much of either as the pig, whilst the ox voids of dry substance in fæces and urine the largest proportion, the sheep somewhat less, and the pig little more than half as much as the sheep, and less than half as much as the ox.

Lastly, the proportion of fat, whether reckoned in relation to the total weight of the body, or to the weight of the increase whilst fattening, is greater in the sheep than in the ox, and greater still in the pig.

Whilst referring to the connexion between the weight and capacity of the stomach and the character of the food, it will not be without interest to call attention to the gradation in the proportion from the ox to the sheep, from the sheep to the pig, and from the pig to man. Below is given the approximate average proportion of stomach, by weight, in 100 live-weight of each.

Oxen.	Sheep.	Pigs.	Man.
3.19	2.44	0.88	0.38

Without assuming that relative weight represents with numerical exactitude relative capacity or size, we nevertheless cannot doubt that these figures have a very obvious significance. Thus, the ox consumes the largest proportion of difficultly digestible or indigestible woody-fibre, the sheep less, the pig scarcely any, but a much larger proportion of comparatively easily digestible starch, whilst man, within certain limits, the better he is fed the less does the non-nitrogenous portion of his food consist of starch, and the more of the much more highly concentrated alimentary substance fat, produced for him from much less concentrated vegetable food-materials by the animals which he feeds for his own consumption.

From the facts which have been briefly stated, it will be obvious that, of the most important animals which we feed for human food, the pig offers many advantages as a subject for the consideration of the source in the food of the fat which he yields. Thus, for a given live-weight he comprises a comparatively small proportion of alimentary organs and contents, and he consumes a large proportion of food, and yields a large proportion both of total increase and of fat, within a given time; his food is, as such, of a high character, yielding, compared with that of oxen or sheep, for a given weight of it much more total increase, much more fat, and much less excreted and necessarily effete matter; whilst his proportion of fat is the greatest, both in a given live-weight and in his increase whilst fattening. It results that changes in his live-weight are in a much less degree likely to be influenced by variation in the amount of the contents of the stomach and intestines, and are therefore much more direct indications of real increase of the substance of the body, and hence that there is much less probable range of error in calculating the amount and composition of the increase in live-weight in relation to the amount and composition of the food consumed.

In fact, from the very opposite characters of the ruminant in these respects, it is very much less appropriate for the purpose of estimating the sources in its food of the fat of its body. It is true, that there is the advantage with the cow, that that important product of the food—the milk—is collected externally to the body, and hence its amount and composition can be easily determined; but the changes of weight of the animal itself, though comparatively small, are due to a greater variety of circumstances, and can, therefore, with less of certainty be properly interpreted than even in the case of either the ox or the sheep. Indeed, when experiments are conducted with cows or oxen, or even with sheep, for periods of a few weeks only, the variation in live-weight may in very great proportion be due to variation in the contents of the alimentary organs merely.

The selection and calculation of results brought to view in Table II. (p. 8) will show that, when experiments are conducted with pigs fed on good fattening food, for periods of not less than eight or ten weeks, the amounts, both of total increase and of fat stored up, are so great in proportion both to the original weight of the animal and to the food consumed, that the data so obtained may be safely relied upon as a means of estimating, with sufficient accuracy for the purposes of the present discussion, from what constituent or constituents of the food the fat of the animals has been derived.

Experiment 1.—In this experiment two pigs of the same litter, of equal weight, and, as far as could be judged, of similar character, were selected. One was killed at once, and the amount of total dry or solid matter, nitrogenous substance, fat, and mineral matter, in its body, determined. The other was then fed for a period of 10 weeks on a good mixed food, containing, however, a more than usually high proportion of nitrogenous substance. It was then weighed and killed, and its composition was determined as in the case of the other animal. The results so obtained supplied an important portion of the data requisite for the calculation of the composition of the increase in the other cases *. The food consisted of a mixture of bean-meal, lentil-meal, and bran, each one part, and barley-meal three parts, given *ad libitum*.

* For further details relating to this and the other experiments, we must refer to our former papers, as follows:—"On the Composition of Foods in relation to Respiration and the Feeding of Animals" Report of the British Association for the Advancement of Science for 1852. "Agricultural Chemistry: Pig Feeding," Journ. Roy. Ag. Soc. Eng. vol. xiv. part 2, 1853. "On the Equivalency of Starch and Sugar in Food," Report of the British Association for 1864. "Experimental Inquiry into the Composition of some of the Animals Fed and Slaughtered as Human Food," Phil. Trans. part 2, 1859.

TABLE II.—Relation of the total Fat in the increase, to the ready-formed fatty matter in the Food, and of the Carbon in the Fat produced within the body, to that in the nitrogenous substance consumed, in experiments with Fattening Pigs.

Experiments.....	1.	2.	3.	4.	5.	6.	7.	8.	9.
Conditions, and actual results of experiment.									
Number of animals.....	1	3	3	3	3	3	3	3	3
Duration of Experiment (weeks).....	10	8	8	8	8	10	10	10	10
Non-nitrogenous substance to one nitrogenous substance in food.....	3.6	5.3	5.0	6.6	6.0	4.1	4.1	4.7	3.9
Final live-weight (lbs.).....	118	429	440	431	448	286	285	281	294
Final live-weight (lbs.).....	191	685	743	652	739	533	533	563	602
Increase in live-weight (lbs.).....	88	266	303	221	291	247	248	272	312
Increase on 100 original weight.....	88.4	68.7	68.9	51.2	64.9	56.4	57.0	56.8	106.8
Increase per 100 lbs. live-weight, per week.....	6.0	5.7	6.4	5.1	6.1	6.0	6.1	6.6	7.0
Calculated for 100 increase in live-weight.									
Fat.....	63.1	73.9	69.6	79.0	71.2	64.1	63.9	69.0	59.9
	15.6	20.4	11.2	26.3	12.4	7.9	7.9	7.3	6.6
	47.5	53.5	58.4	52.7	58.8	56.2	56.0	54.7	53.3
Nitrogenous-substance.	100.0	107.0	138.0	57.0	64.0	81.0	81.0	74.0	82.0
	7.8	6.1	6.7	5.3	6.5	7.5	7.6	8.0	7.8
	92.2	100.9	131.3	51.7	57.5	73.5	73.4	68.0	83.8
Carbon.....	26.8	41.2	48.0	40.6	45.3	43.3	43.1	42.1	41.0
	44.0	48.1	62.6	24.7	27.4	35.1	35.0	31.5	35.2
	+7.4	+6.9	+17.6	15.9	17.9	8.2	8.1	10.6	5.8
Calculated for 100 carbon in estimated "produced" fat.									

Experiments 2 & 3.—In both these experiments the proportion of nitrogenous substance in the food was very large; the relation of non-nitrogenous to one of nitrogenous substance being in Exp. 2 little more than half, and in Exp. 3 little more than one-third as much as is usual in the recognized good fattening food of the animal. In Exp. 2 the food consisted of bran, bean and lentil-meal, and Indian-meal, each given separately, and *ad libitum*; and in Exp. 3 of an equal mixture of bean and lentil-meal only, given *ad libitum*.

Experiments 4 & 5.—In Exp. 4 the food consisted of Indian meal only, and in Exp. 5 of barley-meal only, in each case given *ad libitum*. Barley-meal is undoubtedly the most approved staple fattening food of the pig; and the result was that, in both these experiments, the proportion of non-nitrogenous to nitrogenous substance in the food was very nearly, though rather higher than, the average in that which is recognized as the most appropriate fattening food of the animal.

Experiments 6, 7, 8, & 9.—The peculiarity of this series was, that the food contained less ready-formed fat than was the case in either of the other experiments, and that a large proportion of the non-nitrogenous substance supplied was in the form of either pure starch, pure sugar, or both. In Exps. 6, 7, & 8, a fixed quantity of lentil-meal and bran (averaging nearly $3\frac{1}{4}$ lbs. lentil-meal and about 9 ounces bran) was given per head per day, and, in addition, in Exp. 6 sugar, in Exp. 7 starch, and in Exp. 8 sugar and starch, each separately, *ad libitum*. In Exp. 9 lentil-meal, bran, sugar, and starch were each given separately, *ad libitum*.

The figures given in the Table show that the increase in weight was in no case less than 50, and in several nearly, and in one more than 100 per cent., upon the original weight of the animals; the amounts ranging from 51.3 to 68.9 per cent. when the experiment extended over eight, and from 85.4 to 106.8 per cent. when it extended over ten weeks.

The determined or estimated amount of fat stored up in the increase was also in all cases very large, amounting to 63 per cent. of the total increase in Exp. 1, in which it was experimentally determined, and calculated to be even more than this in several of the other cases. The tendency to error in the calculations would, however, be to give the proportion too low in Exps. 6, 7, 8 & 9, which were conducted over a period of ten weeks, and in which the proportion of increase upon the original weight was very high, and to give it too high in Exps. 3, 4 & 5, conducted only over eight weeks, but more especially in Exps. 2 & 4, in which the proportion of increase upon the original weight was comparatively small. The range of the probable error of calculation here indicated is, however, not such as in any degree to throw doubt upon the validity of

any conclusions which will be drawn from the indications of the figures as they stand.

It is seen that, of the determined or estimated total fat stored up in the increase, the proportion which could possibly have been derived from the ready-formed fat of the food, even supposing the whole of that supplied had been assimilated, was so small as to leave no doubt whatever that a very large proportion of the stored-up fat must have been produced from other constituents than the ready-formed fatty matter of the food. According to the figures given in the Table, the proportion of fat which must have been so produced, ranged from about two-thirds to about eight-ninths of the total amount stored up.

Assuming it, then, to be established beyond doubt, that there was a very large formation of fat within the body from other constituents than the fatty matter of the food, the question arises, whether this large amount of produced fat could possibly have been derived from the nitrogenous constituents of the food? or whether it must of necessity have had its source, in greater or less proportion, in the carbo-hydrates at the same time supplied? The results adduced afford conclusive evidence on this point also.

The figures show that, after deducting from the total amount of nitrogenous substance consumed for the production of 100 lbs. of increase in live-weight, the small amount estimated to be stored up in the increase, there remains a very large proportion available, it may be, for the production of fat with other products.

If we next compare the amount of carbon in the estimated produced fat, with the amount contained in the nitrogenous substance of the food not stored up as increase, minus that contained in the urea which would be one of the final products of the breaking up of this nitrogenous substance (or its equivalent given off), the result shows in some cases an excess, and in others a deficiency, of carbon possibly available from the nitrogenous constituents of the food, compared with that required for the formation of the fat estimated to be derived from other constituents than the ready-formed fat in the food.

Reckoned to the standard of 100 carbon in the estimated produced fat, it is seen, as shown in the two bottom lines of the Table, that in Exps. 1, 2 & 3, in which the proportion of non-nitrogenous to nitrogenous substance in the food was (especially in Exp. 3) considerably less than in such food as experience has shown to be the most appropriate in the fattening of the pig—that is to say, in which the nitrogenous substance was in considerable excess over the amount and proportion usually supplied—there was, according to the calculation, more than sufficient carbon possibly available from the nitrogenous substance of the food for the formation of the fat estimated to be produced.

In Exps. 4 & 5, however, in which the relation of the non-nitrogenous to the nitrogenous substance in the food was much more nearly that in the usual food of the well-fed fattening pig, it is reckoned that there was about 40 per cent. of the carbon of the produced fat which could not possibly have been supplied from the nitrogenous constituents of the food.

In the other experiments (Nos. 6, 7, 8 & 9), in which again the proportion of the non-nitrogenous to the nitrogenous constituents of the food was lower than usual (though not so much so as in Exps. 1, 2 & 3)—in which, in fact, the nitrogenous constituents were in excess—there was still a considerable proportion of the carbon of the produced fat which the nitrogenous constituents of the food could not possibly have supplied.

It is hardly necessary to point out that, according to the mode of illustration we have adopted, the figures show, not only the utmost proportion of the carbon of the stored-up fat which could possibly have had its source in the nitrogenous substance of the food, but even notably more than could possibly have been so derived. Thus, to say nothing of other considerations, it has been assumed for simplicity of illustration, and granted for the sake of argument, that the whole of the ready-formed fatty matter of the food contributed to the fat stored up, that the whole of the nitrogenous substance of the food not stored up in increase would be perfectly digested and become available for the purposes of the system, and that in the breaking up of the nitrogenous substance for the formation of fat no other carbon-compounds than fat and urea would be produced. It is obvious, however, that these assumptions are in part improbable, and in part quite inadmissible, and that the tendency of each of them is to show too large a proportion of the produced fat to have been possibly derived from the nitrogenous constituents of the food.

The amount of fat necessarily derived from other sources than the nitrogenous constituents of the food must therefore be greater than our mode of estimate can indicate; and it is obvious, from the figures given in the Table, that the less the excess of nitrogenous substance in the food, the greater was the proportion of produced fat which must necessarily have had its source in the carbo-hydrates of the food, and that, at any rate in those cases in which the proportion of non-nitrogenous to nitrogenous constituents supplied was the more nearly that occurring in the admittedly most appropriate fattening food of the animal, the proportion of the fat which must necessarily have been derived from the carbo-hydrates was very large, even allowing all that was possible to have been produced from the nitrogenous substance of the food.

That, nevertheless, fat may be produced in the animal body

at the expense of nitrogenous substance, in greater or less degree according to the character of the animal and of the food, not only chemical and physiological considerations, but direct experimental evidence would lead us to conclude. Indeed we have, in former papers already referred to, called attention to the fact that the results of our experiments with fattening animals, when carefully considered, afford evidence in favour of such a conclusion. To discuss the point satisfactorily on the present occasion, by the aid of figures, would, however, unduly extend the limits of our paper.

But, as indicating the bearing of the results referred to, it may be stated, in passing, that in numerous cases, otherwise comparable, but in which the amount and proportion of the nitrogenous constituents consumed varied very greatly, the results clearly showed that, neither the amount of food consumed, nor the amount of increase in live-weight produced, bore any direct relation to the amount of nitrogenous substance supplied. On the other hand, both the amount of food consumed, and the amount of increase produced, bore a very close relation to the supply of digestible non-nitrogenous constituents, and even a closer relation still to the amount of total digestible dry organic substance (that is, nitrogenous and non-nitrogenous taken together); whilst, so far as could be judged from careful observation, the proportion of nitrogenous to non-nitrogenous substance (fat) in the increase did not vary in anything like a corresponding degree with the variation in the proportion of the nitrogenous and non-nitrogenous constituents in the food. The animals consuming excessive amounts of nitrogenous substance did, indeed, show a greater tendency to increase in frame and flesh; but they nevertheless became fat. It would appear, that the excess of nitrogenous substance had acted vicariously in defect of a greater supply of the non-nitrogenous constituents, contributing material not only to meet the respiratory exigencies of the animal, but also for the production of fat.

The main conclusions in regard to the sources of the fat of the animal body to which the evidence adduced has led, may be briefly stated as follows :—

1. That certainly a large proportion of the fat of the *Herbivora* fattened for human food must be derived from other substances than fatty matter in the food.
2. That when fattening animals are fed upon their most appropriate food, much of their stored-up fat must be produced from the carbo-hydrates it supplies.
3. That nitrogenous substance may also serve as a source of fat, more especially when it is in excess and the supply of available non-nitrogenous constituents is relatively defective.

SEWAGE UTILISATION.

BY

DR. J. H. GILBERT, F.R.S., &c.

By permission of the Council.

Excerpt Minutes of Proceedings of The Institution of Civil Engineers,
Vol. xlv. Session 1875-76.—Part iii.

LONDON: PRINTED BY WILLIAM CLOWES AND SONS,
STAMFORD STREET AND CHARING CROSS.

1876.

[*The right of Publication and of Translation is reserved.*]

SEWAGE UTILISATION.¹

Dr. J. H. GILBERT observed, through the Secretary, that there could be no doubt whatever that the interception or dry processes referred to by Mr. Redgrave were an immense improvement upon the old midden-pit or common privy systems; but no such plan could be accepted as a solution of the sewage difficulty. About four-fifths, or more, of the manurial value of human excretal matters were due to the urine. It was a desideratum with all such systems to exclude as much of the urine as possible, and the complete separation of the liquid from the solid dejections had been recommended. Of course, this would much reduce the value of any dry manure so produced, which was already so low as not to be worth more than its carriage beyond the immediate locality. Without such special separation, at the outside about one-third of the urine would be collected with the fæces. Under any such dry system there was, therefore, from two-thirds to the whole of the urine, besides all wash and other house drainage, still to be dealt with; and if the liquid had to go into a stream which served as the water-supply for other populations, sooner or later purification would be enforced. Passing the liquid through land was not only the best mode of purification, but promised the greatest return for the constituents it contained—whether profit to the towns, depended on many local circumstances. Then as to precipitation methods. There could be no doubt that any one of those referred to by Mr. Shelford would be a vast improvement upon doing nothing whatever with sewage that had to be turned into an open stream. No such plan was, however, likely to collect more, and would generally

¹ These remarks were communicated to the Secretary of the Institution of Civil Engineers in reference to the Papers by Mr. Redgrave "On Sewage Interception Systems, or Dry Sewage Processes," and by Mr. Shelford "On the Treatment of Sewage by Precipitation," read March 28, 1876.

collect less, than one-fourth of the nitrogen of the sewage in the solid manure. This one-fourth was, moreover, the least active and least valuable part. These plans also, as a rule, carried down the phosphates, but in a precipitated, not in a soluble, form; and in more than one scheme soluble phosphate had been used, and was converted from this more valuable into the less valuable precipitated condition. The estimate of the value of the nitrogen (reckoned as ammonia) in such a manure, containing only from 1 to 2 per cent. of it, at the same rate as when provided in guano containing about 12 per cent., and in a much more soluble condition, was entirely fallacious; as also was the valuation of precipitated phosphate at 3s. 3d. per unit. Would, then, such precipitated manures pay for their manufacture as such? He thought not. If the process were adopted mainly as a means of purification, what was the result? About three-fourths of the nitrogen of the sewage would remain in the liquid. This would exist, in the main, not as nitrates, but as ammonia and soluble organic compounds. This liquid, with all other house drainage, remained to be dealt with. He did not think that such a liquid would eventually be allowed to run into a water-supply stream. Passing it through the land would best purify it, and would yield the largest return. If the sewage were employed for irrigation, the less taken out of it, beyond the sludge, before use, the better; and if the phosphates were removed, they should be returned either to the sewage or to the land irrigated. In fact, where irrigation was to be eventually adopted, the less effective the precipitation process the better; indeed, the exclusion of the natural sludge was all that was desirable. He was by no means unconscious of the many difficulties involved in the general adoption of sewage irrigation, but he believed if rivers were to be kept from pollution, it would eventually have to be adopted, wherever practicable, before the liquid was discharged into them.


LONDON :
PRINTED BY WILLIAM CLOWES AND SONS,
STAMFORD STREET AND CHANCERY CROSS.

ON SOME POINTS IN CONNECTION WITH ANIMAL NUTRITION.

BEING AN ADDRESS DELIVERED AT SOUTH KENSINGTON,
IN THE BIOLOGICAL SECTION OF THE
SCIENCE CONFERENCES, MAY 26, 1876.

BY DR. J. H. GILBERT, F.R.S., F.L.S., F.C.S.

LONDON :
PRINTED BY VINCENT BROOKS, DAY AND SON,
GATE STREET, LINCOLN'S INN FIELDS, W.C.



ON SOME POINTS IN CONNECTION WITH ANIMAL NUTRITION.

BY DR. J. H. GILBERT, F.R.S., F.L.S., F.C.S.

A few days ago Professor M. Foster wrote to me to say that he intended to bring the subject of nutrition forward on this occasion, and asked me if I would take part in the discussion afterwards ; and as he and I had had a good deal of correspondence and conversation some little time since about the important question of the sources of the fat of the animal body, I concluded it was probably to that subject he wished me to devote my attention. At any rate, I looked up hurriedly the materials which Mr. Lawes and myself have collected in relation to that subject, and some allied points, and propose, with your permission, to lay the facts before you shortly, although Professor Foster has not given you his paper.

Thirty-five years ago, or more, I believe the view generally accepted was, that the carnivora found the fat which existed in their bodies ready-formed in the herbivorous animals they consumed, and that the herbivora in their turn found all the fat of their bodies ready stored up in the plants they consumed. About that time Liebig, in reviewing the composition of vegetable food, came to the conclusion that this was simply impossible, taking into consideration the amount of fat which was stored up by many animals in proportion to the known quantities in the food. He put forward the view that the carbohydrates of the food—starch, sugar, and so on—were important sources of the fat of the herbivora. For a short time this view was opposed, but only for a short time, by Dumas and Boussingault, and some other experimenters in France, though they afterwards accepted it.

The investigations of Mr. Lawes and myself, it must be borne in mind, have always had an agricultural object, so that if they were not conducted exactly in the way which the physiologist will say they might have been, it has been because we had not the same object before us, that is a purely physiological one. Very soon our own experiments led us to believe that Liebig was right in his conclusion on this point, but that he must be wrong on some other points in relation to the feeding of animals which he so ably discussed. We found it was pretty certain, from the consideration of the feeding experiments, that some of the fat must have the source which he assumed.

On the other hand, he assumed that the value of food to the animal was measured by the amount of nitrogen which it contained; that is to say, he maintained that, in the formation of meat, in the formation of milk, and in the exercise of force, the measure of the value of the food required, for these purposes, was the amount of nitrogen it contained; and in the case of the exercise of force, the amount of urea which was eliminated. We found, however, that we could give twice or three times the quantity of nitrogen within a given time to one animal as to another, both at rest, and that the amount of nitrogen eliminated in urea was almost proportional to the amount of nitrogen in the food, and had no direct connection with the amount of force exercised.

The question of which of the constituents in the food, were of the most importance for the exercise of force, and for the making of fat, remained in this condition until the experiments instituted in Munich, about 16 or 17 years ago, with Pettenkofer's beautifully contrived respiration apparatus, a model and drawings of modifications of which are in the next room. I am glad that after very much trouble on my part to get such an apparatus brought to this Exhibition, and entirely failing, it has after all been sent by some one. It consists of a chamber in which an animal can be put, and by a water wheel, or by some other power, the air is gently aspirated through the apparatus, then it passes through gauges, and through solutions, which absorb the carbonic acid, &c., and so the amount of air passing is gauged, and the products of respiration are determined. It is not the apparatus itself, but the results which it has brought out, which I wish to refer to on this occasion. In 1860, Bischoff and Voit published their first results. They kept a dog for many months without change as to movement, without giving it any special exercise, but varied its food immensely, and they found the urea eliminated was almost in proportion to the amount of nitrogen taken in the food. But inasmuch as the then existing view required this to be connected in some way with the exercise of force, they explained that so much more force was exercised in the actions within the body in dealing with the increased amount of nitrogenous substance consumed; so that after all the amount of the urea eliminated was a measure of the exercise of force, although it was in these internal actions, and not in the voluntary exercise of muscular power. I was in Germany at the time that book came out, and went to Munich, hoping to see these gentlemen on the subject. In conversation with Professor Voit, I ventured to call in question the

conclusion at which they had arrived, and I think he considered I was entirely in error. But a few years afterwards it was found by others also that the amount of urea eliminated had no direct connection with the amount of force exercised, and that what is the most pronounced when there is an increased exercise of force, is an increased elimination of carbonic acid by the lungs. I believe there is now no doubt about that matter. Messrs. Fick and Wislicenus, Dr. Frankland, and Dr. E. Smith, brought that prominently forward, and I believe it is now accepted that the elimination of urea is no measure of the muscular force exerted within the body.

After putting forward these views, Messrs. Bischoff and Voit put their dog into a kind of tread-wheel, and they found that the amount of urea eliminated was not in proportion to the exercise of force, but the amount of carbonic acid was so, and eventually they themselves admitted the truth of this.

Then came the question of the sources of animal fat. On this point, again, Voit has worked almost exclusively with the dog, which is a carnivorous, or, at most, an omnivorous animal. He has found, which I do not wish to call in question, that in the case of the carnivora, and in some cases of the herbivora, the fat may be formed from the nitrogenous substance of the food. But from the results obtained with this carnivorous animal he has come to the conclusion, that not only in such cases, but in all, the fat formed within the animal is derived from the albuminous substance of the food or of the body. I have roughly noted a few of the experiments of Voit, which I believe are the strongest or most conclusive for his view of the question. He found that when a dog was fed on starch or sugar alone, or with albumin, or with fat and albumin, the carbon stored up, that is to say, the carbon which was not eliminated in any way from the body, was never more than that in the fat of the food, *plus* that in the albumin which was broken up, as indicated by the amount of urea eliminated. He concluded that this was a proof that fat was not formed from the carbohydrates. In another case, which perhaps was stronger, he fed a dog with starch and a little fat, but no albumin whatever, and the carbon stored up was equal to that of the fat in the food, *plus* that due to the oxidation of albuminous tissue, and when he gave more starch in this food the amount of carbon stored up was reduced; that is to say, he argues that the carbohydrates in this case protected the albumin of the body from disintegration, and did not in any way serve for the production of fat; and that there would have been a greater storing

up of carbon if this additional starch which he gave to the animal had been the source of the fat. He also argued, from a number of experiments, that starch and sugar are quite oxidised in the body yielding carbonic acid, &c., within twenty-four hours. He maintains that the same must occur with herbivora as with carnivora. The carnivora are found absolutely to digest vegetable food, and take into their system as an herbivorous animal; and he argues that, to establish a different source of fat, it must be shewn by experiment that fat is formed in excess of that in the food, *plus* that which can be formed from the oxidated albumin. Now this, I think, I shall be able to show you we have done. We have not accepted the challenge the way of making new experiments for the purpose, but I think we have old experiments which are perfectly conclusive, and do meet exactly the requirement which Voit says is essential to disprove the view which he maintains with regard to the herbivora.

But before entering on our own experiments, I will just say what happened in answer to the challenge in Germany. Weiske and Wildt conceived, as I shall be able to show afterwards was a very right thing to conceive, that the pig was the very best animal to experiment on for this purpose. He is certainly *the* fat-maker of all the animals that feed; and there are other reasons why he is the best of all others to experiment upon in this particular. They had, from a theoretical point of view, a very good conception of what was necessary. They took four pigs, slaughtered two of them, and determined the fat and other constituents in those animals. Then they fed one on food very poor in nitrogenous substance, and one on food exceedingly rich in nitrogenous substance. It happened that the pig fed on food very rich in nitrogen had so much that it became unwell, and that the experiment failed entirely. With regard to the one fed on food poor in nitrogen the food was so poor that the experiment took too long a time; in fact, too much food was passed through the body in proportion to the increase produced; and when eventually they slaughtered that animal and analysed it, so much nitrogen had passed through the body during the time, that they found the whole of the fat that had been formed might be derived from the nitrogenous substance consumed. Weiske and Wildt did not conclude therefrom that it was established that fat could only be produced from the nitrogenous substance, but they admit that the experiment was not conclusive.

In the experiments of Mr. Lawes and myself we have used a great many animals, and we have brought our results into calculations

although the experiments were not at the time arranged with the special view of determining this question. The table shows some results of experiments with sixteen oxen, 249 sheep, and fifty-nine pigs. You will see that the proportion of stomachs and contents in the body is 11·6 per cent. with the oxen, 7·5 with sheep, whilst it is only 1·3 in the pig. The intestines and contents, on the other hand, shew in oxen only 2·7, in sheep 3·6, and in the pig 6·2 per cent.; with it, therefore, very much more than with either of the ruminant animals. We know that the character of the food is such in the case of the ruminants that they must pass an enormous quantity of very crude stuff through their bodies, and must elaborate it first in one stomach and then in another, and the result is they have not only a very large capacity of stomach, but also a very large proportion of contents in relation to the whole body. In the case of the pig, on the other hand, the stomach is exceedingly small; the natural food of the pig is starchy seeds or roots (which are the food of man also), it contains exceedingly little necessarily effete matter, their stomachs have comparatively manageable stuff to deal with, and they have a very small stomach, while on the other hand their intestines are very large. It is known that the transformation of the starch goes on almost throughout the intestinal canal, so that we can easily understand how it is that with such starchy food these animals have an enormous amount of intestines compared with either oxen or sheep. If we look at the proportion in the live weight of the, so to speak, further elaborating organs the heart, the liver, the lungs, the pancreas, and so on, their percentage by weight in the bodies of the three descriptions of animals is almost identical.

Now, for 100 lbs. of live weight the amount of dry substance consumed per week was 12·5 by oxen, 16 by sheep, and 27 by pigs; that is : 100 lbs. live weight of pig will consume much more dry substance of food, and, as I have stated, that food is of a more highly nutritive kind, and more easily digested, than that of oxen or sheep. Again, the increase per week was only 1·13 per cent. on the live weight of oxen, 1·76 of sheep, and 6·43 of pigs. So that the proportion of the increase to the weight of the body is much the greatest with the pig. Then, if we take the facts in relation to the amount of the food, for 100 lbs. of dry substance of food, the ox will give in increase only 5·2 of fat, the sheep 7, and the pig 15·7. Suffice it to say, that there is less effete matter in the food of the pig, and therefore its live weight and its increase indicate more nearly the real increase of the body, and

not the fluctuating matters in the alimentary canal. Its food is of a higher character, so that a larger proportion of it is stored up. That which passes through the system is more completely used, and the amount of fat which is produced is also very much higher. Therefore, I say the pig is by far the best animal to experiment upon for this purpose.

Whilst on this subject I may refer to a portion of the table which vegetarians will perhaps not be much pleased to see. If we are to judge that the size of the stomach indicates to some extent the character of the food, its crudeness or concentration, as no doubt is the case with the other animals, and if we compare oxen, sheep, pigs, and man, we find the proportion of stomach by weight per cent. is, approximately, in oxen 3·2, in sheep 2·44, in pigs about 0·88, and in man only 0·38; so that going from one animal to the other you should have more concentrated and more digestible food in the case of man, than of the pig; and you have animal food as well as starchy seeds, roots, &c.; and the indication is, I think, that man was not made to consume potatoes and cabbages by the bushel.

The next point is as to the indications of merely practical results. Without going into the chemistry of the subject, or discussing whether the food of the animal does contain enough or not enough of nitrogenous substances to yield all the fat produced, I will call attention to some results which will indicate the general relations of the food to the necessities of the body. On the coloured diagram you have the results of thirty separate experiments on pigs. The plan was this: we gave to a certain set a fixed amount of highly nitrogenous food, and let them take whatever they liked of less nitrogenous food. To another set we gave a fixed amount of food low in nitrogen and rich in starch and such matters, and let them make up whatever they wanted with highly nitrogenous food. So we rang the changes in a great many more cases than are here represented, but in this way it will be seen that the animal fixed its own diet according to the necessities of the case; and the question is, was it the nitrogenous substances, was it the non-nitrogenous substances, or was it the total dry substance, nitrogenous and non-nitrogenous together, which guided the amount consumed by a given live weight within a given time, or rather guided—for these were fattening animals—the amount of increase which was produced? The lowest amount of nitrogenous substances consumed by 100 lbs. of live weight of pig per week in any one experiment being taken as 100, in some cases 300 were taken, and

most more than 200. In the same way the lowest amount of non-nitrogenous substance being taken as 100, in no case was nearly as much as 200 consumed, and the average was about 140 parts. When we come to the total dry substance, including both nitrogenous and non-nitrogenous, we find that the quantities ranged more closely together; that is to say, the total digestible organic substance seems to have been the measure of what was required, and that the nitrogenous substance might possibly act for the non-nitrogenous substances if there were enough of them. But it is quite clear that the measure was either non-nitrogenous substances or the total organic substances—certainly not the nitrogenous substances. Then the question arose, whether the same thing would hold in relation to the amount of increase in the weight of animal produced. It was always assumed, until these experiments of Mr. Lawes and myself, that when animals were not fed on highly nitrogenous food the amount they put up with was comparatively small. These experiments show the amount of these three classes of constituents consumed in producing 100 pounds increase of live weight in the different cases. 100 pounds being the lowest amount of nitrogenous substance required, was the highest, the animal fixing his own diet, and in many cases it was over 200; that is to say, more than twice as much as sufficed him when he had enough of other matters to make up. At all rates it would seem that fat can be formed from nitrogenous substances, provided there is a deficiency of non-nitrogenous substances in the food; and I may say that the nitrogenous substances are of a higher food capacity, irrespective of the nitrogen, containing more carbon, more hydrogen, and less oxygen; they have more usefulness in them than an equal weight of starch or any substance of this kind.

Now the question arises, what is the state of affairs when we attempt to calculate these results and to see whether or not the food did contain enough nitrogenous substances, or albuminous matter, to supply the whole of the fat produced? These experiments were not specially arranged to settle that question, but they were calculated afterwards. It was about twenty-six years ago that we took two pigs of the same sort, very carefully selected both by practical and scientific eyes as nearly as possible exactly alike. One was slaughtered, and the total amount of dry matter, fat, nitrogen, mineral matter, and so on determined; and then the other animal was fed. At that time we did not arrive at such distinct conclusions as we did afterwards as to

the desirability of giving a greater proportion of starchy matters. We gave the animal a great deal more than the proportion of nitrogenous substances existing in what may be called the normal fattening food of the pig—barley meal. In the first column of the table, the results of that experiment are calculated out to show whether the food did contain enough nitrogenous matter to yield the fat produced. You will see the proportion of non-nitrogenous matter to one of nitrogenous is 3·6. Now, the proportion in barley meal, which is the best fattening food for the pig, is between 5 and 6 to 1; so that we gave too much nitrogen according to what we now know is the best proportion. There was a considerable amount of increase in ten weeks, eighty-eight pounds, or 85·4 per cent. on the original weight of the body.

The question is, how much fat was in the food? and that is shown in the second division of the table. It is calculated that for 100 pounds increase in the live-weight there were stored up 63·1 pounds of fat. There were of ready-formed fat in the food 15·6 pounds; leaving 47·5 pounds fat to be produced from some material or other. Out of 100 of nitrogenous substances consumed as food, there were stored up in increase 7·8, leaving 92·2 parts of nitrogenous substance which might be used for the production of fat or might not. If we calculate how much carbon there was in the produced fat, and how much there was left available in the nitrogenous substance for the production of fat, we find that there were 7·4 pounds more carbon possibly available from the nitrogenous substance than was necessary for the production of the fat; or, put in another way, there were 120 of carbon available from the nitrogenous substances for 100 required.

According to this mode of calculation, therefore, there was enough nitrogenous substance to justify the conclusion of Voit; or rather, the result does not in any way disprove his conclusion that fat has been produced from the disintegration of nitrogenous substances in the body. This table was calculated some years ago, and we have intentionally put the results in the worst aspect that we could for our own side of the case, that we might not exaggerate the conditions. For instance, we have assumed that the whole of the fat in the food would be taken up, which it certainly would not; and we have assumed that the whole of the nitrogenous substances of the food would be digested, and would come into play, which they certainly would not. If we assume in our own calculations the estimate adopted in Germany, that 100 pounds of nitrogenous substance cannot yield more than

me of fat, even this experiment shows a little deficiency of nitrogenous substances, and would in fact be in favour of our view.

The next two experiments given in the table show a still higher portion of nitrogenous substance in the food; and there was, accordingly, a great deal more carbon available from the nitrogenous substance than was necessary for the formation of the amount of fat produced.

The next two experiments (four and five), were with more natural feeding food of the animal, one entirely Indian corn-meal, and the other entirely barley-meal. A pig requires for rapid fattening very little, if any, more nitrogenous substance than this represents. But we have only 60 per cent. or a little over, 60·8 in one case, and in the other, of the carbon of the fat produced in the animal, possibly derivable from the nitrogenous substance of the food. So that we find in those two cases nearly 40 per cent. of the carbon of the produced fat which could not possibly come from the nitrogenous substances, must have come from the non-nitrogenous matter, in fact from carbohydrates. But an objection may be raised to this calculation—the animals were larger to begin with; and the weights were lower at the end; so that the composition of the lean animal, and of the fat animal, as derived from the direct analyses, does not absolutely agree; but we could not possibly thus get rid of this forty or more per cent. which the calculations would show to be derived from the nitrogenous substance of the food.

The remaining four experiments are also entirely in favour of our view.

The animals were about the same weights as those analysed: the food was more nearly the proper food for fattening, being rather richer in nitrogenous substances, but much higher than in experiments 1 to 5. But even here we found 18·9, 18·8, 25·2, and 14·1 per cent. of the total carbon of the produced fat could not possibly have been derived, and certainly a great deal more was not derived, from the nitrogenous substances of the food.

I need not trouble you further with these results. But I should state that the contrary view has been adopted not only by some physiologists, but in Germany in some text books on agricultural chemistry. I find in my hand one of these text books in which the evidence of the experiments is discarded, and it is assumed that if you cannot experiment with the respiration apparatus the results are good for nothing. I would not wish to depreciate the importance of the results obtained by the respiration apparatus in any way. I have taken the

greatest interest in them, and I think they lead to the most important conclusions; but I also think some observers have come to very erroneous conclusions from the results of such experiments. I submit that if you experiment with *the* fat-producer—the pig—and if you take two carefully selected animals (or more if you like) kill and analyse one, and feed the other as rapidly as possible, that is, let him take as much of the most appropriate food as he will take, you may, without any respiration apparatus, determine this point. It is most important that it should be definitely settled. Since the recent publications on the subject, Mr. Lawes and myself have gone thoroughly into the question, and re-calculated most of our results; those relating to oxen and sheep as well as pigs. They point to this: that the ruminant animals, which have such elaborate machinery, and do so little productive work, do pass so much nitrogenous substance through the body in relation to the amount of increase, that they do not show that fat can be derived from the non-nitrogenous substances of the food; but in the case of pigs the evidence is perfectly conclusive. Having re-calculated our own experiments in this way, and the results being absolutely conclusive so far as the pig is concerned, Mr. Lawes is unwilling to be at the trouble and expense of further experiments on the question; but it really is one of great importance, and one which public institutions might well take up. It is of importance, not only agriculturally, with reference to the proper way of feeding stock, but also in its bearings on the nutrition of man.

[For the tables and diagrams referred to above, see—"On the Sources of the Fat of the Animal Body," *Philosophical Magazine*, December, 1866; and—"On the Formation of Fat in the Animal Body," *Journal of Anatomy and Physiology*, Vol. xi., Part iv.; and for other points, and detail—"Food in its Relations to Various Exigencies of the Animal Body," *Philosophical Magazine*, July, 1866; and the papers therein referred to.]

ON
THE FORMATION OF FAT
IN
THE ANIMAL BODY.

BY
J. B. LAWES, LL.D., F.R.S., F.C.S.,
AND
J. H. GILBERT, PH.D., F.R.S., F.C.S.

FROM THE JOURNAL OF ANATOMY AND PHYSIOLOGY,
VOL. XI., PT. IV.

CAMBRIDGE:
PRINTED AT THE UNIVERSITY PRESS.
1877

Cambridge:

**PRINTED BY C. J. CLAY, M.A.
AT THE UNIVERSITY PRESS.**

ON THE FORMATION OF FAT IN THE ANIMAL BODY¹.

PLATE XXII.

FORMERLY it was supposed that the fat of the Herbivora was derived exclusively from ready formed fat in their vegetable food. Liebig showed that this could not be the case; and he attributed much of the fat of the animal body to the carbohydrates of the food. His views on the point were at first called in question by Dumas, Boussingault, and others, but afterwards accepted. Our own very numerous feeding experiments, commencing about 30 years ago, together with a careful consideration of the experience of practical feeding, afforded strong confirmation of Liebig's conclusions; and more especially in 1852², and subsequently, we pointed out the bearing of the results on the question.

At a meeting of the Congress of Agricultural Chemists held at Munich, in 1865, Professor Voit combatted this view. From the results of experiments with dogs, made in Pettenkofer's respiration-apparatus, he maintained that fat must have been produced from the transformation of nitrogenous substance; and further, that this was probably the chief, if not the only, source of the fat, even of Herbivora. In 1869 he elaborately argued the point, not only in reference to the results of his own experiments with dogs and with cows, but also to the records of those of various other experimenters, with various descriptions of animal; and he has subsequently made further contributions on the subject, conjointly with Professor Pettenkofer.

Their results, obtained with a dog, and their conclusions drawn from them, were to be described in a few words, they might perhaps be so as follows:—When a dog was fed on starch or sugar, alone, or with albumin, or with fat and albumin, the carbon stored up was not more than that in the fat of the food

¹ The substance of this paper was given by one of us at the *Naturforscher Versammlung* (Section für Landwirthschaft- und Agricultur-Chemie) at Hamburg, in September, 1876.

² On the Composition of Foods in relation to Respiration and the Feeding of Animals. *Report of the British Association for the Advancement of Science, for 1872.*

plus that which could be derived from the albumin broken up. There was, therefore, no proof that fat can be formed from the carbo-hydrates. Again, when a dog was fed with starch and a little fat, but no albumin, the carbon stored up was equal to that of the fat of the food *plus* that derived from the transformed nitrogenous substance of the body. More starch reduced the amount of carbon stored up; the carbo-hydrate having protected the albumin of the body from oxidation, and thus limited the formation of fat. They never found fat formed from starch or sugar. They maintain that the same must occur with the Herbivora; and that to establish the formation of fat from the carbo-hydrates, experiments must be brought forward in which the fat deposited is in excess of that supplied by the food *plus* that which could be derived from the transformed albumin.

This, many of our own experiments with pigs do clearly show; and in 1866 we published a short paper¹, in which we illustrated the bearing of some of them on the point. In his paper in 1869², Professor Voit quotes some of those results, and admits that in the experiments in which there was only a medium albuminous supply in the food, there was, as the figures stand, a considerable deficiency for the formation of the fat produced, and a greater deficiency still in the cases in which the relation of the nitrogenous to the non-nitrogenous constituents was such as experience has shown to be the most favourable for pig-fattening; and that, therefore, a considerable amount of fat would, in these instances, appear to have been derived from the carbo-hydrates. Still, Professor Voit says he cannot allow himself to consider a transformation of carbo-hydrates into fat to be proved thereby. He confesses that he has not been able to get a general view of the experiments out of the mass of figures recorded, and suggests several possible sources of error, his reference to some of which shows that he has in fact misunderstood them. At the same time, he proposed that new experiments with geese and with pigs should be made, in order to arrive at a final decision on the question; and in a very recent

¹ On the Sources of the Fat of the Animal Body. *Philosophical Magazine*. December, 1866.

² *Zeitschrift für Biologie*. Band v.

conversation with one of us, he expressed his willingness to undertake a conclusive experiment with pigs.

Weiske and Wildt¹ undertook an investigation to determine the point; which, from a theoretical point of view, was well conceived; but which did not succeed, owing to the oversight of the conditions indicated by experience as essential to the rapid fattening of the animal. They selected four pigs; two were slaughtered to determine the initial composition; one was fed on food so rich in nitrogen that it suffered in health, and the experiment had to be discontinued; the other was fed on food so poor that it fattened extremely slowly; and hence, at the conclusion, calculation showed that there was not enough of the consumed nitrogenous matter available for fat-formation to cover the whole of the fat which had been produced.

Thus, it has been concluded from experiments with animals which are not preeminently fat-producers, that fat is probably either formed in the body from the carbo-hydrates; and some of the experiments with more suitable animals have been, to say the least, inconclusive. Further, it seems to be assumed, that absolute proof on the point can be obtained without the aid of a respiration-apparatus. These views, moreover, have already been adopted, not only by some physiologists, but in the text-books treating of the application of chemistry to the feeding of the animals of the farm. Thus, Professor Emil Wolff, in his *Landwirthschaftliche Fütterungslehre*—although he admits that the amounts of increase produced in relation to constituents of food consumed, which it is established by common observation may be obtained with pigs, and still more those recorded in the direct experiments with those animals, are almost inconceivable without assuming the direct concurrence of the carbo-hydrates in the formation of fat—nevertheless, seems to consider that evidence of the kind in question, and we suppose our own, therefore, is inconclusive. He says that exact respiration-experiments are still wanting; and he suggests that accurate respiration-experiments with pigs should be made, to settle definitively whether the carbo-hydrates, as well as albumin, contribute directly to the formation of fat in the animal body.

¹ *Ib.* Band x.

Since the appearance of Professor Emil Wolff's work, and the publication of the negative results of Weiske and Wildt, we have carefully reviewed and recalculated many of the results of our feeding experiments, including those with oxen, with sheep, and with pigs; in order to satisfy ourselves whether any doubt could be entertained of the views we have previously advocated; and whether, therefore, it was at all incumbent upon us to institute new experiments on the point. The result of this examination, so far as the ruminant animals are concerned, has been to show that, owing to the comparatively small amount of increase obtained with them from a given amount of constituents consumed, the quantity of nitrogenous substance passed through the system for the production of a given amount of increase was, in most, if not in all cases, so large as, in the absence of proof to the contrary, to admit of the assumption that the whole of the fat formed had its source in transformed nitrogenous matter. At any rate, no absolute proof of the derivation of fat from the carbo-hydrates can be obtained from data of the kind in question relating to such animals. In deciding the point in regard to them, the evidence afforded by the analysis of the fæces and of the urine, and by the determination of the products of respiration, must also be brought into consideration. It was quite otherwise, however, in the case of our experiments with pigs; in many of which much more fat was produced than could possibly have been derived from transformed albumin of the food. We concluded, therefore, that we were in no way called upon to institute new experiments, and decided, instead, again to direct attention to the results quoted in the short paper on the subject published in 1866, as already referred to.

The figures given in Table I. of that paper show how much smaller is the proportion of alimentary organs and contents in a given live-weight of the pig than of either oxen or sheep; that, in proportion to a given live-weight, the pig consumes a very much larger quantity of dry substance of food within a given time (whilst his food contains a very much larger proportion of digestible, and therefore, very much less of necessary effete matter); that he gives several times as much increase in relation to a given live-weight within a given time; much more increase in relation to a given quantity of dry substance

od; also a larger proportion of fat in that increase. Further, the most appropriate fattening food of the pig contains a larger proportion of readily digestible carbo-hydrates than that of the ruminant animals. All these conditions indicate the pig to be the most suitable animal for the determination of the matter in question.

The results selected to illustrate the main point are given in Table II. of the same paper. They were all obtained more than 20, and some more than 25 years ago; and the rations were not arranged with a special view to the settlement of the question; but to determine the relations of the different constituents of food to various exigencies of the body, and the amount, and the proportion of different foods which were the most favourable for the feeding of the animals. Accordingly, the series included proportions varying from 2.0 to 6.6 parts of non-nitrogenous to 1 of nitrogenous substance in the food.

In experiment 1, two animals were selected, of the same age, and as nearly as possible alike both in character and weight; the weight of the one being 100 lbs., and that of the other 103 lbs. One was slaughtered at once, and its contents of nitrogenous substance, fat, mineral matter, &c., accurately determined. The other was fed on a mixture consisting of bean-meal, lentil-meal, and bran, each one part, and barley-meal three parts, given *ad libitum*, but accurately weighed, for a period of ten weeks, when it had nearly doubled its weight. The

animal contained, however, a considerably higher proportion of non-nitrogenous to non-nitrogenous constituents than is recognised as the most favourable for the fattening of the pig. The animal was then slaughtered, and analysed, as the other had been. The composition of the food was also determined by analysis. The experiment afforded, therefore, reliable data for determining the amounts of fatty and nitrogenous substance consumed, the amount of nitrogenous substance stored up in the animal, and also the amount of fat stored up.

Eight other experiments were quoted, in each of which a different food-mixture was employed, and in each of which the animals were fed, in some cases for a period of eight, and in others of ten weeks. The average live-weight per head at commencement was, in these eight experiments, respec-

tively, 143, 147, 144, 149, 95, 95, 94, and 97 lbs. Thus, in the first four cases, the average initial weight per head was notably more than that of the two animals of experiment 1; but in the last four experiments it was very nearly the same. In the calculations, the percentage composition of the animals in experiments 2—9 was assumed to be the same at the commencement as that of the unfattened animal in experiment 1, and the same at the conclusion as that of the fattened animal in experiment 1. It was quite obvious, during the progress of the experiments, that the animals having the higher proportions of nitrogen in their food, grew more, and fattened less, than the others; and careful observations, made after slaughtering, entirely confirmed this. The tendency to error in the calculations would be to indicate too low an amount of nitrogenous substance, and too high an amount of fat stored up in the cases with the higher proportions of nitrogenous substance in the food, and too high an amount of nitrogenous substance, and too low an amount of fat stored up with the lower proportions of nitrogenous substance consumed. The range of the probable error here supposed is, however, not such as at all to throw doubt on the validity of the main conclusions which are drawn from the figures as they stand.

A comparison of the amount of ready-formed fat in the food, with that of the determined or estimated total fat stored up in the increase of the respective lots of animals, showed that, even supposing the whole of that consumed had been retained, there remained from two-thirds to nine-tenths of the total amount stored up to be otherwise accounted for. It must have been produced within the body.

The next question was, whether this large amount of produced fat could possibly have been derived from the nitrogenous constituents of the food? or whether it must of necessity have had its source, in greater or less proportion, in the carbohydrates at the same time supplied?

Deducting from the total amount of nitrogenous substance consumed, the small amount estimated to be stored up as such in the increase of the animal, there remained a large proportion available, it may be, for the formation of fat, with other products. In order to give to the nitrogenous substance of the

food not stored up, its fullest possible (and even more than its fullest) value for fat-formation, the whole of its carbon, *minus* that which its nitrogen would require to form urea, is, for the sake of illustration, assumed to be available for fat-formation.

So calculated, the result in experiment 1, and also in two of the other cases in which the proportion of nitrogenous to non-nitrogenous substance in the food was considerably higher than is recognised by experience as the most suitable in the fattening food of the pig, was that more nitrogenous substance was available for fat-formation than was necessary to supply the estimated amount of produced fat. In the cases in which the nitrogenous substance was not so excessive, but still more than is the most appropriate, there was a considerable proportion of the total produced fat which could not possibly have been derived from the nitrogenous substance of the food. Lastly, when the proportion of the nitrogenous to the non-nitrogenous substance in the food was the most appropriate for fattening, there was a much larger proportion (about 40 per cent.) of the total produced fat, which could not possibly have had its source in the nitrogenous substance consumed.

Striking as are these results, it is obvious that a still larger proportion of the produced fat would appear to be formed from the carbo-hydrates, if it were assumed, with Henneberg and Voit, and as is doubtless nearer the truth, that 100 parts of albumin will not yield more than 51·4 parts of fat, instead of, according to the foregoing illustration, about 61 parts.

It will be well, however, briefly to consider, whether an amount of error in the estimates, which would turn the scale, and show that the whole of the produced fat might be derived from the nitrogenous substance of the food, is at all conceivable, at any rate in the cases in which the proportion of the nitrogenous to the non-nitrogenous constituents consumed was the most nearly that which is recognised as the most favourable for pig-fattening, and in which the largest amount of formation from the carbo-hydrates is indicated.

Obviously, the most important point to consider is the range of error admissible in the estimation of the fat stored up in the increase of the animal.

It would be necessary to reduce the estimate of the amount of fat stored up by more than 30 per cent. to bring it low enough to be covered by the fat in the food, *plus* that derivable from the transformed nitrogenous substance, leaving all the other calculations the same. If, however, we were to assume that 100 nitrogenous substance yielded only 51.4 fat, it would be requisite to reduce the estimate of the fat in the increase by more than 40 per cent., to reverse the indication. This is on the assumption that the whole of the fat of the food was stored up in the animal, which would certainly not be the case. It is also on the assumption that the whole of the nitrogenous substance of the food, not stored up as such in the increase, was digested, and available for transformation into fat, &c., but this again is certainly not the case. According to our own experiments, it may be supposed that, with a pig feeding exclusively on good barley-meal, about one-sixth of the total nitrogen voided would be in the fæces. But if it be assumed, according to the estimates of E. Wolff¹, that 20 per cent. of the nitrogenous substance, and 32 per cent. of the fat of the barley, would be voided undigested, and therefore without contributing to the deposition of fat, our estimate of the amount of fat stored up in the increase would have to be reduced by more than 55 per cent., or considerably more than half, to bring it within the amount derivable from the resorbed fat, and the transformed nitrogenous substance of the food.

It is submitted that a range of error in our estimates, at all approaching even the lowest of those above assumed for the sake of illustration, is simply impossible. It is further submitted, with the utmost confidence, that such is the wide margin in the case of pigs fattening rapidly on their most appropriate fattening food, that the question of whether or not the carbo-hydrates contribute to fat-formation may be conclusively settled by a properly conducted experiment with those animals, without any analysis of the fæces or the urine or any determination of the products of respiration. To this end, we would suggest that two animals be selected, of a breed of good fattening quality, and as nearly as possible alike in characters and in weight. A convenient size and weight would

¹ *Landwirthschaftliche Fütterungslehre*, Appendix, Table I.

say about 90 lbs. per head. Let each be fed with ground barley of good quality, giving it, by degrees, as much as it will consume, until both weigh about 100 lbs. Then slaughter one, and determine its total amount of nitrogenous substance, fat, &c. Feed the other in the same way, that is with barley-meal and water) exclusively, as much as it will consume, until it reaches about 200 lbs. in weight. Then slaughter and analyse as the first. The quantity and composition of the food first, of course, also be determined. Such an animal would consume somewhere about 500 lbs. of barley, more or less, and increase from 100 lbs. to 200 lbs. in live-weight, in from 8 to 12 weeks, more or less, according to quality of the animal, quality of the food, &c. &c. It is desirable that the animals selected should have been feeding on fairly good food previously, so that the transition to full fattening food should not be too sudden. It is also, of course, desirable, that the experiment should be made in duplicate if possible.

But, independently of the results of any such experiments, may be asked, what is the lesson of common experience on this matter? We say, unhesitatingly, that the experience of the feeding of animals fully confirms our view.

In reference to this point we would call attention to the coloured diagrams Pl. xxii. which show the proportions of nitrogenous substance (black), of non-nitrogenous substance (yellow), and of total organic substance, nitrogenous and non-nitrogenous together (blue), respectively:—

I—consumed per 100 lbs. live-weight per week,

II—consumed to produce 100 lbs. increase in live-weight,

the case of thirty different feeding experiments with pigs, 15 of which comprised not less than three and some four animals, and in each of which they fixed their own consumption¹. That is to say, various current foods, but containing widely different percentages of nitrogenous substance, being selected, one (or a mixture) of high, or of medium, or of low

¹ "Pig Feeding;" *Jour. Roy. Ag. Soc. Eng.* Vol. xiv. Part 2; Experiments 8, and 12, Series 1; Experiments 1—12, Series 2; Experiments 1—5, Series also "On the Equivalency of Starch and Sugar in Food," *Report of Brit. Ass.* 1854; Experiments 1—4. See also "Experimental Enquiry into the composition of some of the Animals fed and slaughtered as human food." *Phil. Mag.* 1859, Part 2.

percentage of nitrogen, was given, *ad libitum*; or a fixed quantity of one or more was given, and another given *ad libitum*; and so on. In this way the animals fixed their own consumption, and from the results it may be judged by what requirement this was guided.

First, as to the consumption by a given live-weight within a given time; which of course met the collective requirements for both sustenance and increase. Diagram I illustrates this point. The lowest amount of nitrogenous substance so consumed in any one of the thirty experiments is taken as 100; and it is seen that the amount of it consumed ranged, among the thirty dietaries, from 100 parts to more than 300; and it averaged more than 200. Reckoned in the same way, the consumption of non-nitrogenous substance varied from 100 to only 177 parts, and averaged only 141 parts. Again, reckoned in the same way, the consumption of total organic substance (nitrogenous and non-nitrogenous together) ranged from 100 to only 150 parts, with an average of 125 parts.

Secondly, as to the amounts consumed to produce 100 lbs. increase in live-weight. Diagram II shows that, for this result, the consumption of nitrogenous substance ranged from 100 to 282 parts; and it averaged 173 parts. That of the non-nitrogenous substance ranged from 100 to only about 140 parts, with an average of 124 parts; and that of the total organic substance (nitrogenous and non-nitrogenous together) from 100 to only 147 parts, with an average of 122 parts.

It should be explained that, as in the Tables and Diagrams given in the original papers above referred to, the *total* amount of nitrogenous and of non-nitrogenous substance, in the different foods, are taken as the basis of the calculations; no deduction being made for "indigestible" matter; nor is the fat in the food reckoned at any higher value than the other non-nitrogenous constituents. This plan was adopted as best representing the facts actually determined by analysis; but attention was at the same time directed to the varying amounts of indigestible matter in the different foods, and to the greater or less amount of fat which they contained. We have, however, quite recently recalculated the whole of the experiments, making deduction for indigestible or undigested matter, according to E. Wolf's

able already quoted, and with him multiplying the amounts of fat by 2.5, and have constructed Diagrams according to the data so obtained. These still more strikingly illustrate the point in question than the Diagrams herewith given; that is to say, they show a wider range in the amounts of the nitrogenous substance consumed in the different experiments, a less variation (excepting in one case in which there was much fat) in the amounts of the non-nitrogenous substance consumed, and especially a less range in the amounts of the total organic substance consumed. The two methods of calculation show, however, in most of the cases, much less difference in the relation of the nitrogenous to the non-nitrogenous constituents than might have been anticipated. With this explanation, we still adhere to our original plan of calculation, rather than adopt corrections based upon factors as yet not sufficiently established¹. At the same time, we repeat that the points here indicated could be considered in judging of the results as they stand.

It is then perfectly clear, that neither the amount of food consumed in relation to a given live-weight within a given time (which of course covered the requirements for increase as well as sustenance), nor the amount taken to yield a given amount of increase in live-weight (which in its turn covered the requirements for sustenance also), was at all in proportion to the amount of the nitrogenous constituents it supplied. It is quite obvious, that the consumption, both for sustenance and for increase, was much more nearly in proportion to the amount of digestible non-nitrogenous constituents supplied; but it was more nearly still guided by the amount of the total digestible organic substance—nitrogenous and non-nitrogenous together—which the foods contained.

That the great variation in the amount of nitrogenous substance consumed was not due to a deficiency of it in most of the foods employed, is shown by the fact that it was in the experiment in which the food contained the lowest proportion of it, that the smallest amount of nitrogenous matter was actually consumed in relation to a given live-weight within


¹ Professor Emil Wolff has recently determined the proportions undigested of the different constituents of cocoa-nut cake, barley-meal, maize-meal, and rye-meal, in actual experiments with pigs. *Versuchs-Stationen Organ.* Band 1, No. 4, 1876.

a given time, but was required to produce a given amount of increase. It is obvious, that where two or three times as much nitrogenous substance was consumed, it was much in excess of the normal requirement. In fact, the animals consumed almost regardless of the amount of nitrogenous substance supplied, until they had obtained a sufficiency of non-nitrogenous, or of total organic substance. It is further obvious, that the range of variation in the amounts of non-nitrogenous constituents consumed would have been very much less, but for the very variable amount of nitrogenous substance necessarily taken with it, the variable amounts of fat in the foods, and the greater amount of indigestible matter in some of them than in others. The indication is, indeed, that the excess of nitrogenous substance consumed substituted a certain amount of non-nitrogenous constituents; that, in fact, within certain limits, the two classes of constituents may, for the purposes of respiration and fat-formation, mutually replace each other.

Lastly on this point, not only did neither the amount of food consumed, nor the amount of increase in live-weight yielded, bear any relation to the amount of nitrogenous substance supplied, but the more excessive the supply of it the greater was the tendency to grow, and the less the tendency to fatten. There is, of course, a point below which the proportion of nitrogenous substance in the food should not be reduced, but if this be much exceeded, the proportion of the increase, and especially of the fat-increase, to the nitrogenous substance consumed, rapidly decreases; and it may be stated generally, that taking our current fattening food-stuffs as they are, it is their supply of digestible non-nitrogenous, rather than of nitrogenous constituents, which guides the amount, both of the food consumed, and of the increase produced, by the fattening animal.

In conclusion, we repeat that, in many of our experiments with pigs, much more fat was produced than could possibly have been derived from the albumin of the food, and hence the carbo-hydrates must have contributed directly to its formation; further, that experience in practical feeding is entirely in accordance with our views on the point.







Supplement to former Paper entitled—'Experimental Inquiry into the Composition of some of the Animals Fed and Slaughtered as Human Food,'—Composition of the Ash of the Entire Animals, and of certain Separated Parts." By Sir JOHN BENNET LAWES, Bart., LL.D., F.R.S., F.C.S., and JOSEPH HENRY GILBERT, Ph.D., LL.D., F.R.S., V.P.C.S. Received June 11, 1883.

(Abstract.)

In a former paper ("Phil. Trans.," Part II, 1859) the authors had given the actual weights, and the percentage proportion in the entire body, of the individual organs, and of certain more arbitrarily separated parts, of 326 animals—oxen, sheep, and pigs—in different conditions as to age, maturity, fatness, &c. They called particular attention to the wide difference in the proportion by weight of the lungs and intestines in the three descriptions of animal; the proportion of stomach and contents being very much the highest in oxen, considerably less in sheep, and little more than one-tenth as much in pigs as in oxen. On the other hand, the intestines and contents constituted a less proportion to the weight of the body in oxen than in either sheep or pigs; the percentage by weight in pigs being nearly as high as in sheep, and more than twice as high as in oxen. In these very characteristic differences in the proportion of the organs and first laboratories of the food, the other internal organs, especially, as also the blood, contributed a pretty equal proportion to the weight of the entire body, in the three descriptions of animal.

When animals had been selected for the determination of the chemical composition, namely—a fat calf, a half-fat ox, and a fat ox; a fat sheep, a store sheep, a half-fat sheep, a fat sheep, and a very fat sheep; one pig, and a fat pig. In these, in the collective carcass parts, in the collective offal parts, and in the entire bodies, the total nitrogenous substance, the total fat, the total mineral matter, the total dry substance, and the water, were determined; and the results were recorded and discussed in detail.

It was shown that, as the animal fattened, the percentage of nitrogenous substance decreased considerably, whilst that of the fat and of the total dry matter increased in a much greater degree. It was stated that the portions of well fattened animals which would be estimated as human food would contain three, four, and even more parts as much fat as dry nitrogenous substance; and comparing such animal food with wheat-flour bread, it was concluded that, taking into consideration the much higher capacity for oxidation of a given weight

of fat than of starch, such animal food contributed a much higher proportion of non-nitrogenous substance, reckoned as starch, to one of nitrogenous substance than bread. In fact the introduction of staple animal foods, to supplement our otherwise mainly farinaceous diet, did not increase, but reduced, the relation of the flesh-forming material to the respiratory and fat-forming capacity of the food.

Finally, the actual amount, and the percentage, of total ash, in the internal organs, and some other separated parts, were given. It was shown that the percentage of total mineral matter, like that of the nitrogenous substance, decreased, not only in the entire body but especially in the collective carcass parts, as the animals matured. It was the object of the present communication to record the results of the complete analysis of the ashes of the collective carcass parts, of the collective offal parts, and of all parts, of each of the ten animals. Forty complete ash analyses had been made.

As was to be expected, more than four-fifths of the ashes consisted of phosphoric acid, lime, and magnesia; these making up the larger amount in the ash of the oxen, less in that of sheep, and less still in that of pigs. Potash and soda were also prominent constituents. Assuming, for the purposes of illustration merely, that one of phosphoric acid was combined with three of fixed base, the ashes of ruminants showed an excess of base; whereas, according to the same mode of calculation, the ashes of the pigs showed no such excess.

It was, unfortunately, only in the case of the offal parts of the animals that the ash of the chiefly bony, and that of the chiefly soft parts, had been analysed separately. The results showed a considerable excess of acid, especially phosphoric, in the ash of the non-bony portions, presumably, in part at any rate, due to the oxidation of phosphorus in the incineration. In further reference to the point in question may be stated that although the oxen and sheep show a higher percentage of total nitrogenous substance than the pigs, yet the amount of pure ash yielded from the non-bony parts is higher in proportion to that from the bones in the case of the pigs than in that of the ruminants.

Comparing the percentage composition of the ashes of the entire bodies of the different animals, the chief points of distinction were— in the ash of the pigs there is a lower percentage of lime, and a higher percentage of potash and soda, than in the corresponding ashes of the ruminants; there is a somewhat higher percentage of phosphoric acid in the ash of the pigs and of the oxen than in that of the sheep, and there is a higher percentage of sulphuric acid (and somewhat of chlorine also) in the ash of the pigs than in that of the other animals.

A table showing the quantities of total ash, and of each individual mineral constituent, in each of the ten animals analysed was given. Not much stress was laid on the amounts in the particular animals.

analysed; as the actual weights and condition of animals coming under similar designations may vary considerably.

It was of more interest to consider the amounts of the mineral constituents in carcass parts, in offal parts, and in all parts, per 1,000 lbs. fasted live-weight, of each description of animal.

It was shown that a given live-weight of oxen carried off much more mineral matter than the same weight of sheep, and a given weight of sheep much more than the same weight of pigs. With each description of animal the amounts of phosphoric acid, lime, and magnesia, are less in a given live-weight of the fatter than of the comparable leaner individuals. Of both potash and soda, again, the quantity is less in a given live-weight of the fatter animals. The same may be said of the sulphuric acid and the chlorine; in fact, in a greater or less degree, of every one of the mineral constituents.

It was estimated that the loss to the farm of mineral constituents by the production and sale of mere fattening increase was very small. It was greater of course in the case of growing than of only fattening animals. In illustration, the amounts of some of the most important mineral constituents removed annually from an acre of fair average pasture and arable land in various products were compared. Such estimates could obviously be only approximate, and the quantities will vary considerably. With this reservation, it may be stated that, of phosphoric acid, an acre would lose more in milk, and four or five times as much in wheat or barley grain, or in hay, as in the fattening increase of oxen or sheep. Of lime, the land would lose about twice as much in the animal increase as in milk, or in wheat or barley grain; but perhaps not more than one-tenth as much as in hay. Of potash, again, an acre would yield only a fraction of a pound in animal increase, six or eight times as much in milk, twenty or thirty times as much in wheat or barley grain, and more than 100 times as much in hay.

From the point of view of the physiologist, it would doubtless have been desirable that the selection of parts for the preparation and analysis of the ash should have been different, and more detailed. The agricultural aspects of the subject had, however, necessarily influenced the course of the inquiry; and the extent of the essential work had enforced the limitation which had been adopted. The results must be accepted as a substantial contribution to the chemical statistics of the feeding of the animals of the farm for human food.



EXPERIMENTS ON ENSILAGE, CONDUCTED AT ROTHAMSTED;

SEASON 1884-5.

BY

SIR J. B. LAWES, BART., LL.D., F.R.S.,

AND

J. H. GILBERT, LL.D., F.R.S.

This paper is, in the main, a reprint of a series of Articles published in the "Agricultural Gazette," from April 27 to August 10, 1885, as the experiments proceeded; but the order of the sections has been re-arranged, the Tables and text have been revised and corrected, and a more complete summary has been added.

LONDON:
HARRISON AND SONS, ST. MARTIN'S LANE,
Printers in Ordinary to Her Majesty.

1886.



CONTENTS.

	PAGE
1. Introduction	5
2. The construction, and the filling, of the Silos	9
3. Quantity and composition of the Herbage put into the Silos, and of the Silage taken out	10
Silo No. 1 (p. 10)	
Silo No. 2 (p. 15)	
4. The chemical composition of the Silage.....	19
5. Experiments with Fattening Oxen.....	24
6. Food required for mere sustenance, for the production of Milk, and for the production of Fattening Increase.....	28
7. Preliminary Experiments with milking Cows.....	36
8. Plan and arrangement of the experiments with milking Cows	37
9. The results of the Experiments with milking Cows.....	39
10. Characters and composition of the Milk from the different Foods.....	51
11. Summary, and general conclusions.....	55



EXPERIMENTS ON ENSILAGE,

CONDUCTED AT ROTHAMSTED;

SEASON 1884-5.

1.—INTRODUCTION.

According to the "Agricultural Returns," there had, in 1884, been 1183 silos constructed in Great Britain, and in 1885, the number was 1183. It is probable that the greater portion of these are in the hands of tenant farmers, as even at the best of times they are over eager in adopting novelties; and now, when the most depression exists, they may readily hesitate to pay even the cost of erecting upon a building which after trial might be found to be

quite certain that, in some of the statements which have been made by the more enthusiastic advocates of ensilage, there has been exaggeration, and that the revolution which the silo was to bring about in our system of agriculture has yet to take place.

In the early part of 1884, a visit paid to Mr. Gibson's farm at Rothamsted, to show that the process was worthy of careful investigation; Mr. Gibson kindly supplied much valuable information, essential to the investigation as to the points which required to be investigated. Mr. Gibson's system of feeding was, to give the silage mixed with hay when used in this way, with dry food in addition, the milk from the cows yielded by the cows were both of good quality. He had ascertained that clover-silage was superior to that made from grass.

Although Mr. Gibson's experience is of great practical value, it does not afford much help in regard to the important question—what is the value of silage as compared with other foods? A great many points have to be investigated before we can assign to the system of ensilage its proper value in the economy of the farm. We want to know, how good a food silage is generally—and especially the silage made from various crops—both as compared with roots, and with the same crops used in the ordinary way. We further want to know, what are the changes that take place in the pit, and how many pounds of dry matter are taken out for 100 lbs. put in; for neither sour nor sweet silage can be made without the destruction of some of the food.

Any one who has had much experience with cows, and the products of a dairy, will understand at once that where the object of experiments is to ascertain the value of one food compared with another, in the production of milk or butter, the subject is surrounded with difficulties. In the feeding of fattening animals, if a sufficient number of them are under experiment, and the food is carefully weighed, it may be expected that the increase in live-weight will give an approximately correct indication of the feeding value of the foods used. With cows, however, not only has every change of food some influence on the yield of milk, but the same food, given in the same quantity, will produce very different amounts of milk with different cows.

It is also well known that milk may be produced at the expense of the substance of the cow herself. The animal may not be even losing weight for a short time, and still may be parting with solid matter which is replaced by water. This being the case, all experiments over short periods, in which one food is substituted for another, are liable to the serious objection, that it is not possible to say what is the source of the change in the milk production. As every cow has a certain milk producing capacity, which cannot be exceeded by any manner of feeding, the animal may be putting on flesh as well as yielding milk with one food, while it may only yield milk without putting on flesh with another food. It is evident that, in the first case, the food possessed a higher nutritive value than in the second; and, consequently, that it might have been reduced without lessening the production of milk. What has been said will be sufficient to show how difficult is the task of carrying out trustworthy experiments for the purpose of determining the influence of different foods on the production of milk or butter.

In addition to the inquiries respecting the comparative feeding value of silage made from different crops, it is very desirable that investigations should be carefully carried out in regard to the rela-

of food to the milk produced. In the Rothamsted dairy, as usually in all dairies, some rough and ready plan is adopted for increasing or diminishing the more costly foods according to the amount milk or butter yielded; but we are not aware of any experiments which the rise and fall of both milk and food have been made the subject of careful inquiry.

Our experiments on the feeding of cows with silage were commenced in December, 1884, with red clover-silage, followed later withadow-grass-silage, and they were continued until the cows went to grass in the spring. Next year we propose to repeat some of experiments, with different foods, as it is very desirable that the comparison between a corn crop cut green and put into a silo, and same crop left to ripen, should be made the subject of investigation. Oats will probably be the crop selected for this purpose.

It is contrary to our usual plan, to publish the results of our experiments before they are completed, and we have fully made up our minds as to the lessons to be learnt from them. There were, however, reasons why an exception seemed desirable in the case of our ensilage experiments. We could at any rate supply sufficient information to enable practical farmers to draw their own conclusions from the evidence brought before them. Any conclusions so put forward, however, should be accepted as possibly subject to modification as the investigation proceeds.

Although many careful experiments are carried out on parts of the farm at Rothamsted without regard to cost, on other parts the object is to make a profit just as much as on any other farm. The dairy is an ordinary working dairy, intended to produce milk as cheaply as possible. A contract is generally made to furnish so many gallons of milk per day for a year, and it may be mentioned that for the past year the contract has been 17d., 21d., and 22d. for 17 pints delivered at a railway station in London. Within certain limits, the quantity contracted for must be delivered. If more is produced it must be got rid of some other way; and if there is a deficiency the alternative is to purchase more cows.

In addition to the cows, there are stock of all ages, bred on the farm, and about 40 Irish oxen are generally purchased in the autumn to consume the rough grass left by the cows. The land is mixed arable and pasture, about equally divided. Much of the pasture is young, and that which is old is not good. Sheep's fescue and *Agrostis alba*, while white clover and the better grasses can only be obtained by high animal manuring. For six months of the year the stock are in the fields, while during the other six months other food must be provided for them. A certain amount of succulent as well as dry food is therefore required. For the former, mangels grown on the

farm, are generally depended upon; and for the latter, hay and straw, with a certain amount of purchased food.

Under these circumstances, which are as ordinary and commonplace as can well be imagined—and although milk only is sent to London, it would be the same if butter were made—the question to consider is: What benefit will be derived from the use of silage?

Each cow in milk requires about 7 tons of mangels during the 6 months, say, 60 cows \times 7 420 tons
100 head of other stock, each $3\frac{1}{2}$ tons mangels or swedes 350 tons

Total 770 tons

Estimating the dry matter in the roots to average 12 per cent., the 770 tons will give about $92\frac{1}{2}$ tons of dry substance of food.

Ten tons of fresh cut grass, clover, or tares—equal to nearly $2\frac{1}{2}$ tons of hay per acre—may be considered a full yield, and about one-fifth of the fresh weight would be dry matter. There would therefore, be required 462 tons of the fresh green produce, or the produce of about 46 acres, to yield the same amount of dry matter as in the 770 tons of roots.

The question next arises—What crops should be grown for ensilage purposes? Should some of the corn crops be cut in the green state? should less roots be grown? or should clover, or pasture grass, be put into the silo?

With regard to the clover or pasture, there is the difficulty that a certain amount of hay is required for winter consumption, and if, in addition, a quantity of green grass were mown for the silo, less stock must be kept in the summer, which would be an objection. It is true that a rather better price is obtained for milk and butter in the winter; but, on the other hand, the cost of keep and attendance is much greater. Possibly it may be suggested that tares instead of roots should be grown; but the roots generally follow a second, and sometimes a third corn crop. Winter tares are an excellent crop, and of these a certain number of acres are always grown, but they are not to be compared with roots as a cleaning crop. In fact the substitution of a silage crop for roots, would be equivalent to giving up taking a barley crop after wheat.

If 25 tons of mangels were grown per acre, 28 acres would yield about $87\frac{1}{2}$ tons dry substance of food, whilst of ensilaged crops it would require more than 40 acres to yield the same amount. The relative feeding values of these two descriptions of food form, therefore, a very serious matter for consideration. Of course, all the difficulties mentioned could be got over, if it were clearly shown that silage is a more profitable food. But without roots the land cannot

be cleaned, unless by means of summer fallow, which is out of the question; and that land should be fairly clean is, certainly, an essential point in successful arable farming.

Practical farmers will thus see that there are difficulties, which are by no means fanciful, in the way of an extended adoption of ensilage; and that, notwithstanding all that has been said in favour of the process, there is a simplicity, and economy, in feeding off clover on the land, and the summer feeding of grass, which it will not be easy to supersede. Some may even go so far as to say: "I don't care how good silage may be as a food, I cannot see my way to adopt it, for I must grow roots." Others may say, "It ought to be a very good food indeed for me to use it, for it has a most disagreeable smell, which pervades the whole farm." The general view will probably be that some further information, derived from carefully conducted experiments, is very desirable.

2.—THE CONSTRUCTION, AND THE FILLING, OF THE SILOS.

In May (1884), the ground was excavated, and the building of the silos, of which there are two, was commenced. Each measured 15 feet by 13 feet 10 inches; the depth being 22 feet, 17 feet below the surface of the ground, and 5 feet above it. The silos are built of brick and cement, and are, in fact, water-tight tanks, with a corrugated iron roof over them. Each silo is calculated to hold from 100 to 120 tons of compressed silage, and it was intended to fill one with red clover, and the other with meadow-grass and oats; but as the oats advanced rather beyond the point which was considered desirable for the purpose, this part of the experiment was postponed until another year.

The first crop of red clover—which was pure clover without rye grass—was a very fine one. There was no rain during the filling of the silo, although, owing to the delay caused by the weighing and sampling, the operation extended from June 23 to June 30. The operations were as follows: every cart, as it came from the field, after passing over the weigh-bridge, was brought alongside the silo, and the clover was then passed through a chaff-cutter driven by steam; a sample of the chaff being taken from every load. Of the first crop, 98 tons were weighed in, several men and boys being employed within the silo to spread the chaff evenly, and tread it down. The material was first covered with boards fitting tolerably close, and then pressure was applied by means of hard Staffordshire bricks, weighing from 9 lbs. to 10 lbs. each. The pressure per square foot amounted to about 90 lbs.

At the end of August the boards were removed, and about 21 tons of a second crop of clover were weighed, chaffed, put in, and sampled.

After the pressure had been again applied the material did not reach within 4 feet of the top. The contents then measured 3706 cubic feet. The silage of the upper 4 feet weighed $45\frac{1}{2}$ lbs., and that of the lower 4 feet, $59\frac{1}{2}$ lbs. per cubic foot, giving an average weight of 53.6 lbs. per cubic foot.

The second silo was filled in exactly the same way as the first, the meadow-grass with which it was filled being taken from four different fields. It will be evident that any information regarding the cost of filling the silo would be of no value, as accuracy, and not economy, was the main object to be attained; and the operations are so simple that any practical farmer can form his own estimate on the subject. It must, however, be borne in mind, that in carrying fresh cut crops, four or five times as much weight has to be carted as would be the case if the crop were made into hay. Roughly speaking, 4 tons of fresh cut clover will make 1 ton of hay; so in all movements of the crop, either in or out of the silo, calculation for the extra amount of labour must be made accordingly.

3.—QUANTITY AND COMPOSITION OF THE HERBAGE PUT INTO THE SILOS, AND OF THE SILAGE TAKEN OUT.

Silo No. 1.

Table I (p. 11) shows—the dates of filling Silo No. 1, the number of loads, the fresh weights of the clover as put in, the percentages of dry matter, ash, and nitrogen, in it, and the actual quantities of dry matter, ash, and nitrogen, put in.

Table II (p. 12) summarises the quantities of fresh clover, and of its various constituents, put into the silo; and it also shows—the quantities of silage, and of its various constituents, taken out; the actual loss of the various constituents; the percentage loss of each constituent, reckoning the total of each as 100; and the percentage loss of each constituent, reckoned on the total Fresh = 100.

We have thought it desirable that these results should be given as fully as possible, in order that those who will take the trouble to examine them, may form their own conclusions respecting them. This is especially desirable, as upon one very important point, namely the loss of constituents which occurs in the silo, we, and probably most of those who are interested in what may be called the chemistry of ensilage, have formed our opinions mainly on results which have been published by German or French chemists.

It would appear to be a very simple thing to estimate the loss which takes place in a silo. It may be said—"You have only carefully to weigh everything that goes in, and everything that comes out, and it is done." It will be seen, however, that a good deal more

FIRST AND SECOND CROP RED CLOVER.

TABLE I.—Quantity and Composition of the Herbage put into Silo No. 1.

Field.	When put in.	No. of Loads.	Fresh Weight.	Percentage composition.			Total weight.		
				Dry matter.	Ash in dry matter.	Nitrogen in dry matter.	Dry matter.	Ash.	Nitrogen.
First Crop Clover.									
by acres.....	1884.		lbs.	Per cent.	Per cent.	Per cent.	lbs.	lbs.	lbs.
	June 23	20	34,776	16·35	10·80	2·942	5,686	614	167·3
	" 23	20	39,035	16·66	11·80	2·866	6,503	767	186·4
	" 23	5	8,827	19·99	9·68	2·781	1,765	171	49·1
	" 24	15	24,462	18·40	10·72	2·920	4,501	483	131·4
	" 24	14	25,554	19·26	10·92	2·793	4,941	540	138·0
mow	" 28	21	34,965	23·02	10·28	2·578	8,049	827	207·5
	June 28	9	12,628	24·07	9·64	2·423	3,040	293	73·7
	28-30	15	23,670	21·32	8·98	2·382	5,046	451	120·3
	" 30	9	13,677	22·83	8·87	2·385	3,122	277	74·5
	Total ...	128	217,694	42,653	4,423	1148·1
Second Crop Clover.									
mow	Aug. 25	4	6,950	19·13	8·61	2·616	1,330	115	34·8
	" 25	2	3,590	22·66	8·92	2·704	813	73	22·0
	" 26	8	12,831	17·10	10·45	2·688	2,194	229	69·0
	" 26	3	5,088	17·92	10·15	2·668	912	93	24·3
	" 26	4	5,483	18·66	8·45	2·602	1,018	86	25·5
	" 26	3	4,786	18·72	9·22	2·623	896	83	23·5
shed off	" 26	6	9,032	21·90	9·15	2·524	1,978	181	49·9
	Total	30	47,760	9,141	860	239·0
	...	1	1,136	(19·43)	(9·28)	(2·618)	221	21	5·8
Total	29	46,624	8,920	839	233·2	
Summary.									
1st crop clover.....	...	128	217,694	42,653	4,423	1148·1
2nd crop clover..	...	29	46,624	8,920	839	233·2
Total		157	264,318	51,573	5,262	1381·3
Tons, cwts.			118 0	23 0½	2 7	0 12½

than this is required. So far as regards the total loss we have no difficulty. We weighed in 118 tons, and weighed out $88\frac{1}{2}$ tons, thus showing a loss of $29\frac{1}{2}$ tons. But the results further show that, of this loss, more than 28 tons consisted of water, and only about $1\frac{1}{2}$ ton of dry or solid substance; the total loss of dry substance amounting to about 5 per cent. of that put in. It may be mentioned that this is very nearly the same amount of loss as that which we found had taken place in a large rick of about 40 tons of hay, after standing two years.

Careful experiments have been made in Germany, by Weiske and

TABLE II.—Quantity and Composition—of the Herbage put into the Silo No. 1, the Silage taken out; and actual, and Percentage, Loss of Constituents in the

	Authority for Analyses.	Fresh.	Dry Matter.	Ash.	Nitro- gen.	Dry Organic.	Crude Nitrogenous Substance = N x 6.25.	Crude Non- Nitrogenous Substance.
Herbage as put into the Silo No. 1.								
Second crop clover ...	Rothamsted	lbs. 46,624	lbs. 8,920	lbs. 839	lbs. 233	lbs. 8,081	lbs. 1,456	lbs. 6,625
First crop clover	"	217,694	42,653	4,423	1,148	38,230	7,175	31,055
Total	"	264,318	51,573	5,262	1,381	46,311	8,631	37,680
Silage taken out of the Silo No. 1.								
Second crop clover	Rothamsted Voelcker ...	37,739	8,178 8,069	717 766	211 203	7,461 7,303	1,319 1,259	6,142 6,034
	Mean ...		8,124	742	207	7,382	1,294	6,088
First crop clover ...	Rothamsted Voelcker ...	160,731	40,791 40,881	3,548 3,767	1,062 1,059	37,243 37,114	6,638 6,619	30,805 30,495
	Mean ...		40,836	3,657	1,061	37,179	6,629	30,550
Total	Rothamsted Voelcker ...	198,470	48,969 48,960	4,265 4,833	1,273 1,262	44,704 44,417	7,957 7,888	36,747 36,429
	Mean ...		48,960	4,399	1,268	44,561	7,923	36,638
Actual Loss of Constituents.								
Second crop clover	Rothamsted Voelcker ...	8,885	742 851	122 73	22 30	620 778	137 187	483 591
	Mean ...		796	97	26	699	162	537
First crop clover ...	Rothamsted Voelcker ...	56,963	1,862 1,772	875 656	86 89	987 1,116	537 556	450 569
	Mean ...		1,817	766	87	1,051	546	505
Total	Rothamsted Voelcker ...	65,848	2,604 2,623	997 729	108 119	1,607 1,894	674 743	933 1,151
	Mean ...		2,613	863	113	1,750	708	1,042
Per Cent. Loss—Total of each Constituent=100.								
Second crop clover	Rothamsted Voelcker ...	19.1	8.3 9.5	14.5 8.7	9.4 12.9	7.7 9.6	9.4 12.8	7.3 8.9
	Mean ...		8.9	11.6	11.2	8.7	11.1	8.1
First crop clover ...	Rothamsted Voelcker ...	26.2	4.3 4.2	19.8 14.8	7.5 7.8	2.6 2.9	7.5 7.7	1.4 1.8
	Mean ...		4.3	17.3	7.6	2.8	7.6	1.6
Total	Rothamsted Voelcker ...	24.9	5.1 5.1	18.9 13.9	7.8 8.6	3.5 4.1	7.8 8.6	2.5 3.1
	Mean ...		5.1	16.4	8.2	3.8	8.2	2.8
Per Cent. Loss—Total Fresh=100.								
Second crop clover	Rothamsted Voelcker ...	19.1	1.6 1.8	0.3 0.1	0.05 0.06	1.3 1.7	0.3 0.4	1.0 1.3
	Mean ...		1.7	0.2	0.06	1.5	0.4	1.1
First crop clover ...	Rothamsted Voelcker ...	26.2	0.9 0.8	0.4 0.3	0.04 0.04	0.5 0.5	0.3 0.3	0.2 0.3
	Mean ...		0.9	0.4	0.04	0.5	0.3	0.2
Total	Rothamsted Voelcker ...	24.9	1.0 1.0	0.4 0.3	0.04 0.04	0.6 0.7	0.3 0.3	0.3 0.4
	Mean ...		1.0	0.3	0.04	0.7	0.3	0.4

Schulze, on the changes and losses in ensilaging; and as their results are familiar to all those who have studied the chemistry of ensilage, we propose merely to refer to them so far as to point out that with lupines, maize, and lucerne, the loss during fermentation amounted to between 22 and 36 per cent. of the dry matter. In some of the experiments one-third or more of some of the food-constituents contained in the green crops put in, was destroyed, while another portion was reduced in value. The experiments were made in tight casks, and in some cases the filling in and the putting on of the pressure were done at once; whilst in others the filling was extended to six or seven days. The latter was about the time occupied in the process in our own experiments.

In the German experiments the loss of water was very small, but there was a considerable loss of mineral matter. It has been generally held that, as in the fermentation of vegetable substances, which contain both organic and mineral matter, the destruction of the former necessarily increases the percentage of the latter, the difference between the amount of mineral matter in the fresh crop put into the silo and that in the silage taken out, would, in some degree, be a measure of the loss of organic matter during fermentation, and also be some check on the accuracy of the work. Weiske and Schulze point out, however; that owing to the sand, and other adventitious mineral matters, with which bulks of herbage from the field are always contaminated, the difference in the amounts cannot be relied upon as a basis of calculation. Our own results fully confirm this view. Indeed, they show a loss, and not a gain, of mineral matter in the silage, as compared with that in the clover put in. It is true that our loss of dry substance would only amount to 1 per cent. of the fresh herbage, and the loss of mineral matter to about one-third of 1 per cent.; but small as this loss appears to be, according to one set of analyses it amounts to 729 lbs., and to the other to 1000 lbs. in the total quantity of mineral substances contained in the 118 tons of clover that were put into the silo.

The complete explanation of this loss is not very obvious. It may be pointed out, however, that of the 16 samples of clover which were taken as the crop was chaffed and put into the silo, the variations in the percentage of ash are very great; the lowest being a little below $8\frac{1}{2}$, and the highest nearly 12 per cent. of the dry substance.

The depth of the silage in the pit was between 17 and 18 feet; and its removal for feeding purposes was a daily operation extending over three months. It was taken out in four layers, each of 4 to 5 feet in depth. From each layer five, six, or more samples were taken for analysis. A quantity of 1000 lbs., or more, was weighed, and rapidly mixed on an asphalt floor, so as to avoid loss by evaporation as much

as possible; and from this samples were taken. To ascertain the range of loss by evaporation in such an operation several experiments were made, of which the following is an example:—

		lbs.
January 2.	Weight 11.30 A.M.	200
	" 3 P.M.	199 $\frac{1}{4}$
	After turning over and mixing	199 $\frac{1}{4}$
January 3.	Weight 10 A.M.	198
	Loss.....	2

On a very windy day the loss amounted to 2 $\frac{1}{4}$ per cent., and on a still, damp day, it was less than $\frac{1}{4}$ per cent. On the 25th April, the weighing machine was taken into the silo, and a block of silage, about 3 feet by 2 feet by 3 feet deep, was cut out and immediately weighed in baskets, when the weight was found to be 814 lbs. It was then taken to the mixing floor, where it was turned over three times, and samples were taken from it. At the end of an hour the silage was again weighed, and the loss was found to be not quite 1 $\frac{1}{4}$ per cent. It was noticed that the floor, although previously quite dry, had become wet. It may also be mentioned that the weather was much warmer when this experiment was carried out, and that the silage was taken from the second silo, the first having been emptied some weeks previously.

It should be added that the silos, which, as already explained, were built of brick and cement, were filled very shortly after they were finished. This may to a certain extent have been a source of loss, as a good deal of the floor rested upon the chalk, which would carry away water very rapidly if there were any crack in the cement. It is quite possible, therefore, that some loss by drainage did take place. The late Dr. Voelcker gave the composition of the drainage from a silo as 90 per cent. of water, and 10 per cent. solid matter, of which more than one-third was mineral.

It seems somewhat difficult to suppose that the very large loss of water in the silo was wholly due to evaporation, when the loss of organic matter, by fermentation, was so small. But it is to be observed that the lower layers were removed more slowly than the others, so that the cut surfaces were longer exposed; whilst, as a matter of fact, the silage was drier towards the bottom. On the other hand, some drainage was observed soon after the filling and the application of the pressure, so that some at any rate of the loss was due to drainage. If we were to assume that drainage had taken place sufficient to carry off the whole of the apparent loss of minerals—amounting to 863 lbs.—in a solution of the strength of the liquid which was pressed out of a silo, and was analysed by the late Professor Voelcker, the loss of water by drainage would be 8 per cent., and by

evaporation 17 per cent.; but the actual loss by drainage was probably less than here supposed.

It is satisfactory to know that the Council of the Bath and West of England Society have voted a sum of money for the purpose of investigating the loss of nutritive substances which takes place during the fermentation of green crops in a silo; and it is to be hoped that the experiments will be carried out with the necessary care. It is evident that, until this important matter has been thoroughly sifted, all estimates in regard to the economy of ensilaging, as compared with the ordinary operations of the farm, are little better than guess-work.

Silo No. 2.

Silo No. 2 was similar in all respects to Silo No. 1, as already described; and was only divided from it by a 9-inch wall. From June 25 to July 2 (1884), about $45\frac{1}{2}$ tons of first-crop meadow-grass, from four different fields, were first put in. The whole of the grass was of fairly good quality; but that from the second field was much the ripest. It was intended to follow with winter oats; but the hot weather brought the crop too forward to be suitable for silage; so the full pressure was applied.

The silo was opened on August 23, when, from that date to August 26, about $30\frac{1}{2}$ tons of second-crop red clover were put in.

In the middle of October the weights were again removed, and about $6\frac{1}{2}$ tons of second-crop meadow-grass were put in. Part of this grass was from land which had already been mown once; and the rest was from a field which had been fed, and consisted chiefly of the portions which, having grown too strong, had been left by the stock.

As in the case of Silo No. 1, all the crops were chaffed, and the material was evenly spread and well trodden down, as it was put in.

Table III (p. 16) gives the fresh weights of each lot of herbage put into the silo; also the percentages, and the actual quantities, of dry matter, mineral matter, and nitrogen, which each contained.

Table IV (p. 17) summarises the amounts of fresh herbage, and its various constituents, put into the silo, and gives the amounts of silage, and of its various constituents taken out. It also shows the actual loss of each of the constituents in the silo, the loss per 100 of each, and the loss of each per 100 of fresh herbage put in.

Comparison of the results given in Table III, with those formerly given in Table I, relating to the herbage put into Silo No. 1, will show that whilst the dry matter in the first-crop clover put into Silo No. 1 ranged from under 17 to 24 per cent., and that of the second-crop clover from over 17 to under 23 per cent., the dry matter of the first-crop meadow-grass put into Silo No. 2 ranged from $24\frac{1}{2}$ to nearly 32 per cent., that of the second-crop grass was more than 38 per cent.,

FIRST AND SECOND CROP MEADOW-GRASS, AND SECOND CROP CLOVER.

TABLE III.—Quantity and Composition of the Herbage put into Silo No. 2

Field.	When put in.	No. of Loads.	Fresh Weight.	Percentage composition.			Total weight.		
				Dry matter.	Ash in dry matter.	Nitrogen in dry matter.	Dry matter.	Ash.	Silo ga.
First Crop Meadow-Grass.									
	1864.		lbs.	Per cent.	Per cent.	Per cent.	lbs.	lbs.	cu. ft.
Broadbalk and Appletree	June 25-26	29	46,502	24.72	8.47	1.685	10,086	851	104
Dr. Gilbert's meadow	" 25-26	12	19,609	31.56	7.67	1.340	6,247	479	87
Park and Grass-wick	" 26	11	13,639	27.30	7.44	1.628	3,728	277	47
Park and Lodge Meadow	July 2	20	27,990	26.66	8.14	1.670	7,446	608	81
Total		72	101,980	27,502	2,216	401
Second Crop Clover.									
Thirty acres	August 23	35	32,405	39.27	8.14	2.620	12,727	1,036	221
"	" 23	10	8,363	24.36	9.65	2.435	2,037	184	69
"	" 25	3	3,880	19.63	11.01	2.907	762	84	21
"	" 25	1	2,631	19.41	10.02	2.775	394	39	17
Geencroft	" 25	2	2,090	19.88	9.05	2.649	415	38	19
"	" 25	2	2,160	19.01	9.13	2.269	411	38	19
"	" 25	2	2,008	19.50	10.90	2.557	391	42	19
"	" 25	1	1,040	18.36	9.40	2.798	191	18	10
"	" 26	5	5,790	21.90	9.15	2.524	1,268	116	23
Thirty acres	" 26	7	7,218	32.67	7.97	2.639	2,358	188	47
Total		68	66,985	20,954	1,784	364
Brought from Silo 1		1	1,136	(19.43)	(9.28)	(2.618)	221	21	11
Total		69	68,121	21,175	1,805	375
Second Crop Meadow-Grass.									
Park and Park Field	October 17	17	14,858	38.13	9.04	1.890	5,665	512	107
Summary.									
First crop grass		72	101,980	27,502	2,216	401
Second crop clover		69	68,121	21,175	1,805	375
Second crop grass		17	14,858	5,665	512	107
Total		158	184,959	54,342	4,533	1,073
Tons, cwt...		82 1 1/2	24 5 1/2	2 0 1/2	8 1/2

and that of the second-crop clover ranged from over 18 to over 39 per cent. Indeed, more than two-thirds of the clover put into the Silo No. 2 averaged nearly 36 per cent. dry matter. It is thus seen that the herbage put into Silo No. 2 contained a very much less proportion of water than that put into Silo No. 1. It was, in fact, upon the whole, riper, and doubtless more woody, especially so far as the meadow-grass was concerned.

TABLE IV.—Quantity and Composition of the Herbage put into Silo No. 2; of the Silage taken out; and Actual and Percentage Loss of Constituents in the Silo.

	Fresh.	Dry Matter.	Ash.	Nitro- gen.	Dry Organic.	Crude Nitrogenous Substance = N × 6.25.	Crude Non- Nitrogenous Substance.	Water.
<i>Herbage as put into the Silo.</i>								
seed crop grass.....	lbs. 14,858	lbs. 5,665	lbs. 612	lbs. 107	lbs. 5,153	lbs. 669	lbs. 4,484	lbs. 9,193
seed crop clover	68,121	21,175	1,805	552	19,370	3,450	15,920	46,946
wt crop grass.....	101,980	27,502	2,216	438	25,286	2,738	22,548	74,478
Total	184,959	54,342	4,533	1,097	49,809	6,857	42,952	130,617
<i>Silage taken out of the Silo.</i>								
seed crop grass.....	14,473	5,068	525	91	4,543	568	3,975	9,405
seed crop clover	65,670	20,651	1,809	551	18,842	3,444	15,398	46,019
wt crop grass.....	90,798	28,861	2,086	369	21,275	2,306	18,969	67,437
Total	170,941	49,080	4,420	1,011	44,660	6,318	38,342	121,861
<i>Actual Loss of Constituents.</i>								
seed crop grass.....	3.5	597	+ 13	16	610	101	509	+ 212
seed crop clover	2,451	524	+ 4	1	528	6	522	1,927
wt crop grass.....	11,182	4,141	130	69	4,011	432	3,579	7,041
Total	14,018	5,262	113	86	5,149	539	4,610	8,766
<i>Loss per Cent.—Total of each Constituent = 100.</i>								
seed crop grass.....	Per cent. 2.6	Per cent. 10.6	Per cent. +2.6	Per cent. 15.1	Per cent. 11.8	Per cent. 15.1	Per cent. 11.4	Per cent. +2.3
seed crop clover	3.6	2.5	+0.2	0.2	2.7	0.2	3.3	4.1
wt crop grass.....	11.0	15.1	5.9	15.8	15.9	15.8	15.9	9.6
Total	7.6	9.7	2.6	7.8	10.3	7.8	10.7	6.7
<i>Loss per Cent.—Total Fresh = 100.</i>								
seed crop grass.....	2.6	4.0	+0.09	0.11	4.1	0.7	3.4	+1.4
seed crop clover	3.6	0.8	+0.006	0.001	0.8	0.009	0.8	2.8
wt crop grass.....	11.0	4.1	0.1	0.07	3.9	0.4	3.5	7.0
Total	7.6	2.8	0.06	0.04	2.8	0.3	2.5	4.7

The great difference in the composition of the various crops put into the silos renders it extremely difficult to determine with certainty the changes and the losses which take place in them. There is also considerable difficulty arising from the depth of the silage in the pits, and the necessity of taking it out in several separate layers, and occupying weeks, or it may be months, in the process. The experience gained

this first year will probably enable us the better to guard against some of these possible sources of error in the future.

The columns of Table III (p. 16), giving the percentages of ash and of nitrogen in the dry matter of the herbage put into Silo No. 2, show that, whilst in the first crop of grass the ash varied from $7\frac{1}{2}$ to $8\frac{1}{2}$ per cent., and in the second crop of grass it was about 9 per cent., in the second crop of clover it varied from about 8 to about 11 per cent. of the dry matter. There is a still greater difference in the percentages of nitrogen in the dry matter of the two descriptions of herbage. Thus, whilst in the dry matter of the meadow-grass, the nitrogen ranged from 1.34 to 1.89 per cent., in that of the clover it ranged from 2.27 to 2.91 per cent.

Turning now to Table IV (p. 17) it is seen that the proportion of water lost in the silo was comparatively small, both in the meadow-grass and in the clover. The loss was in fact much less than in Silo No. 1. The crops put into the Silo No. 2 contained, it is true, a much less proportion of water than those put into Silo No. 1. But the amount of loss by fermentation was much greater, and hence it would be supposed that the greater heat developed would have caused more evaporation. It would seem, therefore, as was supposed, that part of the greater loss of water in Silo No. 1 was due to drainage, and part to the length of time taken in using the silage, and the consequent long exposure of the cut surfaces. In the second crop of clover in Silo No. 2 the loss of water indicated is only 4.1 per cent., whilst in the second crop of clover in the Silo No. 1 it amounted to over 20 per cent. Further, it will be observed that there was an actual gain of water by the second crop of grass, which formed the top layer in Silo No. 2, it having doubtless absorbed watery vapour from the layers below it.

The table (IV) shows that the losses of dry substance, and of its several organic constituents, were much greater, both in the first crop of grass which was at the bottom of the silo, and in the second crop of grass which was at the top, than in the clover between them. Of the 21,175 lbs. dry substance in the clover put into the silo, there was a loss of only 524 lbs., or of only 2.5 per cent.; whilst in the first crop grass below it there was a loss of 15.1 per cent., and in the second crop grass above it, a loss of 10.5 per cent. of the dry matter put in.

Of mineral matter, or ash, the figures show a gain in the second crop grass, and in the second crop clover, but a loss in the first crop grass. An examination of the ashes of the herbage put in, and of the silage taken out, clearly shows that, as before assumed, the irregularity is dependent on the variable amounts of adventitious mineral matter in the ashes.

Of the total nitrogen, the loss in the first crop of grass below the

clover was 15·8 per cent., and in the second crop grass above the clover it was nearly the same, namely, 15·1 per cent.; whilst in the clover itself, between them, the percentage of nitrogen in which is so much higher, the loss of it was extremely small, amounting to less than a quarter of 1 per cent.

Thus, contrary to what might have been expected, the results indicate much more loss of dry matter, and of its several constituents, in the meadow-grass than in the clover. Further reference to the various changes and losses will be made when we come to consider the chemical composition of the herbage put in, and of the silage taken out of the silos. It is obvious that, so far as the results of these first experiments are to be relied upon, the losses of the food-constituents of the crops are far less than in the German experiments before referred to.

4.—THE CHEMICAL COMPOSITION OF THE SILAGE.

Although a very large amount of analytical work has been executed in connection with our ensilage experiments, the data at present at command are insufficient to enable us to discuss at all adequately the changes, and the losses, which have taken place in the herbage put into the silo. Still, the results do bring to view some points of interest respecting which there can be little doubt.

Confining attention on the present occasion to Silo No. 1, it may be stated that, of the herbage put in, nine different mixed samples were made of the first-crop red-clover, and seven of the second-crop red-clover, as chaffed and put into the silo. Each of these samples represents the number of loads, and the weight of herbage, as shown in Table I, page 11; and is a mixture of a given quantity taken from each individual load. Unfortunately, the pressure of other work in the Rothamsted Laboratory rendered it quite impossible to determine the composition of these mixed samples in the green, undried condition. Each was, however, carefully sun-and-air dried; and in each, the dry matter, the ash, and the total nitrogen, were determined at Rothamsted. In proportionally mixed samples, representing, respectively, the whole of the first crop, and the whole of the second crop, the amounts of total nitrogen, of albuminoid nitrogen, and of woody fibre, have also been determined. The analyses of these mixed samples are, however, incomplete, but we are able to give, further on, a general indication of their bearing.

The first crop of clover put into the silo formed a layer of silage of 13 feet 10 inches in depth; and the second crop clover above it a layer of 4 feet of silage.

This latter, the second crop clover-silage, was taken out in one layer

to its full depth; and six samples were taken from it, namely—on December 13, 15, 18, and 30, 1884, and January 2 and 7, 1885. In each of these, the dry matter, the ash, and the total nitrogen, were determined at Rothamsted; and of three of them, namely, those taken on December 13, 18, and 30, more complete analyses were made by Dr. John A. Voelcker.

The first crop clover-silage was taken out in four layers, respectively of 4 feet 2 inches, 4 feet, 5 feet, and 8 inches, in depth. From the first layer five samples, namely, on January 14, 17, 20, 23, and 28, were taken; from the second layer five samples were taken, namely, on February 2, 6, 10, 17, and 19; from the third layer six samples were taken, namely, on March 6, 12, 14, 18, and 26, and April 4; and from the fourth layer five samples, namely, on March 10, 13, 17, and 20, and April 2. In each of these first crop clover-silage samples, the dry matter, the ash, and the total nitrogen, were, as before, determined at Rothamsted; and more complete analyses were made by Dr. John A. Voelcker, of the samples of January 17 and 23, of the first layer; of February 6 and 17 of the second layer; and of March 12 of the third layer. It was intended that another complete analysis of the silage of the third layer, and at least one of that of the fourth layer, should have been made, but this has not been accomplished.

Tables V and VI (pp. 21 and 22) give the results of the three complete analyses of the second-crop clover-silage, and of the five of the first-crop clover-silage; and, in the lower part of each table, is given a summary showing, side by side with Dr. Voelcker's determinations of dry matter, ash, and nitrogen, those made at Rothamsted in the same samples. Table V shows the percentage composition of the silage in the moist condition in which the samples were taken; and Table VI shows the percentage composition of the dry substance of the silage.

It will be seen that, upon the whole, Dr. Voelcker's and the Rothamsted determinations of dry matter, and of nitrogen, in the silage, agree very closely. Dr. Voelcker's determinations of mineral matter, or ash, are, however, always higher than those made at Rothamsted; the difference being probably due to the fact that, in the Rothamsted ashes, all visible adventitious matter was carefully picked out, whilst this was not done in the case of Dr. Voelcker's ashes.

It is frequently assumed that, although there is more or less loss of certain food-constituents in the conversion of succulent herbage into silage, yet this is more than compensated by the conversion of indigestible into digestible matter. The results at present at our command certainly do not seem to bear out this view.

ANALYSES OF THE SILAGE FROM SILO NO. 1; FIRST AND SECOND CROP RED CLOVER;
COMPLETE ANALYSES BY DR. J. A. VOELCKER;
PARTIAL ANALYSES MADE AT ROTHAMSTED.

TABLE V.—*Per Cent. in Fresh Silage. Detailed Analyses of each Layer.*

Constituents.	2nd Crop Red Clover.				1st Crop Red Clover.					
	Depth of Silage 4 feet.				1st depth, 4 feet 2 inches.		2nd depth, 4 feet.		3rd depth, 5 feet.	Mean.
	Sample 1 taken Dec. 13, 1884.	Sample 2 taken Dec. 18, 1884.	Sample 3 taken Dec. 30, 1884.	Mean.	Sample 1 taken Jan. 17, 1885.	Sample 2 taken Jan. 23, 1885.	Sample 1 taken Feb. 6, 1885.	Sample 2 taken Feb. 17, 1885.	Sample 1 taken Mar. 12, 1885.	
	P. cent.	P. cent.	P. cent.	P. cent.	P. cent.	P. cent.	P. cent.	P. cent.	P. cent.	
Water	78.39	79.17	78.30	78.62	76.24	76.49	73.84	73.88	73.82	74.85
Water-soluble Albuminoids	0.12	0.44	0.49	0.35	0.31	0.50	0.88	0.75	0.56	0.60
Water-soluble Albuminoids ..	1.94	1.75	1.69	1.79	1.81	1.81	1.93	1.75	2.19	1.90
Total	2.06	2.19	2.18	2.14	2.12	2.31	2.81	2.50	2.75	2.50
Insoluble Ash	1.28	1.37	1.28	1.31	1.45	1.40	1.88	1.69	1.58	1.60
Water-soluble Ash	0.76	0.64	0.75	0.72	0.67	0.69	0.71	0.88	0.78	0.75
Total	2.04	2.01	2.03	2.03	2.12	2.09	2.59	2.57	2.36	2.35
Water-soluble Fibre	5.37	5.29	5.51	5.39	5.93	6.13	6.56	7.06	6.45	6.42
Water-soluble Fibre	7.02	6.41	6.15	6.53	6.72	6.70	6.48	6.50	6.67	6.61
Total	12.39	11.70	11.66	11.92	12.65	12.83	13.04	13.56	13.12	13.03
Free acid	0.44	0.19	0.62	0.42	0.56	0.76	0.63	0.70	0.73	0.68
Free acid	0.86	1.18	0.85	0.96	1.10	0.84	1.11	0.79	0.81	0.93
Total	1.30	1.37	1.47	1.38	1.66	1.60	1.74	1.49	1.54	1.61
Water-soluble Carbohydrates, Limes (1), Chloro- phyl, &c.	3.82	3.56	4.36	3.91	5.21	4.68	5.23	6.00	6.41	5.66
Total of all	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Non-albuminoid Nitrogen	0.33	0.35	0.35	0.35	0.34	0.37	0.45	0.40	0.44	0.40
Non-albuminoid Nitro- gen	0.21	0.17	0.20	0.19	0.17	0.22	0.23	0.26	0.28	0.23
Total	0.54	0.52	0.55	0.54	0.51	0.59	0.68	0.66	0.72	0.63

Summary.

1st crop Red Clover.	Voelcker	21.61	20.83	21.70	21.38	23.76	23.51	26.16	26.12	26.18	25.15
	Rothamsted *	21.38	20.44	21.62	21.15	23.84	22.74	25.00	25.29	25.15	24.40
	Rothamsted †	21.82	20.63	22.24	21.66	24.40	23.50	25.63	25.99	25.88	25.08
2nd crop Red Clover.	Voelcker	2.04	2.01	2.03	2.03	2.12	2.09	2.59	2.57	2.36	2.35
	Rothamsted ..	1.92	1.81	1.93	1.89	2.02	1.90	2.36	2.44	2.11	2.17
3rd crop Red Clover.	Voelcker	0.54	0.52	0.55	0.54	0.51	0.59	0.68	0.66	0.72	0.63
	Rothamsted ...	0.56	0.53	0.57	0.55	0.54	0.54	0.65	0.67	0.69	0.62

* Actual result by drying at 100° C.

† Actual result + volatile (acetic) acid.

ANALYSES OF THE SILAGE FROM SILO NO. 1; FIRST AND SECOND CROP RED CLOVER

COMPLETE ANALYSES BY DR. J. A. VOELCKER;
PARTIAL ANALYSES MADE AT ROTHAMSTED.

TABLE VI.—Per Cent. in Dry Matter. Detailed Analyses of each Layer.

	2nd Crop Red Clover.				1st Crop Red Clover.					Mean.
	Depth of Silage 4 feet.				1st depth, 4 feet 2 inches.		2nd depth, 4 feet.		2nd depth, 5 feet.	
	Sample 1 taken Dec. 13, 1884.	Sample 2 taken Dec. 18, 1884.	Sample 3 taken Dec. 30, 1884.	Mean.	Sample 1 taken Jan. 17, 1885.	Sample 2 taken Jan. 23, 1885.	Sample 1 taken Feb. 6, 1885.	Sample 2 taken Feb. 17, 1885.	Sample 3 taken Mar. 12, 1885.	
	P. cent.	P. cent.	P. cent.		P. cent.	P. cent.	P. cent.	P. cent.	P. cent.	
Soluble Albuminoids	0.56	2.11	2.26	1.64	1.30	2.13	3.36	2.87	2.14	2.19
Insoluble Albuminoids	8.97	8.40	7.79	8.39	7.62	7.70	7.38	6.70	8.37	7.90
Total	9.53	10.51	10.05	10.03	8.92	9.83	10.74	9.57	10.51	9.49
Soluble Ash	5.92	6.58	5.90	6.13	6.10	5.96	7.19	6.47	6.03	6.45
Insoluble Ash	3.52	3.07	3.46	3.35	2.82	2.93	2.71	3.37	2.98	2.94
Total	9.44	9.65	9.36	9.48	8.92	8.89	9.90	9.84	9.01	9.69
Digestible Fibre	24.85	25.40	25.39	25.21	24.96	26.07	25.08	27.03	24.64	25.44
Woody Fibre	32.48	30.77	28.34	30.53	28.28	28.56	24.77	24.89	25.48	29.09
Total	57.33	56.17	53.73	55.74	53.24	54.57	49.85	51.92	50.12	52.26
Acetic Acid	2.04	0.91	2.86	1.94	2.36	3.23	2.41	2.68	2.79	2.60
Lactic Acid	3.98	5.67	3.91	4.52	4.63	3.57	4.24	3.02	3.09	3.90
Total	6.02	6.58	6.77	6.46	6.99	6.80	6.65	5.70	5.89	6.49
Soluble Carbohydrates, Amides (1), Chloro- phyll, &c.	17.68	17.09	20.09	18.29	21.93	19.91	22.86	22.97	24.48	21.69
Total of all	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Albuminoid Nitrogen ...	1.53	1.68	1.61	1.61	1.43	1.57	1.72	1.53	1.68	1.59
(1) Non-albuminoid Nitro- gen	0.97	0.82	0.92	0.90	0.72	0.94	0.88	1.00	1.07	0.95
Total	2.50	2.50	2.53	2.51	2.15	2.51	2.60	2.53	2.75	2.54

Summary.

Total Dry Matter.	Voelcker	21.61	20.83	21.70	21.38	23.76	23.51	26.16	26.12	26.18	26
	Rothamsted* ..	21.38	20.44	21.62	21.15	23.84	22.74	25.00	25.29	25.15	24
	Rothamsted † ..	21.82	20.63	22.24	21.56	24.40	23.50	25.63	25.99	25.88	25
Total Ash in Dry Matter.	Voelcker	9.44	9.65	9.35	9.48	8.92	8.89	9.90	9.84	9.01	9
	Rothamsted ..	8.80	8.77	8.68	8.75	8.28	8.09	9.21	9.39	8.15	8
Total Nitrogen in Dry Matter.	Voelcker	2.50	2.50	2.54	2.51	2.15	2.51	2.60	2.53	2.75	2
	Rothamsted ..	2.55	2.55	2.57	2.56	2.21	2.30	2.54	2.57	2.65	2

* Actual result by drying at 100° C.

† Actual result + volatile (acetic) acid.

The percentage of "*woody fibre*" is considerably higher in the dry substance of the silage than it was found to be in that of the herbage put in; and, so far as the results enable us to calculate, there was no reduction in the total amount of woody fibre put into the silo. The differences in the percentages of woody fibre in the different samples of silage which the analyses show, might, at first sight, lead to the conclusion that there has been a reduction in its percentage, and in its amount, in the lower layers; and the results might further be supposed to indicate a somewhat corresponding increase in the soluble extractive matters (soluble carbohydrates, &c.). But a careful consideration of the differences in the condition of maturation, and in the composition accordingly, of the herbage contributing to the several layers of silage, shows that the differences in the composition of the latter in these respects, are obviously connected with differences in the character of the herbage put in.

As to the relative proportions of soluble and insoluble albuminoids in the silage, it is obvious that they, too, are largely dependent on the conditions of succulence, or of ripeness, of the herbage put in; though some of the results seem to favour the view that a portion of the insoluble albuminoids has become soluble. According to the results given in Table II (p. 12), there was a loss of about 8 per cent. of the total nitrogen of the herbage put into the silo; and the analytical results now given show that, of the total nitrogen in the silage from the first crop clover, there was an average of 36.8 per cent. in a non-albuminoid condition; whilst initiative results show that, in the herbage put in, less than 20 per cent. of the total nitrogen was non-albuminoid. Again, of the total nitrogen of the second-crop clover-silage, there was an average of 36 per cent. non-albuminoid; whilst in the very ripe herbage put in, only about 10 per cent. of the total nitrogen was non-albuminoid. These results are quite in accordance with those obtained by Professor Kinch, Mr. Clifford Richardson, the late Dr. Voelcker, Mr. W. H. Jordan, and others.

Thus, not only is there a loss of nitrogenous food-material, but a very considerable proportion of the nitrogenous substance which remains is degraded into compounds, some of which are of no value as food (ammonia for example), whilst others, forming a much larger proportion, are, to say the least, of reduced food-value. Further, besides the loss, and the degradation, of nitrogenous substance, it has been shown that there was also more or less loss of non-nitrogenous matter; whilst there is no evidence that woody fibre of a certain degree of induration has been rendered more soluble.

It remains to see what are the results of the experiments on the feeding of animals with the silage.

5.—EXPERIMENTS WITH FATTENING OXEN.

In the beginning of December, 1884, 10 oxen were selected from a herd of 40 Irish shorthorns, of good quality, which cost about 16*l.* a head delivered at Rothamsted a short time previously. On December 19th, they were weighed, and divided into two lots of five each; care being taken that the two lots should correspond as far as possible, both as to the character of the animals, and in weight. The difference in the average weight per head of the two lots, was, at the commencement of the experiment, only 12 lbs. For the purpose of the experiment they were placed in ten boxes under the same roof; one lot of five facing the other lot of five.

The experiment was arranged to compare the feeding value of red-clover-silage with that of an ordinary winter food for fattening stock—namely, a mixture of clover-hay-chaff and swedes. Besides these foods, which were to be tried one against the other, each animal received the same description and amount of purchased or saleable food; namely, 6 lbs. of cake, and 4½ lbs. of barley-meal, per head per day. Of the experimental foods, one lot received an average of rather more than 65 lbs. of clover-silage per head per day; and the other lot 12 lbs. of clover-hay-chaff, and an average of about 50 lbs. of swedes, per head per day. As far as we could calculate beforehand, the quantity of dry substance in the silage given to the one lot, would be nearly the same as that in the clover-hay and swedes given to the other lot.

Table VII (p. 25) shows the exact amounts of the various foods consumed per head per day by each lot. In the particulars of the foods weighed off it will be noticed that some cake was included. This consisted of the harder pieces of decorticated cotton-cake, which has been unusually hard of late. It would have been better to have picked this out, and weighed it separately. It will be seen, however, that the oxen on silage had more than 75 lbs. of the mixed foods per head per day, of which, on the average, only about 1½ lb. was weighed off. Beckoning one-fourth of this to have been cake and three-fourths silage, and that of the food weighed off from Lot 2, one-third was cake, one-third chaff, and one-third swedes, calculation shows that the silage oxen consumed about ½ lb. more dry substance of food per head per day than the others; but this larger amount of dry substance contained a somewhat larger proportion of woody fibre. The total amount of dry substance consumed was, in both cases, on the average, between 24 and 25 lbs. per head per day; and the quantity of nitrogen consumed was practically identical in the two cases.

EXPERIMENTS ON THE FEEDING OF OXEN.

TABLE VII.—FOOD CONSUMED PER HEAD PER DAY.

Lot 1.—5 Oxen; *Experimental Food—Clover Silage.*

<i>Food given per Head per Day.</i>			
Periods.	Clover-silage.	Cake.*	Barley meal.
	lbs.	lbs.	lbs.
December 19—December 31 = 13 days.	65	6	4½
January 1—February 10 = 41 „ .	70	6	4½
February 11—March 16 = 34 „ .	65	6	4½
March 17—April 11 = 26 „ .	65	5	4½
Total..... 114 days.	—	—	—

Food Weighed off.

January 7	20 lbs.	Chiefly silage.
January 26	90 „	Chiefly silage.
February 10	168 „	Silage and cake.
February 20	92 „	Silage and cake.
February 28	124 „	Silage and cake.
March 9	38 „	Chiefly silage.
March 16	116 „	Silage and cake.
April 9	78 „	Chiefly silage.
April 11	57 „	Silage and a little cake.
Total	783 lbs.	Per head per day	1·36 lb.

Lot 2.—5 Oxen; *Experimental Food—Clover-chaff and Swedes.**Food given per Head per Day.*

Periods.	Clover-chaff.	Swedes.	Cake.*	Barley meal.
	lbs.	lbs.	lbs.	lbs.
December 19—December 31 = 13 days.	12	40	6	4½
January 1—January 7 = 7 „ .	12	60	6	4½
January 8—March 16 = 68 „ .	12	50	6	4½
March 17—April 11 = 26 „ .	12	50	5	4½
Total	—	—	—	—

Food Weighed off.

January 7	35 lbs.	Chaff and swedes.
March 16	85 „	Chaff, swedes, and cake.
March 23	46 „	Chaff and swedes.
April 11	34 „	Chaff, swedes, and cake.
Total	200 lbs.	Per head per day	0·35 lb.

* December 19 to February 4 (48 days) an equal mixture of linseed and decorticated cotton-cake; February 5 to April 11 (66 days) decorticated cotton-cake only.

The following Table (VIII) shows the weight of each animal at the commencement, at two intermediate periods, and at the conclusion of the experiment; also the gain in weight within each period, and over the total period of 16 weeks and 2 days. It further shows, for each lot, and for each period, the average increase per head per week, and the average increase per 1000 lbs. live-weight per week.

Taking the results for the whole period, whether we compare the total increase, the average increase per head, the increase per head per week, or per 1000 lbs. live-weight per week, there is a very close agreement between the two lots, the one receiving clover-silage, and the other very nearly the same amount of dry substance in clover-hay-chaff and swedes. The silage has slightly the advantage; but the difference is not more than might be expected in two lots of oxen fed on precisely the same food. Both lots did remarkably well; the silage oxen giving an average increase of rather more, and the others of rather less, than $1\frac{1}{2}$ per cent. of their live-weight per week. There can be

EXPERIMENTS ON THE FEEDING OF OXEN.

TABLE VIII.—*Actual Weights, and Increase in Weight of the Oxen.*

Oxen.	Actual weights.				Increase in weight.			
	Dec. 19, 1884.	Jan. 30, 1885.	Feb. 28, 1885.	April 11, 1885.	Dec. 19 to Jan. 30, 43 days.	Jan. 30 to Feb. 28, 29 days.	Feb. 28 to April 11, 42 days.	Total Dec. 19 to April 11, 114 days.
<i>Lot 1.—5 Oxen; Experimental Food—Clover-silage.</i>								
Nos.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
1.....	1055	1145	1183	1281	90	38	98	226
2.....	1132	1330	1421	1547	198	91	126	415
3.....	1086	1224	1263	1344	138	39	81	258
4.....	1048	1213	1260	1365	166	47	105	317
5.....	1020	1171	1253	1351	151	82	98	331
Total.....	5341	6083	6380	6888	742	297	508	1547
Average	1068	1217	1276	1378	149	59	102	310
Increase per head per week					24.3	14.2	17.0	19.4
Increase per 1000 lbs. live-weight per week					21.2	11.4	12.8	15.4

Lot 2.—Oxen; Experimental Food—Clover-chaff and Swedes.

1.....	1055	1231	1265	1347	176	34	82	292
2.....	1068	1229	1260	1327	161	31	67	259
3.....	1020	1166	1166	1288	146	20	102	268
4.....	1125	1257	1270	1393	132	13	123	268
5.....	1132	1334	1390	1519	202	56	129	387
Total.....	5400	6217	6371	6874	817	154	503	1474
Average	1080	1243	1274	1375	163	31	101	296
Increase per head per week.....					28.5	7.5	16.8	18.1
Increase per 1000 lbs. live-weight per week					22.8	5.9	12.7	14.4

doubt, therefore, that well made red-clover-silage is a very good food for fattening oxen.

About 25 years ago, from the results then at command, we concluded that fattening oxen, liberally fed on good food, composed of a moderate proportion of cake or corn, some hay or straw-chaff, and roots, would, on the average, consume 12 to 13 lbs. of the dry substance of such mixed food per 100 lbs. live-weight per week, and could give 1 lb. of increase in live-weight for 12 or 13 lbs. dry substance of food so consumed; that is 1 per cent. increase in live-weight per week. In these new experiments the oxen receiving silage consumed rather over, and those receiving clover-chaff and swedes rather under,

10 lbs. of dry substance of food per 100 lbs. live-weight per week; and the former gave 1 lb. increase for a little over 9 lbs., and the latter for about $9\frac{1}{2}$ lbs. dry substance of food consumed. In other words, both lots consumed rather more dry substance of food per 100 lbs. live-weight per week than according to our former estimates; but they gave considerably more increase, both upon a given live-weight within a given time (about $1\frac{1}{2}$ instead of 1 per cent. per week), and for a given amount of dry substance of food consumed.

It will be seen that the amount of cake and meal given was rather large, together amounting to $10\frac{1}{2}$ lbs. per head per day. It was, however, thought desirable that the fattening should be completed, or nearly completed, when the experiment closed. It is a question how far the better result than formerly is due to the large proportion of cake and meal; how far to improvement in the fattening qualities of stock since that time; and how far to the fact that, with earlier maturity, the increase in live-weight represents a considerable proportion of growth as well as fattening increase. It may be added that four of the oxen were sold by auction at Watford, and fetched £. 6s. 3d. per head; whilst the others were sold at intervals during seven weeks, and averaged 24l. 12s. 6d. per head.

So far then as the results of a single experiment can be relied upon, it would seem that, as food for fattening oxen, a given amount of dry substance in red-clover-silage is quite equal to the same amount of dry substance in a mixture of clover-hay-chaff and swedes, given in proportion of 12 parts chaff to 50 parts swedes.

It will be of interest to consider—what would be the difference in cropping of the farm, to produce clover-silage on the one hand, or clover-hay-chaff and swedes, on the other, in the proportions used in these experiments with oxen. Supposing that, to give a herd of 12 65 lbs. of clover-silage per head per day, we had to produce 12 tons of the silage; and for the 12 lbs. of clover-hay and 10 lbs. of swedes, per head per day, we had to produce 12 tons of clover-hay, and 50 tons of swedes—how much land would be

required in the two cases? A fairly good crop of red clover, cut twice, would weigh about 10 tons per acre in the green or fresh state; and according to the results with No. 1 Silo, this would yield only about $7\frac{1}{2}$ tons of clover-silage, so that it would require $8\frac{2}{3}$ acres to produce the 65 tons of silage. The 10 tons of first and second crop green clover would make about $2\frac{1}{2}$ tons of clover-hay; so that it would require about $4\frac{2}{3}$ acres to produce the 12 tons of clover-hay. There would thus remain about 4 acres at disposal for the production of the 50 tons of swedes.

6.—FOOD REQUIRED FOR MERE SUSTENANCE, FOR THE PRODUCTION OF MILK, AND FOR THE PRODUCTION OF FATTENING INCREASE.

Before referring to the plan and the results of the experiments with cows, it may be well, with a view to the better understanding of the subject, to make some remarks on food generally, and especially to call attention to the distinction between the amount of food required for the mere sustenance of the animal, and the amount required for the production of the milk.

In the various accounts which have been published of experiments on silage, nothing is more striking than the extraordinary results which are stated to have been obtained by its use. In several cases, the use of an amount of the material which, according to calculation, would not contain more dry substance of food than would be sufficient to support the life of the cow, has, it has been stated, been followed by a remarkable increase in the production of milk and butter. The authors of these statements have, probably, no doubt of their accuracy; but they depend upon others for the record of the facts; and those employed usually try to make these come out so as to meet the views of their masters, if they take an interest in the new food.

The ordinary dry foods of the farm—hay, straw, and corn—contain about one-sixth of their weight of water, and five-sixths of real *dry substance*. Whenever the foods given to cows have been carefully weighed, and the dry matter determined, it has been found that a cow of ordinary size, and in ordinary milking condition, will consume not less than 25 lbs. of "*dry substance*" of food daily.

For example, 40 years ago, Drs. Thomas and Robert Dundas Thomson carried out some feeding experiments at Glasgow, with cows of the Ayrshire breed, weighing about 1000 lbs. each. They used grass alone, or hay, with various dry foods. The cows were yielding about 2 gallons (about $20\frac{2}{3}$ lbs.) of milk per head per day, and the dry substance of food varied from 25 lbs. to 30 lbs. per head per day. The higher amount, which was that consumed when an additional

ntity of corn or linseed was given, yielded only a slight increase the milk.

About 25 years ago we conducted experiments with cows and on, at Rugby, for the Royal Sewage Commission, trying un- vaged against sewaged grass, and the same with oilcake in dition. The various experiments were made in 1861, 1862, and 63. We will confine attention here to the experiments with cows. e average weight of the animals was between 1000 and 1100 lbs. e average of five experiments with grass alone—some unsewaged d some sewaged—showed that 28·7 lbs. of dry substance of food e consumed per head per day, with an average yield of $26\frac{1}{2}$ lbs. of lk per head per day. And the average of five experiments with ss and oilcake showed a consumption of 28·4 lbs. of dry substance of d per head per day, with a yield of $25\frac{1}{2}$ lbs. of milk per head per day. Again, in the course of experiments which we conducted for the ard of Trade on the relative values of unmalted and malted barley, food for stock, cows, oxen, sheep, and pigs were experimented on; and in the case of the 20 cows, of which the average weight s 1140 lbs., and the average yield of milk about 23 lbs. per head day, the average consumption, over a period of 10 weeks, during winter of 1863-4, amounted to about $29\frac{1}{2}$ lbs. dry substance of d per head per day.

In 1884, we assisted Mr. Edwards, who has a large dairy of ghly-bred shorthorns at St. Albans, in carrying out some experi- nts upon ensilage; and it was found, on calculation, that, in the three periments, the food supplied $28\frac{1}{2}$ lbs., 27 lbs., and 27 lbs., per head day, of real *dry substance*; whilst the three lots were yielding, pectively, an average of $19\frac{3}{4}$ lbs., 17 lbs., and scarcely 17 lbs. of milk, e head per day. It may be mentioned that these were unusually ay cows, one indeed weighed over 1700 lbs., and the 26 cows ve an average of 1413 lbs.

Again, in the spring of 1884, the dry food consumed by the thamsted cows, when yielding 30 lbs. of milk per head per day, s about 26 lbs.

It is usually said that it requires about three acres of grass to support cow for a year. If we estimate an acre of ordinary meadow land, en fed or mown, to yield a produce equal to $1\frac{1}{2}$ tons of hay per e, three acres of such produce would furnish a cow with an average lbs. of *dry substance* of food per day during the year; if the pro- ce were equal to $1\frac{3}{4}$ tons of hay per acre, 3 acres would supply an erage of 26·8 lbs. of *dry substance* per day for a year; or if the yield e equal to 2 tons of hay per acre, it would supply an average of 7 lbs. of *dry substance* of food per day, the year round. The d would of course yield much more than its average quantity

during the summer, and much less during the winter; whilst a cow would consume more than its average amount when yielding milk, and less when dry.

It will be seen that there is a fairly general agreement between the figures brought out in the foregoing cases; certainly quite sufficient, not only to raise doubts as to the accuracy of any results which differ materially from them, but also sufficient to provide practical farmers with some basis upon which to calculate the requirements of their stock during the six winter months.

So far, we have only treated of the amount of dry substance of food consumed by a cow that is yielding milk. It is evident, however, that the different foods must vary considerably in their composition, and consequently in their feeding qualities, as also they vary in their cost; and as, in all dairies, the yield of milk varies considerably, not only according to the milk-yielding capacity of the individual cows, but also owing to the necessary fluctuations in the yield from the time of calving to dryness, an accurate knowledge of the feeding qualities of different descriptions of food ought to lead to economy in their use.

For many years past, important investigations have been carried on in Germany, especially at Weende, by Henneberg and others, relating to the digestibility of the various constituents contained in different foods, and also as to the amounts of the digestible substances which are required to support life.

What is called the "sustenance food" of an animal, is the amount which will supply the waste of the body in a state of rest, without either gain or loss of weight. For example, it was found that a ration of $19\frac{1}{2}$ lbs. of clover hay supplied to an ox weighing 1000 lbs., was sufficient to keep up this weight, without adding to it, the animal doing no work, and the temperature of the stall being kept at about 51° Fahrenheit. Of this $19\frac{1}{2}$ lbs. of hay, $\frac{5}{8}$ lb. of nitrogenous, and nearly $7\frac{3}{4}$ lbs. of non-nitrogenous substances, were digested. Other rations were also experimented upon—such as clover hay, straw, and rape cake, in different proportions, and clover hay, mangels, straw, and rape cake. The mean of five experiments, including that with the clover hay alone, showed 0.57 lb. of digestible nitrogenous, and 7.4 lbs. of digestible non-nitrogenous substances, to be required for the support of the animal without gaining or losing weight; but in two of the cases the temperature of the stall was about 62° F., and in the other two it was about 69° F. From results of this kind, both Professor Julius Kühn and Professor Emil v. Wolff have constructed tables which give the amounts of digestible substances that may be expected to be present in a great variety of foods.

Owing to the great difference in the feeding qualities of the same description of food—whether it be roots, hay, corn, or purchased food—it is evident that these tables require to be used with caution; still, when so used, they are of considerable value. We now propose, therefore, to apply them to the results obtained at Rothamsted in the spring of 1884, from 30 cows, each of which consumed daily 4 lbs. decorticated cotton-cake, $3\frac{1}{2}$ lbs. bran, 3·6 lbs. hay chaff, 7·2 lbs. oat-straw chaff, and 81 lbs. mangels, and yielded 30 lbs. of milk per head per day.

The following Table (IX) shows that the food supplied an average of about 25 $\frac{1}{2}$ lbs. dry substance per head per day. It also gives the quantities of digestible nitrogenous and non-nitrogenous substances in the foods, calculated according to our own estimates of average composition, and to Emil v. Wolff's estimates of the proportion of the several constituents which is digestible. It further shows—the amounts which, according to the German estimates, would be required for the sustenance of one of the cows, the average weight of which was about 1290 lbs.; the amounts required for the production of 30 lbs. of milk; and lastly, the estimated excess in the food.

TABLE IX.—*Amount and Distribution of Food, and Food Constituents, per Head per Day.*

	Total dry substance.	Digestible.		
		Nitrogenous substance.	Non-nitrogenous substance.*	Total, nit. and non-nit.-substance.
	lbs.	lbs.	lbs.	lbs.
4 lbs. cotton-cake.....	3·6	1·38	1·93	3·31
$3\frac{1}{2}$ lbs. bran	3·0	0·42	1·41	1·83
3·6 lbs. hay chaff.....	3·0	0·19	1·52	1·71
7·2 lbs. oat-straw chaff ..	6·0	0·10	2·86	2·96
81 lbs. mangels.....	10·1	1·30	7·40	8·70
	25·7	3·39	15·12	18·51
Required for sustenance } of 1290 lbs. live-weight }	—	0·74	9·55	10·29
		2·65	5·57	8·22
Required for 30 lbs. of } milk	—	1·10	3·90	5·00
Estimated excess in food..	—	1·55	1·67	3·22

* Reckoned as starch.

Of the 25½ lbs. of dry substance of food consumed, about 18½ lbs. are reckoned as digestible; and of the non-nitrogenous or respiratory and fat-forming portion of the food, about 9½ lbs. out of the 15½ are estimated to be employed in supporting the life of the animal. It is, however, probable that somewhat more than this was actually used for the purpose, as the temperature of our sheds would be lower than in the cases of the German experiments above referred to. Thus, whilst for the mere sustenance of the cow the demand upon the non-nitrogenous constituents of the food is very large, the requirement for the nitrogenous constituents is small.

In the case of all the foods except the roots, only that portion of the nitrogenous and non-nitrogenous constituents which is estimated to be digestible is entered in the table; but as regards the mangels all the constituents are included, as the German calculations do not make any allowance for indigestible matter in their case. It is generally assumed, however, that it is only those compounds which contain nitrogen in the form of albuminoids that are competent to form flesh and the nitrogenous compounds of milk; whilst it is certain that a very large proportion of the nitrogen in roots is not in that form. In connection with ensilage this subject becomes of great importance, as there seems to be no doubt that during the fermentation much of the albuminoid matter is destroyed.

It will be observed that of the 10·1 lbs. of dry substance of the mangels, 1·3 lb. is given as digestible nitrogenous matter; but of this total quantity as little as one-fifth may, and pretty certainly not more than two-fifths would, consist of albuminoids. Out of the 10·1 lbs. of total dry substance, 7·4 lbs., or about 74 per cent., are recorded as digestible non-nitrogenous, or respiratory and fat-forming matter. We thus see what constitutes the great value of the root-crops. It is the fact that they furnish the essential respiratory constituents of food in very large quantities; though, even when grown under very favourable conditions of soil and season, they are not adapted for fattening, unless used in conjunction with other foods. For the purpose of mere sustenance however—as, for instance, in the case of cows which are out of milk, but are kept in warm yards or sheds—roots are a very suitable food used with straw chaff.

It may here be mentioned incidentally, that the loss of lambs born dead, which is often attributed to the manure used for the roots, is more probably due to the want of sufficient albuminoids in the roots. It should be further understood that while the nitrogenous constituents of the food can be used by the animals for respiratory purposes, the non-nitrogenous constituents cannot be used for the production of albuminoids; and as a certain amount of albuminoids is essential, if the food is deficient in them the animal must wastefully consume the

non-nitrogenous substances in excess in order to obtain enough of the albuminoids.

The quantities given in the table are calculated for a cow weighing 1290 lbs.; but if as many sheep as would represent an equivalent weight were substituted for the cow, the amount of sustenance food necessary to keep them without increasing or losing weight, would be considerably higher; partly because more surface would be exposed to the cold; and partly because there is always more or less growth of wool.

Out of the 15½ lbs. of total digestible non-nitrogenous, or respiratory and fat-forming matter in the food, the home produce of the farm furnished rather more than 11½ lbs., and the purchased cake and bran about 3½ lbs.; and as the sustenance of the animal is reckoned to require only about 9½ lbs., there remains a surplus of 2½ lbs. of the home produce, and all the respiratory constituents of the purchased food, available for the production of milk or increase. Instead of the ½ lbs. required by the cow for sustenance, sheep of the same weight would probably require the whole of the digestible non-nitrogenous substance of the farm-produce for sustenance alone; and they would be entirely dependent upon the purchased food for increase in live-weight.

It has been shown that the food requirements of a cow for the purposes of sustenance only, are very small so far as the nitrogenous constituents are concerned, but large for the respiratory, or non-nitrogenous matters. For a cow yielding 30 lbs. (nearly 3 gallons) of milk per day, however, the requirement for the nitrogenous constituents is very much greater. Foods which are rich in nitrogen, such as oil-cakes and leguminous seeds, are therefore specially adapted for the production of milk.

It is somewhat remarkable that the composition of bran, which has special value in the eyes of all dairy-men, bears a very close relation to that of milk in the proportion of the digestible nitrogenous and non-nitrogenous constituents, as will be seen in the following table:—

	Digestible nitrogenous matter.	Digestible non-nitrogenous matter.
3½ lbs. of bran will supply	0·42	1·41
11½ „ of milk will contain. . . .	0·42	1·49

Table IX (p. 31) shows that, in the 4 lbs. of cake and 3½ lbs. of bran, which were daily supplied to the cows, the total digestible constituents would be approximately the same as those required to produce the 30 lbs. of milk; but the nitrogenous would be in excess, and the non-nitrogenous in about corresponding deficiency. As, however, the

amount of milk varies very much, owing to the difference in the milk-yielding capacity of different cows, and as every cow, whatever its capacity may be, has a maximum yield which is followed by an almost daily decline, it appeared desirable, while the ensilage experiments were going on, that the cake and bran, which obviously contributed so largely to the production of the milk, should be supplied to each cow somewhat in proportion to its yield.

The following Table (X) shows, not only how very large, but how very variable, may be the amounts of total solid matter, nitrogenous matter, fat, &c., in the milk yielded by different cows, or even by the same cow at different periods between calving and dryness. For comparison there is also given, the estimated composition of the increase in live-weight of a fattening ox weighing 1000 lbs., the amount of which increase would frequently be only about 10 lbs., and would rarely exceed 15 lbs. per week.

The table shows the amounts of the various solid constituents in the milk yielded per week by cows giving respectively 4, 8, 12, 16, and 20 quarts per day = 7, 14, 21, 28, and 35 gallons per head per week, assuming the milk to contain 12·50 per cent. of total solids; consisting of 3·65 per cent. albuminoids, 3·50 per cent. butter-fat, 4·60 per cent. sugar, and 0·75 per cent. mineral matter. The table also shows, the estimated constituents in the weekly increase in live-weight of a fattening ox weighing 1000 lbs.:—

TABLE X.

Milk per week (one gallon reckoned to weigh 10·33 lbs.).	Total solid matter.	Nitrogenous substance.	Fat.	Non-nitrogenous substance not fat.	Mineral matter.
	lbs.	lbs.	lbs.	lbs.	lbs.
7 gallons.....	9·04	2·64	2·53	3·33	0·54
14 „	18·08	5·28	5·06	6·66	1·08
21 „	27·12	7·92	7·69	9·99	1·62
28 „	36·16	10·56	10·12	13·32	2·16
35 „	45·20	13·20	12·65	16·65	2·70
10 lbs. increase in live-weight.....	7·25	0·75	6·35	—	0·15
15 lbs. increase in live-weight.....	10·88	1·13	9·63	—	0·22

It may be observed that whilst the meat-producing power of an ox is confined within comparatively narrow limits, the milk-producing capacity of a cow has a very wide range. Another very remarkable fact is the extremely small amount of both nitrogenous and mineral

matter which is stored up in the increase of an ox, compared with that carried off in the milk of a cow. Hence, a dairy, where milk is exported, is very exhausting.

It is evident that there is far more scope for economy in the regulation of the diet of a cow producing milk, than in that of a fattening ox. During the period in which an ox advances from the store condition to fatness, at an average rate of increase which may be estimated at from 1 to $1\frac{1}{2}$ per cent. of its live-weight per week, a cow may be yielding 5 gallons of milk per day at one time, and at another less than one-fourth of that quantity.

Starting with the fact that the two lots of 20 cows each, which were under experiment, were receiving, per head per day, 4 lbs. of cake, and $3\frac{1}{2}$ lbs. bran, with a fixed amount of chaff, and, in addition, in one case silage, and in the other roots, and were yielding an average of 28 lbs. of milk per head per day, it was decided that whilst each lot of 20 cows should continue to receive the total of 80 lbs. of cake daily, the amount should be so apportioned among the 20 that each cow should receive more or less than 4 lbs. daily, accordingly as its yield during the preceding week averaged more, or less, than 28 lbs. of milk per day. The average yield of milk of each cow was, therefore, made up at the end of the week; and, for the succeeding week, 4 lbs. of cake were given to every cow which had yielded 28 lbs. of milk daily: and to each cow which yielded more or less than this quantity, the amount of cake was increased, or reduced, in the proportion of $\frac{1}{4}$ lb. of cake for each 2 lbs. of milk yielded more or less than 28 lbs.

Thus, if a cow yielded 50 lbs. of milk per day (nearly 5 gallons), that is 22 lbs. more than the standard amount of 28 lbs., it received an extra allowance of $2\frac{3}{4}$ lbs. of cake, or in all $6\frac{3}{4}$ lbs. per day; if it yielded 40 lbs. (nearly 4 gallons) or 12 lbs. in excess of the average, it received $1\frac{1}{2}$ lb. cake extra, or a total of $5\frac{1}{2}$ lbs.; if 30 lbs., or 2 lbs. extra, in all $4\frac{1}{4}$ lbs. cake; if only 20 lbs., or 8 lbs. deficiency, only 3 lbs. of cake; or if only 10 lbs., or 18 lbs. deficiency, it received $2\frac{1}{4}$ lbs. less cake, or in all only $1\frac{3}{4}$ lb., and so on.

We cannot at present discuss the results of this experiment; but anyone who looks at the table showing the difference between the amount of solid matter contained in 7 gallons and in 35 gallons of milk, will admit that, without some regulation of the diet, one of two things must take place. Either the yield of some cows must be stinted for want of sufficient food, or others must be receiving food which cannot be turned to profitable account. The highest average yield of milk of any cow during our experiments was 51 lbs., or nearly 20 quarts per day, whilst several were yielding, at the same time, not much more than one-fifth of this quantity.

It is no unusual thing to give fattening oxen 15 lbs. to 18 lbs. of

cake and corn per day. An ox receiving this quantity of food must be consuming a very large amount of nitrogenous substance, of which but a very small percentage is found in the increase in live-weight. A certain amount of that which is digested may be employed in the production of fat, the nitrogen being found in the urine in soluble compounds. In this form, however, the nitrogen nitrifies so rapidly in the soil, that it has little more permanence as manure than that in a salt of ammonia, or in nitrate of soda.

On the other hand, the bulk of the manure of a fattening ox possesses a durability which is well established in practice. Indeed, it may be considered that "condition," or "unexhausted fertility" of a soil, is chiefly due to the constituents of the food which pass, with but little change, in the solid excrements of the fattening animals; but partly also to those in the litter. To *accumulate* nitrogen in the soil, it must be in combination with carbon. In a field where we grow barley continuously, manured with rape cake, although the cake largely increases the crop, a residue still accumulates in the soil, sufficient to be measurable by analysis, and so to establish a claim for unexhausted fertility.

7.—PRELIMINARY EXPERIMENTS WITH MILKING COWS.

In the beginning of February, 1884, one of the laboratory staff was instructed to take the weight of each cow's milk, morning and evening, and to ascertain the amount of food consumed. This was before a brick of the silos was laid, but in anticipation of experiments which it was intended to carry out in the following winter.

From the 4th of February to the 9th March, the average yield of milk per cow per day was 29 lbs. 13 oz., and the average consumption of food per head per day was estimated to be as follows:—

Decorticated cotton cake.....	4 lbs.
Bran	3½ "
Mixed hay and straw chaff	14 "
Mangels	80 "

The fluctuations in the average daily yield of milk were very small, the highest of any one day being just below 32 lbs.; and the lowest a little above 27 lbs. The weights of food were not, however, taken with sufficient accuracy to be altogether reliable, as will be evident from the results obtained in the following month, when it was found that between March 10th and April 7th the average yield of milk was 30 lbs. 5 oz. The same amounts of cake and bran were consumed, but of chaff it was found, on accurate weighing (instead of measuring), that the consumption amounted to only 10½ lbs. per cow,

or $3\frac{1}{2}$ lbs. per head per day less than the amount estimated to be given in the previous month. The amount of dry substance of food consumed per cow daily was now $25\frac{1}{2}$ lbs., whilst the figures for the previous month would give $28\frac{1}{2}$ lbs.; but it is probable that not more than from 25 lbs. to 26 lbs. of dry substance was really consumed.

The grass was so forward on April the 8th, that the cows were turned out from 11 o'clock to 3 o'clock each day. This was continued until the 17th, during which time they had the same cake and bran as before, but only about half the amount of chaff and rather less mangels. On April 17 they were turned out altogether; but continued to have the same quantity of cake; the bran, chaff, and mangels, being gradually reduced; the two latter being stopped altogether in the middle, and the bran at the end of May. The average yield of milk only increased slightly, the amount being $31\frac{1}{2}$ lbs. as against $30\frac{1}{2}$ lbs.; and the highest increase during the summer was to 35 lbs. 3 oz. in the week ending May 17th.

It does not appear that, if cows have been well fed during the winter, any large increase of milk is obtained when they are turned out to grass; and later on, when the weather became hot and dry, the milk fell off very much, the flies having been very troublesome during part of the time. With the exception of the cotton cake, 4 lbs. of which were given to each cow daily, no record of food consumed was possible until the cows again went into their winter quarters in November, but in the meantime the milk was carefully weighed.

8.—PLAN AND ARRANGEMENT OF THE EXPERIMENTS WITH MILKING COWS.

In the beginning of December, 1884, 48 cows were in milk, 40 standing in one house—20 facing the other 20—the food being supplied from a gangway down the centre. We proposed to try only two experiments; and after much trouble in calculation, and finding it hopeless to match each individual cow in one lot with one in the other which had been approximately the same length of time in milk, and was yielding the same amount of milk, it was finally decided to select from the herd two lots, of 20 each, which should agree as nearly as possible in the following conditions: (1) the *average* number of weeks since calving; (2) the *average* yield of milk. This involved the shifting of some of the cows—a proceeding which they strongly resented. Eventually two lots of 20 each were selected, which gave between 14 and 15 weeks as the average date since calving, and an average daily yield of milk of between 30 lbs. and 31 lbs. during the preceding 10 weeks; or less, if they had not been in milk so long.

Of course these averages were made up of very great differences, both in the length of time since calving, and in the yield of milk;

but such is the case in all herds where it is of importance to obtain, as nearly as possible, an uniform supply of milk all the year round. As the experiments were to go on for some months, it was evident that, in both lots, some cows would become dry, and would have to be replaced by others which had recently calved. Still it was considered that results obtained under these circumstances were far more likely to be trustworthy than if the experiment had been made on individual, or on a very few, cows.

Although a certain amount of interest attaches to the substitution of silage for hay, it will be admitted by every practical farmer, that the real value of silage must greatly depend upon its capability of wholly, or at any rate largely, superseding root-crops. To those who are out of the reach of brewer's grains, roots are the only succulent food available for fully half the year; and it is tolerably evident that, without some succulent food, neither meat nor milk can be produced profitably during the winter months.

As already shown, in the experiments with fattening oxen, clover-silage was tried against a mixture of clover-hay-chaff and swedes; but in the experiments with milking cows, now to be described, the comparison is limited to the trial of silaged crops against mangels.

In arranging comparative feeding experiments, it is always desirable that those foods which are not the subject of comparison should be the same for the different lots of animals. Silage, and especially that made from red clover, with which we began our experiment, differs from roots in some important respects. It contains a much higher percentage of dry substance, its dry substance contains a considerably higher proportion of albuminoid compounds, and also very much more indigestible woody fibre. It was impossible, therefore, so to arrange the two rations as to supply exactly the same constituents in each; and it was decided that equal quantities of *dry substance* should be given in the silage, and in the mangels.

The following amounts of other foods were allotted per head per day to each of the 40 cows:—

Cake	4 lbs.
Bran, 3½ lbs., afterwards raised to	4 „
Chaff (half hay and half straw)	10 „

Thus far the same foods were given to each lot; but, in addition, one lot received, per head per day, at first 42 lbs. of clover-silage, gradually raised to 50 lbs., and the other lot at first 75 lbs. of mangels, gradually increased to 90 lbs., as the silage given to the others was increased, and it was found that lower down in the pit it contained a higher percentage of dry matter. Most of the cows took to the silage readily, though a few had a distinct aversion to it, and in order not to

influence their yield of milk injuriously these were supplied with a limited quantity of mangels as well as silage, the former being reduced and the latter increased until their repugnance to the food had ceased.

Two or three weeks elapsed before the disturbance caused by the necessary changes in the position of some of the cows, and in the alteration of their food, was overcome; and it was not until the 24th of December that the animals were weighed.

9.—THE RESULTS OF THE EXPERIMENTS WITH MILKING COWS.

Table XI (p. 41) shows the average amounts of food consumed per head per day, by each lot of 20 cows, within each of the 13 weeks of the experimental period.

Table XII (pp. 42-3) shows, for each of the 20 cows receiving clover-silage, and Table XIII (pp. 44-5), for each of the 20 receiving mangels, the dates of calving, the average yield of milk per day prior to the commencement of the experiment, and the average yield during each of the 13 weeks of the experimental period. They also show, the average yield per day, and the total yield, both in lbs. and gallons, of each cow, over the whole of the experimental period.

Milking cows are exceedingly sensitive to change in anything they have been accustomed to; and after the few necessary transpositions of the animals had been made in accordance with the allotments, and the experiment had fairly commenced, they were treated exactly as if they had not been under experiment at all; the only difference in the management being that both the food and the milk were carefully weighed. Several of the cows of each lot became dry during the experiment; and they were replaced by others, as explained in the foot notes to Tables XII and XIII. The results proved that the new cows were not all equally good milkers; but the differences are to some extent neutralised by the large number of animals in each lot.

Table XI (p. 41), relating to the food, shows that during the first few weeks, until the animals had become accustomed to the silage, a small quantity of mangels was given with it; and this was done again towards the end of the experiment, in the case of a few cows which did not consume all their silage, when it was found that, with the admixture of a small quantity of mangels, they generally consumed their silage also. Most of the animals, however, took to the silage quite readily.

At page 35, it was stated that each lot of 20 cows received the same total amount of cake and bran, but that, after a time, first the cake, and afterwards the bran also, were allotted to the individual cows within each lot in proportion to their yield of milk. That is to

say, instead of each cow receiving 4 lbs. of cake, and $3\frac{1}{2}$ lbs. or 4 lbs. of bran, it received more or less than these quantities, in proportion as its yield of milk was more or less than the average of the whole lot. We shall consider the results of this apportionment of the purchased foods according to the yield of milk at some future time.

Tables XII and XIII show that, over the whole experimental period of 13 weeks, the average yield of milk of the cows receiving clover-silage was 25 lbs, 12 ozs. per head per day, against 27 lbs. 5 ozs., yielded by the cows receiving mangels. This corresponds to a difference over the whole period of 14 gallons per head, or of 281 gallons in the lot of 20 cows, in favour of those receiving mangels.

It is probable that part of this difference was due to the fact that two of the cows brought into the mangel lot during the progress of the experiment (Nos. 21 and 40), turned out to be much better milkers than any two brought into the silage lot. But it is probable that it was also in part due to the more succulent mangels being a more appropriate addition to the dry food for milking cows, than so much silage, which contained nearly twice as high a percentage of dry substance as the mangels. That this was so would appear from the fact that the cows on the silage drank an average of about $1\frac{3}{4}$ gallon more water per head per day than those on the mangels.

Upon the whole the evidence points to the conclusion that if a portion of the dry matter of the clover-silage, say one-fifth or more, had been replaced by a corresponding amount in mangels, not only would some of the cows have consumed the silage better, but such a mixture would doubtless have been more appropriate for milk-yielding. It was feared, however, that if a mixture of the two foods to be compared were given, the results might not be sufficiently distinct. It may be added that the general impression among the attendants was that the cows on the clover-silage showed more tendency to fatten than the others; whilst, as a matter of fact, they did give rather more increase in live-weight. To this point we shall recur further on.

It will be seen that the anticipation we expressed that there would be considerable difficulty in getting strictly comparable results in experiments with dairy cows has been fully realised. But, as the whole of the data are now before the reader, he can form his own conclusions respecting them.

EXPERIMENTS WITH COWS.

TABLE XI.—AVERAGE FOOD CONSUMED PER HEAD PER DAY.

13 WEEKS; DECEMBER 14, 1884, TO MARCH 14, 1885.

Week ending	Average food given per head per day.					Total food weighed off.
	Cake.*	Bran.	Chaff.	Clover-silage.†	Mangels.‡	
Lot 1.—20 Cows ; <i>Experimental Food—Clover-silage.</i>						
1884-5.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Dec. 20 }	4	3½	6½	42½	13½	410½
„ 27 } §	4	3½	9½	44½	8½	0
Jan. 3 }	4	3½	10	46½	4½	0
„ 10.....	4	3½	10	48½	(3) 13½	0
„ 17.....	4	3½	10	50	0	0
„ 24.....	4	3½	10	50	0	0
„ 31.....	4	3½	10	50	0	0
Feb. 7.....	4	3½	10	50	0	0
„ 14.....	4	3½	10	50	0	0
„ 21.....	4	3½	10	50	(1) 5½	0
„ 28.....	4	3¾	10	50	(2) 7½	415
Mar. 7.....	4	4	10	50	(2) 2½	307
„ 14.....	4	4	10	50	0	353
Per head per day....	—	—	—	—	—	0½

Lot 2.—20 Cows; *Experimental Food—Mangels.*

1884-5.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
Dec. 20 } 	4	3½	11	—	75	—
" 27 } 	4	3½	10½	—	75½	—
Jan. 3 }	4	3½	10	—	80	—
" 10	4	3½	10	—	80	—
" 17	4	3½	10	—	82½	—
" 24	4	3½	10	—	90	—
" 31	4	3½	10	—	90	—
Feb. 7	4	3½	10	—	90	—
" 14	4	3½	10	—	90	—
" 21	4	3½	10	—	90	—
" 28	4	3½	10	—	90	—
Mar. 7	4	4	10	—	90	—
" 14	4	4	10	—	90	—
Per head per day....	—	—	—	—	—	—

* December 14 to January 5, half linseed, and half decorticated cotton-cake; afterwards decorticated cotton-cake only.

† Second-crop clover-silage, December 14 to January 10; afterwards first-crop clover-silage.

‡ First three weeks some mangels were supplied to all the silage-fed cows; the figures in parentheses show the number of cows receiving mangels at other periods.

§ Averages for 19 cows only; cow No. 19 various foods.

|| Averages for 19 cows only; cow No. 35 various foods.

EXPERIMENTS

TABLE XII.—YIELD OF MILK

Lot 1.—20 Cows.

Names of cows.	Date of calving.	First milked.	Average milk per head per day.								
			Prior to experiment.						Experimental period		
			From calving.	Previous 10 weeks (or less).		Last week (or less).			Week ending		
									Dec. 20.	Dec. 27.	Jan. 3.
	1884.	1884.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.
1. Strawberry.....	Oct. 28	Oct. 29	47 13	(48 10)	45 11	40 3	39 1	38 13			
2. Phoebe*.....	Mar. 31	Apr. 5	36 7	17 11	12 8	11 11	11 14	11 3			
3. Isabella†.....	Sept. 2	Sept. 3	22 5	20 6	19 2	17 12	16 8	16 13			
4. Penelope.....	Aug. 19	Aug. 21	29 8	25 13	20 7	19 12	20 7	20 1			
5. Beauty‡.....	June 6	June 10	29 11	23 6	18 12	16 6	16 6	16 0			
6. Joan of Arc.....	Sept. 29	Oct. 1	33 5	33 10	29 15	25 7	24 1	24 1			
7. Rothamsted Maid	Sept. 23	Oct. 1	24 3	23 9	16 12	16 0	16 1	16 2			
8. Parody.....	Nov. 2	Nov. 5	39 11	(40 0)	38 7	35 15	34 11	33 0			
9. Queen Bess.....	Aug. 17	Aug. 21	32 0	29 2	21 15	23 2	22 14	22 1			
10. Cherry Ripe.....	Oct. 17	Oct. 21	45 3	(45 14)	41 2	37 14	36 3	35 5			
11. Granny.....	Aug. 18	Aug. 19	26 7	24 13	20 3	20 12	20 12	19 13			
12. Helen.....	Nov. 18	Nov. 19	35 3	(36 12)	35 6	29 7	34 0	31 5			
13. Polly.....	June 19	June 20	36 14	28 0	22 3	20 12	21 6	21 9			
14. Sally.....	Nov. 20	Nov. 21	39 11	(41 15)	41 1	36 2	41 2	44 2			
15. Wonder§.....	July 22	July 23	21 10	19 6	14 15	13 1	14 14	13 14			
16. Fairy Queen 	Feb. 24	Feb. 26	36 0	23 8	20 4	18 5	17 14	18 1			
17. Buttercup.....	Oct. 2	Oct. 4	32 12	32 10	35 0	32 5	33 0	30 0			
18. Narcissus¶.....	June 22	June 24	22 10	16 10	10 8	6 10	{ 2 12 }	23 15			
19. Lucy.....	Dec. 9	Dec. 13	(37 8)	(37 8)	(37 8)	33 3	40 11	42 6			
20. Victoria†.....	Sept. 9	Sept. 10	39 3	37 5	30 0	26 3	26 14	26 15			
Average.....			33 6	30 5	26 9	24 1	25 8	25 12			

* Replaced January 11th, by Ann, calved January 3rd.

† Bought as newly-calved at this date; exact date of calving not known.

‡ Replaced February 5th, by Gypsy, calved January 30th; Gypsy (ill) replaced February 16th, by Rosemary, calved February 12th.

WITH COWS.

HEAD PER DAY (LBS.).

Experimental Food—Clover-silage.

Average milk per head per day.											Summary for experimental period. 13 weeks, Dec. 14—Mar. 14.		
Experimental period.													
Week ending											Average per day.	Total yield.	
Nov. 10.	Jan. 17.	Jan. 24.	Jan. 31.	Feb. 7.	Feb. 14.	Feb. 21.	Feb. 28.	Mar. 7.	Mar. 14.				
no. of cows.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs.	galls.
5 2	36 13	35 6	36 9	36 5	35 4	34 10	33 7	33 4	31 11	35 15	3271½	317	
2 8	35 4	37 1	39 11	38 6	38 1	36 2	36 6	35 13	33 3	28 14	2624½	254	
6 6	16 7	15 11	15 11	15 9	15 14	15 8	15 12	15 12	15 4	16 1	1461½	142	
5 5	19 11	19 6	19 8	19 7	19 0	18 8	17 13	15 15	15 8	18 13	1712½	166	
8 8	14 1	13 8	12 12	12 11	13 7	17 15	32 13	33 4	33 4	19 11	1794	174	
12 12	21 8	21 11	21 0	19 12	19 15	19 15	19 5	18 13	18 1	21 3	1926½	186	
8 8	15 5	14 13	14 2	13 15	13 7	12 15	12 9	11 14	11 4	14 2	1287½	125	
9 9	33 5	33 8	33 3	33 8	32 0	31 7	31 0	30 6	29 13	32 10	2970½	287	
3 3	21 13	20 11	19 14	18 15	18 1	16 15	15 15	14 13	12 12	19 4	1750½	169	
14 14	33 9	32 11	32 13	32 0	31 12	31 2	31 6	31 6	31 2	33 3	3017½	292	
18 18	17 11	16 15	17 10	18 5	16 15	15 5	13 10	14 15	14 2	17 6	1579½	153	
4 4	31 6	29 8	27 12	28 8	27 11	28 9	26 11	26 14	25 2	29 3	2653½	257	
11 11	21 3	20 15	20 5	20 10	20 3	20 4	20 1	19 14	19 7	20 9	1871	181	
10 10	44 5	44 0	44 15	45 10	46 2	46 13	44 1	44 10	43 12	43 11	3978½	385	
7 7	12 13	12 2	12 12	13 9	13 4	13 10	14 11	20 6	21 4	14 11	1335½	129	
15 15	19 2	19 6	18 10	19 4	19 8	19 10	18 13	18 15	17 8	18 11	1701½	165	
6 6	32 11	32 7	33 15	32 12	32 9	33 10	32 12	33 0	32 13	32 8	2956	286	
4 4	34 10	35 1	34 0	31 15	30 4	30 15	28 9	24 11	24 11	28 6	2580½	250	
8 8	43 10	44 7	43 13	43 3	44 1	44 12	44 6	42 14	42 10	42 8	3865½	374	
7 13	28 13	28 6	29 9	28 14	28 11	28 2	28 7	27 0	25 5	27 12	2526½	245	
5 2	26 11	26 6	26 7	26 3	25 13	26 4	26 0	25 12	24 15	25 12	2343½	227	

§ Replaced February 25th, by Daffodil, calved February 20th.

|| Slipped calf October 15th, but continued milking.

¶ Replaced December 23rd, by Kate, calved December 20th.

EXPERIMENT

TABLE XIII.—YIELD

Lot 2.—20 Cc

Names of cows.	Date of calving.	First milked.	Average milk per head per day.								
			Prior to experiment.						Experimental period.		
			From calving.	Previous 10 weeks (or less).	Last week.	Week ending					
						Dec. 20.		Dec. 27.		Jan. 3.	
	1884.	1884.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	
21. Susan*	June 12	June 14	28 5	18 4	11 7	10 1	8 7	5 9	40 2	39 4	
22. Countess	Nov. 23	Nov. 28	45 8	(45 7)	46 1	44 10	41 10	40 2	39 4	39 4	
23. Nina	Oct. 31	Nov. 8	37 0	(37 0)	34 10	32 5	31 1	30 4	29 15	29 15	
24. Jane	Nov. 29	Nov. 30	23 15	(24 0)	27 5	29 9	28 8	27 15	26 8	26 8	
25. Charmer	Aug. 30	Sept. 3	36 8	33 2	26 1	24 12	23 15	22 8	21 8	21 8	
26. Nancy†	Sept. 9	Sept. 10	30 15	28 5	21 10	13 15	13 3	12 8	11 8	11 8	
27. Nelly	Dec. 2	Dec. 6	39 12	(41 0)	41 0	40 15	39 1	38 0	37 0	37 0	
28. Princess‡	April 29	May 5	22 8	17 5	13 1	12 7	10 9	9 2	8 2	8 2	
29. Bright Eye	June 17	June 23	32 10	28 0	25 15	25 4	23 15	24 2	23 15	23 15	
30. Darling	July 5	July 5	34 12	29 6	23 8	21 13	21 0	20 2	19 15	19 15	
31. Empress	Oct. 18	Oct. 21	40 6	(40 12)	37 0	38 12	38 3	36 15	35 15	35 15	
32. Julia	Nov. 16	Nov. 17	27 0	(29 4)	34 10	30 15	32 13	32 15	31 15	31 15	
33. Scrutable	Nov. 3	Nov. 5	35 15	(36 6)	33 9	31 3	29 15	28 15	27 15	27 15	
34. May Day	July 4	July 5	27 4	22 7	21 11	22 7	23 6	22 8	21 8	21 8	
35. Liddy	Dec. 11	Dec. 14	—	—	—	32 12	43 0	50 15	49 15	49 15	
36. Welcome	Nov. 24	Nov. 28	45 12	(46 3)	46 5	47 13	47 3	44 15	43 15	43 15	
37. Ne plus ultra§	July 1	July 5	30 15	23 6	20 5	20 6	21 1	20 15	19 15	19 15	
38. Cowslip	April 1	April 4	31 9	21 4	15 13	18 4	15 14	14 15	13 15	13 15	
39. Russet Belle	March 18	March 20	38 9	27 9	21 0	19 14	19 4	19 15	18 15	18 15	
40. Ophir¶	July 26	August 1	34 1	28 6	20 3	19 3	17 10	15 4	14 15	14 15	
Average			33 14	30 6	27 7	26 14	26 7	25 15	24 15	24 15	

* Replaced January 11th, by Dewdrop, calved January 6th.

† Bought as newly-calved at this date; exact date of calving not known. Replaced February 22nd, by Gypsey, calved January 30th. (See Note ‡, Lot 1).

‡ Replaced January 24th, by Emma, calved January 2nd.

H COWS.

PER HEAD PER DAY (LBS.).

Experimental Food—Mangels.

Average milk per head per day.										Summary for ex- perimental period. 13 weeks, Dec. 14—Mar. 14.			
Experimental period.													
Week ending										Aver- age per day.	Total yield.		
Jan. 17.	Jan. 24.	Jan. 31.	Feb. 7.	Feb. 14.	Feb. 21.	Feb. 28.	Mar. 7.	Mar. 14.					
lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs. oz.	lbs.	galls.	
8 51 10	48 14	51 0	53 8	54 7	52 4	51 0	49 9	47 12	37 11	3432½	332		
15 38 7	40 10	40 2	39 12	39 3	37 4	38 14	38 2	38 12	39 10	3608½	349		
7 31 10	30 14	31 8	31 6	31 1	30 8	29 15	28 10	27 15	30 10	2789	270		
10 31 11	30 12	30 8	31 3	31 2	31 0	30 15	29 13	28 12	30 5	2759½	267		
4 21 12	20 11	20 7	20 12	20 3	21 1	20 3	19 3	19 8	21 6	1944½	188		
13 11 4	10 9	10 9	10 8	10 0	9 0	17 14	20 14	20 5	13 6	1216½	118		
7 38 2	33 11	39 8	34 15	34 8	34 5	34 14	33 6	34 7	36 6	3298½	319		
0 4 0	{ 3 4 7 6 }	12 0	18 15	20 15	20 7	21 3	20 11	20 9	14 1	1279½	124		
8 23 13	24 4	23 0	23 6	22 14	22 0	22 15	21 11	21 8	23 4	2115½	205		
12 20 7	20 15	19 15	21 1	21 0	20 12	26 6	29 0	29 10	22 9	2054½	199		
6 34 4	32 12	32 10	34 3	33 4	32 5	33 12	33 12	32 1	34 4	3116½	302		
4 30 5	29 5	27 14	26 13	25 7	24 10	25 9	27 10	27 11	28 7	2589	251		
14 26 0	25 5	23 11	24 10	22 12	22 12	22 4	21 13	21 14	24 14	2266½	219		
4 21 4	21 4	19 8	19 11	20 5	19 14	19 0	18 1	17 6	20 7	1860½	180		
14 49 5	44 5	48 12	49 15	44 7	45 6	47 11	47 13	49 1	46 3	4201	407		
9 40 13	39 10	35 10	35 12	35 14	34 2	32 12	30 0	29 11	37 13	3443½	333		
1 17 7	15 7	12 5	10 14	34 5	36 8	35 10	37 0	34 11	24 3	2202½	213		
14 6 0	{ 5 8 11 0 }	20 5	23 10	23 7	22 12	21 13	21 9	20 10	17 7	1589½	154		
14 16 14	16 13	16 10	15 15	16 4	16 3	14 15	15 10	15 14	17 1	1551½	150		
14 13 8	12 4	11 3	{ 9 14 29 11 }	43 12	44 12	47 6	47 0	48 3	26 14	2444½	237		
5 26 7	25 7	26 6	27 2	29 4	28 14	29 12	29 9	29 5	27 5	2488½	241		

§ Replaced February 8th, by Butterfly, calved February 3rd.

|| Replaced January 24th, by Rose, calved January 1st.

¶ Replaced February 6th, by Milkmaid, calved February 4th.

We will next give the results obtained on changing from clover-silage to meadow-grass-silage:—

On March 14 the experiment with cows on clover-silage terminated, and that on meadow-grass-silage at once commenced, and was continued for a period of six weeks. In order not to interfere with the yield of milk by a sudden change of food, a mixture of three parts clover-silage and one part grass-silage was given during the first seven days, and a mixture of half clover- and half grass-silage during the next seven days, after which grass-silage only was given.

Table XIV shows the average amounts of the different foods consumed per head per day, within each week, by each of the two lots of 20 cows.

EXPERIMENTS WITH COWS.

TABLE XIV.—AVERAGE FOOD CONSUMED PER HEAD PER DAY.
SECOND PERIOD; SIX WEEKS, MARCH 15 TO APRIL 25.

Week ending	Average food given per head per day.					Total food weighed off.
	Cake.	Bran.	Chaff.*	Silage.	Mangels.†	
Lot 1.—20 Cows ; Part Clover-silage, part Meadow-grass-silage.						
1885.	lbs.	lbs.	lbs.	lbs.	lbs.	lbs.
March 21	4	4	9½	50	(1) 11½	408
„ 28	4	4	8	50	(1) 20	197
April 4	4	4	7½	50	(2) 20	329
„ 11	4	4	7	50	(4) 11½	395
„ 18	4	4	7	42½	(20) 6½	434
„ 25	4	4	7	41½	(20) 17½	75½
Average per head } per day	2.19
Lot 2.—20 Cows ; Mangels as before.						
March 21	4	4	10	—	90	—
„ 28	4	4	10	—	90	—
April 4	4	4	10	—	90	—
„ 11	4	4	10	—	90	40
„ 18	4	4	10	—	90	—
„ 25	4	4	10	—	90	—
Average per head } per day	0.06

* ½ hay and ½ oat straw.

† The figures in parentheses show how many cows received mangels.

‡ During this period most of the unconsumed food was distributed among the other cows of the lot, and consumed.

It will be seen that both lots received the same amount of cotton-seed cake and bran as before; and Lot 2 also the same amount of chaff and mangels as before. In fact, there was no change whatever in the food of the 20 cows receiving mangels. The cows receiving silage were also at first supplied with the same amounts of both chaff and silage as before; but they did not consume them, and it was found necessary to reduce the quantity of chaff, until, during the last three weeks of the period, the allowance was only 7 lbs. instead of 10 lbs. per head per day; and during the last fortnight the amount of silage was reduced by nearly one-fifth, and some mangels were given to all instead of only to a few, of the cows receiving silage. It will be observed, indeed, that throughout the experiment a good deal of food, chiefly silage and chaff, was left unconsumed.

In the case of the experiment with clover-silage as against mangels, it was concluded that the silage, when given to cows in quantity containing an amount of dry matter corresponding to that in the mangels, was not so suitable for milk production as the succulent mangels, which only contained about half the percentage of dry matter, and a very much less proportion of woody fibre. It was also concluded that probably a better result would have been obtained if a smaller quantity of the silage, with some mangels, had been given. The meadow-grass-silage contained a still higher percentage of dry matter than the clover-silage, and no doubt a much higher percentage of woody fibre. It also contained only about two-thirds as much nitrogen as the clover-silage; though the quantity of it consumed contributed nearly as much nitrogen as the quantity of mangels consumed by the other cows; and doubtless the silage supplied a larger amount of albuminoid nitrogen than the mangels, that is, a larger amount in a condition available as food.

It is not to be wondered at, therefore, that with silage containing still more dry matter, and especially still more woody fibre, the cows could not consume so much, and that they should also require much less chaff.

Table XV (p. 48) shows, for each of the 20 cows receiving the meadow-grass-silage, the average yield of milk per day, during each of *x* weeks of the experiment; and Table XVI (p. 49) shows the same particulars for the 20 cows receiving mangels. In each case, there is also given the average yield per day during the preceding 13 weeks of those of the cows which had gone through the whole of the previous experiment, or for a shorter period for those which had been brought in during its progress; also the average yield during the immediately preceding week. Finally, there is given the average yield per day, and the aggregate yield (both in lbs. and in gallons), of each cow, and of each lot, during the whole period of six weeks of the second experiment.

EXPERIMENTS WITH COWS.

TABLE XV.—YIELD OF MILK PER HEAD PER DAY, &c. SECOND PERIOD; MARCH 15 TO APRIL 25.
 Lot 1.—20 Cows; part Clover-silage and part Meadow-grass-silage 2 weeks; Meadow-grass-silage 4 weeks.

Names of cows.	Yield of milk per head per day.														6 weeks, March 15 to April 25.	
	Previous period.				Experimental period.										Average per day.	Total yield.
	13 weeks (or less) Dec. 14 to March 14.				Week ending											
	lbs.	oz.	lbs.	oz.	March 21.	March 28.	April 4.	April 11.	April 18.	April 25.	lbs.	oz.				
1. Strawberry	35	15	31	11	30	15	27	7	24	10	23	2	24	11	118½	108
2. Ann	(36	10)	33	3	33	4	32	6	31	6	31	7	31	7	1847½	130
3. Isabella*	16	1	15	4	14	1	12	4	10	2	9	9	42	2	969	94
4. Penderpet	18	13	15	8	14	4	12	3	9	10	20	13	22	2	686½	62
5. Rosemary.	(32	5)	33	4	33	10	32	12	31	3	28	10	27	9	1257½	122
6. Joan of Arc.	21:	3	18	1	17	2	16	8	14	9	14	6	13	11	622½	60
7. Beatrice†	32	10	29	13	35	1	33	3	35	12	29	14	34	8	1464½	142
8. Parody	19	4	12	12	29	5	27	14	27	11	28	5	26	2	1153½	112
9. Queen Bees‡	33	3	31	2	10	15	8	8	31	9	35	7	35	2	1118½	108
10. Cherry Bipe	17	6	14	2	39	11	29	12	23	11	27	6	26	12	1177½	114
11. Granny‡	29	3	25	2	13	9	12	8	11	2	9	7	7	12	724½	70
12. Helen	30	9	19	7	25	2	25	8	24	8	21	11	18	2	942½	91
13. Polly...	43	11	43	12	41	12	41	7	41	3	37	13	35	7	739	72
14. Sally	(19	10)	21	4	22	8	23	1	22	7	21	8	21	7	1644½	159
15. Daffodil	18	11	17	8	16	15	15	10	15	11	15	4	15	3	925½	90
16. Fairy Queen	32	8	32	13	33	6	32	5	32	2	32	6	26	3	635½	63
17. Buttercup	(20	13)	24	11	42	10	45	14	20	8	18	14	13	7	1238½	122
18. Kate	42	8	42	10	22	12	45	9	45	11	37	10	34	13	861	82
19. Lucy	27	12	25	5	24	4	22	10	21	7	20	1	19	2	1719½	166
20. Victoria	27	13	25	10	25	10	25	2	25	1	23	15	25	9	880½	85
Average ..	27	13	25	10	25	10	25	2	25	1	23	15	25	9	1060½	102

* Replaced April 10, by Princess, calved April 6.

† Replaced April 10, by Alice, calved March 28.

‡ Beatrice calved March 10, brought into the experiment on March 16.

§ Replaced March 28, by Delance, calved March 19.

¶ Replaced April 16, by Caroline, calved April 10.

TABLE XVI.—YIELD OF MILK PER HEAD PER DAY, &c. SECOND PERIOD; MARCH 15 TO APRIL 25.
Lot 2.—20 Cows; Mangels as before.

Names of cows.	Yield of milk per head per day.										6 weeks, March 15 to April 25.	
	Previous period.			Experimental period.							Average per day.	Total yield.
	13 weeks (or less) Dec. 14 to March 14.			Last week, March 8 to 14.				Week ending				
	lbs.	oz.	lbs. oz.	March 21.	March 23.	April 4.	April 11.	April 18.	April 25.	lbs. oz.		
21. Dewdrop	47	12	45 6	44 9	44 11	39 10	39 11	43 13	42 15	1803 7	170	
22. Countess	39	10	37 7	38 8	37 9	36 12	36 2	36 12	37 3	1661 4	161	
23. Nina	30	10	27 15	28 14	26 15	23 4	23 8	25 10	25 9	1074 4	104	
24. Jane	30	6	29 0	28 1	27 1	27 5	26 12	26 6	27 7	1151 1	112	
25. Charnier	21	6	19 8	18 14	16 6	15 0	14 1	13 3	15 8	651 1	63	
26. Gypsy	(19 11)	20 5	19 14	20 0	20 10	21 5	20 15	21 1	20 10	866 4	84	
27. Nelly	36 6	34 7	32 6	29 4	29 9	27 7	27 3	28 6	29 0	1218 4	118	
28. Emma	(19 0)	20 9	20 7	20 9	20 14	23 2	20 1	20 6	20 9	863 4	84	
29. Brighteye*	23 4	21 8	21 7	20 10	19 13	17 12	14 7	{ 8 14 4 }	16 3	763 1	74	
30. Darling	22 9	29 10	28 0	28 5	25 13	27 4	27 13	27 10	27 8	1163 4	112	
31. Empress	34 4	32 1	32 3	32 13	31 15	32 0	33 0	31 2	32 3	1451 1	131	
32. Juliet	28 7	27 11	25 13	24 9	24 12	18 6	16 1	33 1	23 10	795 1	77	
33. Scrutable	24 14	21 14	20 12	20 10	19 12	19 10	19 8	19 7	19 15	837 1	81	
34. Mayday†	20 7	17 6	16 0	15 7	15 10	13 13	{ 11 10 1 }	38 14	19 4	575 2	55	
35. Liddy	46 3	49 1	46 0	45 0	45 6	44 7	45 7	44 6	45 2	1893 3	183	
36. Welcome	37 13	29 31	28 3	28 3	27 5	27 13	27 7	27 4	27 11	1163 4	112	
37. Butterfly	(25 10)	34 11	35 10	35 13	35 10	32 9	34 1	34 0	34 10	1453 1	140	
38. Rose	(21 13)	20 10	19 15	19 6	19 4	18 11	18 10	19 1	19 3	804 1	78	
39. Russet Belle	17 1	15 14	14 9	14 11	15 10	15 12	14 15	16 4	15 2	635 1	62	
40. Milkmaid	(45 5)	48 3	48 1	43 1	43 0	38 11	39 8	39 7	41 15	1762 1	171	
Average	30 4	29 5	28 5	27 11	27 6	25 15	25 12	27 15	27 3	1119 1	108	

* Replaced April 24, 1885, by Sylvia, calved April 10, 1885.

† Replaced April 19, 1885, by Flora, calved April 18, 1885.

‡ Replaced April 18, 1885, by Phoebe, calved April 10, 1885.

In the first place, Table XV (p. 48) shows that, of the 20 cows on silage, four became dry during the experiment; one in the second, two in the fourth, and one in the fifth week, and they were replaced by others; three out of the four of which gave a higher yield of milk than the average of the lot. Table XVI (p. 49) shows that, of the 20 cows on mangels, three became dry, one in the fifth, and two in the sixth week; and they were in each case replaced by others which yielded more than the average of the lot.

In the week prior to the commencement of this second experiment, the average yield of the silage-fed cows was nearly $3\frac{1}{2}$ lbs. per head per day less than that of the cows on mangels; and although during the six weeks of the experiment the cows receiving the meadow-grass-silage gave a less actual quantity of milk than those on mangels, they fell off in yield less, and gave, on the average of the whole period, only 2 lbs. instead of as at the beginning nearly $3\frac{1}{2}$ lbs. less milk per head per day than the mangel-fed cows. Both lots fell off in yield during the fourth week, probably under the influence of the weather. But whilst the silage-fed cows recovered again in the next week, coincidently with a reduction in the amount of silage and its replacement by some mangels, aided by the introduction of fresh cows, those on mangels did not recover until a week later, and then not to their original yield, notwithstanding three high-yielding cows had been brought in.

So far as the yield of milk is concerned, the evidence is, therefore, not adverse to the silage, when used in moderate quantity, according to the amount of solid and digestible matter it contains.

In the first experiment, the cows receiving clover-silage in rather large amount gave rather less milk than those on mangels; but, on the other hand, they gave a considerable increase in live-weight, whilst those on mangels as a whole lost in weight. Owing to the numerous changes in the cows during the course of the experiments, it happens that there are only 12 of the original silage-fed cows, and only 11 of the original mangel-fed cows, the weights of which can be traced through both periods of experiment. The cows were weighed soon after the commencement of the experiment with clover-silage, in December, 1884; again at the end of that experiment, when that with the meadow-grass-silage commenced; and, lastly, at the conclusion of the second period. We do not propose to enter fully into the subject of the increase or loss in weight of the animals on the present occasion, but only to refer to the general character of the results in their connection with the greater or less yield of milk by the different lots, over the different periods.

As already said, the cows on clover-silage increased in weight con-

rably, and of the 12 of the original 20 which continued through both experiments, every one increased when on the clover-silage. On the other hand, of the 11 of the mangel-fed cows which remained throughout the two periods, only 4 increased during the first period, and on the average the lot lost in weight. During the second period, however, when the meadow-grass-silage was used, only 3 of the 12 increased at all, two of them very little, and the lot as a whole lost in weight considerably; whilst of the 11 cows of the mangel lot, 9 increased, and the lot as a whole increased in weight.

From the whole of the results of the experiments on the feeding of cows with silage, it would appear that the clover-silage given in such a quantity was more favourable for meat-production, and less for milk-production, than the mangels. On the other hand, when the meadow-grass-silage was used, the animals did not give more milk, but they lost in weight. There can be no doubt that the grass-silage is inferior as food to the clover-silage; and it is to be supposed that the unfavourable maintenance in the yield of milk on the grass-silage compared with that of the cows on mangels over the same period, is in any rate partly to be attributed to a drain on the flesh and fat previously stored up by the animals. So far as this may take place, it obviously does not necessarily follow that there will be an immediate drop off in the yield of milk on changing to an inferior food. It is known that a badly-fed cow will reduce in condition very much when milking a calf; and the same thing may happen when the cow is milked. In fact, the comparative values of different foods for cows must not be measured by the yield of milk alone; but the increase or decrease in weight must also be taken into account.

—CHARACTER AND COMPOSITION OF THE MILK FROM THE DIFFERENT FOODS.

owing to the large amount of additional work in the Rothamsted Laboratory in connection with other branches of the ensilage experiments, it was found impossible to devote sufficient time and attention to obtain a complete series of comparative analyses of the milk yielded by the cows on the different foods. Still, sufficient has been done to indicate the comparative characters of the different milks. Table XVII (p. 52) shows the percentage of total solids, in 13 samples of the morning milk, of each of the two lots of cows, taken at intervals during the 10 weeks from January 14 to March 25; Lot 1 on lucerne silage, and Lot 2 mangels containing a corresponding amount of dry matter; each lot having the same amount of other foods in addition. The table also shows the percentages of mineral matter, or ash, in the last eight samples for each lot.

TABLE XVII.—PERCENTAGES OF TOTAL SOLIDS, AND MINERAL MATTER (ASH), IN THE MORNING MILK OF THE COWS.

Lot 1.—*Silage.* Lot 2.—*Mangels.*

Dates.	Total solids.		Mineral matter (ash).	
	Lot 1. Silage.	Lot 2. Mangels.	Lot 1. Silage.	Lot 2. Mangels.
A.M.	Per cent.	Per cent.	Per cent.	Per cent.
January 14	12·13	12·36	—	—
" 19	12·01	12·12	—	—
" 21	11·96	12·35	—	—
" 26	11·95	12·05	—	—
" 28	11·80	12·30	—	—
February 9	11·86	12·19	0·75	0·78
" 11	11·87	12·15	0·79	0·77
" 23	11·92	12·15	0·61	0·65
" 25	11·77	12·12	0·72	0·73
March 9	11·94	12·21	0·71	0·73
" 11	11·91	12·36	0·71	0·73
" 23	11·93	12·52	0·73	0·73
" 25	12·02	12·68	0·71	0·73
Mean	11·93	12·27	0·72	0·73

Table XVIII (p. 53) shows, not only the percentages of total solids and of mineral matter, but also the percentages of butter-fat, and the specific gravity, of both the *morning* and the *evening* milk of each lot of cows, on six different days from April 7 to April 22. It may be stated that the percentage of fat was frequently determined by Marchand's lacto-butyrometer; but the results given in the table are calculated by Fleischmann and Morgen's formula, from the amount of total solids, and the specific gravity. The results obtained by the lacto-butyrometer, although very useful, are admittedly only approximative; and, as Dr. P. Vieth has shown by a very comprehensive investigation of the subject, the deviations from the amounts as determined by more exact methods, are greater in the case of results obtained by it, than in those obtained by calculation from the total solids and the specific gravity.

Referring to the results in Table XVII, it should be observed that Lot 1 were receiving clover-silage during the periods of sampling from January 14 to March 11, and a mixture of clover-silage and meadow-grass-silage when the samples of March 23 and March 25 were taken. They were also receiving meadow-grass-silage at the six periods of sampling to which the results given in Table XVIII refer. Lot 2, however, received mangels throughout.

TABLE XVIII.—COMPOSITION OF THE MORNING AND THE EVENING MILK OF THE COWS.
 Lot 1.—*Meadow-grass-silage.* Lot 2.—*Mangels.*

Dates.	Average yield of milk per head.		Specific gravity of the milk (corrected to 60° F.).			Total solids.			Butter-fat.			Mineral matter (ash).	
	A.M.	P.M.	Total.	A.M.	P.M.	Direct daily average.	A.M.	P.M.	Direct daily average.	A.M.	P.M.	A.M.	P.M.

Lot 1.— <i>Meadow-grass-silage.</i>													
April 7	14 4	10 0	24 4	Sp. gr. 1·0833	Sp. gr. 1·0821	Sp. gr. 1·0828	P. cent. 12·29	P. cent. 12·67	P. cent. 12·45	P. cent. 3·03	P. cent. 3·61	P. cent. 0·71	P. cent. 0·71
" 9	13 6	9 2	22 8	1·0831	1·0828	1·0830	12·41	13·06	12·67	3·18	3·78	0·73	0·72
" 13	14 6	10 0	24 6	1·0831	1·0820	1·0826	12·18	12·68	12·39	2·99	3·64	0·70	0·71
" 15	15 9	10 12	26 5	1·0834	1·0825	1·0830	11·97	12·79	12·31	2·74	3·62	0·72	0·73
" 20	15 6	10 11	26 1	1·0822	1·0824	1·0823	11·82	12·79	12·22	2·86	3·64	0·64	0·67
" 22	15 15	10 10	26 9	1·0825	1·0823	1·0824	12·09	12·64	12·31	3·02	3·64	0·71	0·71
Mean ..	14 13	10 3	25 0	1·0829	1·0824	1·0827	12·13	12·77	12·39	2·97	3·64	0·70	0·71

Lot 2.— <i>Mangels.</i>													
April 7	14 13	11 1	25 14	1·0835	1·0839	1·0837	12·99	13·54	13·23	3·58	3·96	0·75	0·75
" 9	15 1	10 12	25 13	1·0844	1·0838	1·0841	12·85	13·43	13·09	3·26	3·89	0·73	0·74
" 13	14 11	10 13	25 8	1·0845	1·0839	1·0842	12·54	13·40	12·90	2·99	3·84	0·73	0·73
" 15	15 0	10 11	25 11	1·0844	1·0843	1·0844	12·64	13·49	12·99	3·08	3·82	0·74	0·75
" 20	16 5	11 12	28 1	1·0840	1·0839	1·0839	12·14	13·53	12·72	2·74	3·95	0·75	0·69
" 22	16 0	11 12	27 12	1·0841	1·0840	1·0841	12·73	13·39	13·01	3·22	3·81	0·74	0·74
Mean ..	15 5	11 2	26 7	1·0841	1·0840	1·0841	12·65	13·46	12·99	3·15	3·88	0·72	0·73

It is seen that the mean percentage of total solids in the 13 samples of the morning milk of the silage-fed cows, taken from January 14 to March 25, is 11·93 against 12·13, the mean of the six samples of their morning milk taken from April 7 to April 22. Again, the mean of the 13 samples of the morning milk of the mangel-fed cows, taken from January 14 to March 25, was 12·27, against 12·65 during the period from April 7 to April 22. There was thus, in both cases, an increase in the percentage of total solids as the season advanced. But, as the increase was even greater with the mangel- than with the silage-fed cows, it is obvious that the result is due to the season, and not, in the case of Lot 1, to the change from clover- to meadow-grass-silage.

Still confining attention to the results relating to the morning milk, which is well known to be poorer than evening milk, it is seen that at each of the 19 periods of sampling to which Tables XVII and XVIII refer, the milk of the mangel-fed cows shows a higher percentage of total solids than that of the silage-fed cows. It also shows, almost without exception, a slightly higher percentage of mineral matter.

Turning now to the more detailed composition, as recorded in Table XVIII, it is seen that the milk of both lots shows, on the average, indeed in nearly every individual case, a slightly lower specific gravity in the afternoon than in the morning. The afternoon milk shows, on the other hand, uniformly a considerably higher percentage of both total solids and butter-fat; and generally a somewhat higher percentage of mineral matter also. Thus, whilst the mean percentage of total solids in the morning milk of the silage-fed cows is 12·13, in their evening milk it is 12·77, or 0·64 higher; and whilst the mean percentage of total solids is 12·65 in the morning milk of the mangel-fed cows, it is 13·46, or 0·81 higher, in their evening milk. Again, whilst the mean percentage of butter-fat in the morning milk of the silage-fed cows is 2·97, in their evening milk it is 3·64, or 0·67 higher; and whilst the mean percentage of butter-fat in the morning milk of the mangel-fed cows is 3·15, in their evening milk it is 3·88, or 0·73 higher. It is thus seen that, in the case of both lots of cows, the higher percentage of total solids in the evening milk very closely corresponds with the increased amount of butter-fat in the evening milk.

The first three columns of Table XVIII, show the average yield of milk per head of each lot of cows, in the morning, in the evening, and for the whole day, at each of the six periods of the sampling and the analysis of the milk. It will be seen that there was little more than two-thirds as much of the richer evening milk as of the poorer morning milk.

The "*direct daily averages*," of specific gravity, and of the percentages of total solids, butter-fat, and mineral matter, as given in the

third column of the respective divisions of the table, are calculated from the quantities, and the composition, of the morning and the evening milk, respectively. According to these results, the average percentage of total solids in the total daily milk of the silage-fed cows was, over the six periods, 12·39, against 12·99, or 0·60 higher, in that of the mangel-fed cows. And the average percentage of butter-fat in the daily milk of the silage-fed cows is 3·24, against 3·45, or 0·21 higher, in the milk of the mangel-fed cows. There is thus less difference between the average amount of total solids in the milk of the two lots of cows, than there is between the morning and evening milk of the same cows; and there is very much less difference in the average amount of butter-fat in the milk of the two lots, than there is between the amount in the morning and the evening milk of the same cows. It may be added that the average percentages of total solids, and of butter-fat, agree very fairly with those given by Dr. Vieth for the average milk of the same period of the year; those of the silage-fed cows being somewhat below, and those of the mangel-fed cows above his average.

Upon the whole, the analytical results clearly show that the milk of the mangel-fed cows throughout contained higher amounts of both total solids, and butter-fat, than that of the silage-fed cows. Yet, quite consistently with the observation of others on the same point, the milk of our silage-fed cows was judged, both by colour and by taste, to be richer than that of the mangel-fed cows. The milk of the silage-fed cows possessed a slight, but not at all disagreeable, flavour, which may be described as *hayey*, and which could readily be detected by some, but not by others. The butter from the milk of the silage-fed cows was also much yellower than that from the milk of the mangel-fed cows; but there was no perceptible distinction between the two as to taste.

With these results we close the record of our silage experiments for the present.

SUMMARY, AND GENERAL CONCLUSIONS.

1. It would require a larger area of land to obtain a given quantity of dry substance of food in crops grown for ensilaging, than to obtain the same quantity in roots.
2. The substitution of ensilage-crops for roots on a large scale would necessitate a considerable change in the course of cropping. It would lessen the area under cleaning crops, and consequently lessen the area suitable for growing grain for the market.
3. Where ensilaging is only adopted instead of hay-making, it is improbable that it will be substituted for it entirely, and if only

partially, the process would only have a comparatively limited application. Or, if it be extended to the natural and artificial grasses usually fed on the land, an expensive mode of feeding would be substituted for an economical one.

4. Neither in the case of red-clover, nor in that of meadow-grass, was the loss of dry substance of food in the silo so great as has frequently been supposed. It was, so far as can be judged, much about the same as in a hay rick.

5. There was some total loss of nitrogenous substance; but there was a much larger amount of it degraded from the albuminoid condition to compounds incapable of forming the nitrogenous substances of animal increase, or of milk.

6. The results obtained do not confirm the conclusion that woody fibre of a given degree of induration is rendered more soluble, and consequently more digestible.

7. In a comparative experiment with fattening oxen, a given amount of dry substance in red-clover-silage was found to be practically equal to the same amount of dry substance in a mixture of clover-hay-chaff and swedes, given in the proportion of 12 parts chaff and 50 parts swedes.

8. The amount of dry substance of food required by a cow of say about 1200 lbs. live-weight, when in milk, will vary considerably according to the character of the food, the temperature to which the animal is exposed, its yield of milk, and other circumstances. But it will seldom be less than 25 lbs. per head per day, and will seldom exceed 30 lbs.

9. When a milking cow is supplied with fair proportions of chaff and roots, with fair proportions of purchased food, such as cake and bran, in addition, the home-grown food will approximately supply the requirements of the animal for mere sustenance, and approximately the whole of the purchased food will remain for milk production. It is a question for consideration, therefore, how far the supply of the purchased foods should be graduated according to the yield of milk of each individual cow.

10. In ordinary liberal feeding, food is always consumed in excess of the amount which, according to calculation, is actually required for sustenance, and for the production of fattening increase, or of milk.

11. Two or three times as much total dry substance, and six, seven, or more times, as much nitrogenous substance, will be contained in the milk of a cow yielding 3 gallons per day, as in the fattening increase of an ox of the same weight.

12. In a comparative experiment with milking cows, clover-silage was tried against mangels. Two lots of cows, of 20 each, had the same amounts of cake, bran, and chaff; one lot receiving besides, an

average of nearly 49 lbs. of clover-silage, and the other lot an average of nearly 86 lbs. of mangels per head per day. Over a period of 13 weeks, the cows having clover-silage yielded an average of 25 lbs. 12 ozs. of milk per head per day, and those having mangels gave an average of 27 lbs. 5 ozs. per head per day.

13. There was more nitrogenous substance, and rather more total dry substance, but more of this was woody fibre, in the quantity of clover-silage consumed, than in the mangels. The silage was, therefore, when given in such large proportion, less suitable for milk-production than the more succulent mangels. The result was that the cows on the silage showed more tendency to fatten; and, in fact, though giving less milk, they gained in live-weight, whilst the mangel-fed cows slightly lost in weight.

14. It is probable that, if a portion of the clover-silage had been replaced by an amount of mangels containing a corresponding quantity of dry matter, the yield of milk would have been greater, and the tendency to increase in live-weight would have been less.

15. In a second experiment with the two lots of cows of 20 each, and extending over six weeks, the same standard foods were given as before; but, in addition, the one lot now received meadow-grass-silage, and the other mangels as before. Compared with the clover-silage, the meadow-grass-silage would contain still more woody fibre, and it contained little more than two-thirds as much nitrogenous substance. It was accordingly found that the animals would not consume so much chaff, nor the whole of the silage given to them. Some mangels were therefore given with the silage, when they consumed it better. Both lots of cows slightly declined in yield of milk during this period, the silage-fed cows rather less than the others. There was, however, but little difference in the yield of the two lots; but whilst the silage-fed cows now lost in weight, the mangel-fed cows slightly gained.

16. When analysed, the milk of the mangel-fed cows invariably showed a higher percentage of both total solid matter, and butter-fat, than that of the silage-fed cows.

17. The afternoon milk of both lots of cows always gave higher percentages of both total solid matter, and butter-fat, than the morning milk; and the excess of solid matter in the afternoon milk very closely corresponded with its excess of fat.

18. There was more difference in the percentage of both total solid matter and fat, between the morning and the evening milk of the same lot of cows, than between the milk of the two lots taken at the same time of day.

19. There can be no doubt that good food may be preserved in a favourable state for future use by being properly ensilaged. But the

results obtained at Rothamsted do not favour the idea that produce which is itself not good food, can be made good food by being ensilaged.

20. Good ensilage, given in such amount as to supply the same quantity of dry substance as would be given in chaff and roots, is no doubt a very good food for fattening oxen.

21. Good ensilage, given in less proportion, and in conjunction with roots, with the ordinary dry foods in addition, is no doubt a very good food for milking cows.

22. In conclusion, it is hoped that the details which have been given of the first year's experiments on ensilage at Rothamsted, will afford some useful basis for the consideration of those who may be deliberating whether or not to adopt the system.





